

Self-consistent intermediate Hamiltonians: A coupled cluster type formulation of the singles and doubles configuration interaction matrix dressing

I. Nebot-Gil and J. Sánchez-Marín

Departament de Química Física, Facultat de Química, Universitat de València, c/Dr. Moliner, 50 46100 Burjassot (Valencia), Spain

J. P. Malrieu, J. L. Heully, and D. Maynau

Laboratoire de Physique Quantique, URA 505 du CNRS, Université Paul Sabatier, 118, Route de Narbonne, 31062 Toulouse Cedex, France

(Received 27 January 1995; accepted 27 April 1995)

This paper presents a new self-consistent dressing of a singles and doubles configuration interaction matrix which insures size-consistency, separability into closed-shell subsystems if localized molecular orbitals (MOs) are used, and which includes *all* fourth order corrections. This method yields, among several schemes, a reformulation of the coupled cluster method, including fully the cluster operators of single and double excitations, and partially those of the triples (Bartlett's algorithm named CCSDT-1a). Further improvement can be easily included by adding exclusion principle violating corrections. Since it leads to a matrix diagonalization, the method behaves correctly in case of near degeneracies between the reference determinant and some doubles. Due to its flexibility this formulation offers the possibility of consistent combination with less expensive treatments for the study of very large systems. © 1995 American Institute of Physics.

I. INTRODUCTION

The size-consistency requirement is a fundamental aspect of the quantum many-body problem.¹⁻⁴ While truncated configuration interactions (CI) are variational and suitable for rational and flexible selections, they are not size-consistent.⁵ A more correct scheme is the coupled cluster (CC)⁶⁻¹² expansion. At its singles and doubles approximation (CC-SD), it ignores part of the triples and thus does not insure the Rayleigh-Schrödinger fourth order. The complete inclusion of the triples is rather expensive, and thus perturbative treatments of the triples are frequently proposed (CCSD(T)),¹³⁻¹⁵ although they do not behave properly when single bonds are broken. Although the full CCSDT model has been developed¹⁶⁻¹⁸ the calculations performed keep rather illustrative character due to the cost of the method. Moreover, since the CC equations are nonlinear, they must be solved iteratively. The numerical algorithms for solving sets of nonlinear equations suffer for risk of poor convergence and require rather large number of iterations.^{19,20}

Somewhere in between CI and CC one must mention the approximate size-extensive CEPA (coupled electron pair approximation)²¹⁻²⁶ and CPF (coupled pair function)^{27,28} models, which exist in several versions and are essentially based on a single reference and truncation to double excitations, plus an approximate cancellation of the unlinked corrections.

In a previous work,²⁹ the size extensivity of the lowest root of any selected CI has been obtained by a proper dressing (or change) of the diagonal energies of the CI matrix. This dressing is self-consistent, depending on the coefficients of the doubles, so that the method has been labeled (SC)²CI (size-consistent self-consistent CI). Its implementation is straightforward with negligible extra costs in terms of memory and computation time, and the efficiency is really

impressing,^{30,31} specially when a direct selected CI algorithm is used.³²

The method was derived in terms of the intermediate Hamiltonian Theory³³ (a generalization of effective Hamiltonians³³), but it may be seen as the most exact CEPA scheme when the model space is the SDCI space or as a generalized CEPA scheme for arbitrary CI's. It insures the strict separability of the energy

$$E_{AB} \rightarrow E_A + E_B \quad \text{when } r_{AB} \rightarrow \infty \quad (1)$$

for the separation of an *AB* system into *A* and *B* closed shell subsystems provided that localized MO's are used. The method consists in *adding unlinked* effects of the outer space onto the diagonal energies in order to cancel *all unlinked* terms produced by the diagonalization. Of course, if the model space consists in the SDCI space, the linked contributions of triples and quadruples are not taken into account so that the method is poorer than CC-SD which correctly treats the linked contribution of the quadruples.

The linked effects of the triples and quadruples have been added once³⁵ *a posteriori* as the mean value of an additional dressing operator, taken on the vector resulting from the (SC)²SDCI. Actually, this was an approximate application of a fundamental idea, namely the total dressing of the model space by the linked and unlinked effects of the outer space.

The preceding paper³⁶ has presented a simple definition of a diagonal dressing in terms of the coefficients of the desired vector on the outer space determinants, together with two perturbative evaluations of these coefficients (and therefore of the dressing). The test calculations were convincing, especially when high-order exclusion principle violating (EPV) corrections were included. The present paper explores the possibility to replace these perturbative evaluations of the

dressing by coupled cluster type estimates, which are expected to behave more satisfactorily when near degeneracies occur in the model space. A formulation is found, which is equivalent to CCSD method, if only the unlinked contributions to triples and quadruples are taken into account, or analogous to CCSDT-1a,^{37,38} if the linked contributions to Triples are perturbatively accounted for. Therefore, the method insures the strict separability into closed shell subsystems when localized MO's are used. Although leading to a matrix built on the SDCI space, the method treats all the fourth order effects of the triples and quadruples.

II. SELF-CONSISTENT INTERMEDIATE HAMILTONIAN WITH HERMITIAN DRESSINGS

A. Matrix formulation of the intermediate Hamiltonians

We shall not recall the definition and advantage of intermediate Hamiltonians.³³ We suppose that we only search the exact energy and the projection of the exact eigenvector onto the model space of a single root, this eigenvector having the largest amplitude on the *main* model space determinant Φ_0 (or reference determinant). The intermediate model space is spanned by (at least) all the determinants Φ_i interacting with Φ_0 . Thus, Φ_0 and all the Φ_i built the model space S . Then, if P is the projector on the model space S

$$P = |\Phi_0\rangle\langle\Phi_0| + \sum_{i \in S} |\Phi_i\rangle\langle\Phi_i|, \quad (2)$$

it leads to the diagonalization of the dressed matrix

$$\mathbf{P}(\mathbf{H} + \mathbf{\Delta})\mathbf{P}, \quad (3)$$

where $\mathbf{\Delta}$ is the dressing operator.

If we define $\mathbf{Q} = \mathbf{1} - \mathbf{P}$ as the projection on the outer space, and \mathbf{c} as the vector of the coefficients of the exact eigenvector:

$$\Psi_0 = \Phi_0 + \sum_{i \in S} c_i \Phi_i + \sum_{\alpha \notin S} c_\alpha \Phi_\alpha, \quad (4)$$

the exact Schrödinger equation $(\mathcal{H} - \mathcal{E}_0)\Psi_0 = 0$ may be written for the rows associated to the model space in a matrixial formulation:

$$\mathbf{P}\mathbf{H}\mathbf{P}\mathbf{c} - \mathcal{E}_0\mathbf{P}\mathbf{1}\mathbf{P}\mathbf{c} + \mathbf{P}\mathbf{H}\mathbf{Q}\mathbf{c} = 0. \quad (5)$$

The last term $\mathbf{P}\mathbf{H}\mathbf{Q}\mathbf{c}$ is a vector \mathbf{V} the elements of which are

$$V_i = \sum_{\alpha \notin S} c_\alpha \langle \Phi_i | \mathcal{H} | \Phi_\alpha \rangle \quad (6)$$

so that Eq. (5) may be written

$$\mathbf{P}\mathbf{H}\mathbf{P}\mathbf{c} - \mathcal{E}_0\mathbf{P}\mathbf{1}\mathbf{P}\mathbf{c} = -\mathbf{V}. \quad (7)$$

The dressed matrix eigen-equation is

$$[\mathbf{P}(\mathbf{H} + \mathbf{\Delta})\mathbf{P} - \mathcal{E}_0\mathbf{P}\mathbf{1}\mathbf{P}]\mathbf{P}\mathbf{c} = 0. \quad (8)$$

The condition under which the above equation will provide the exact energy and the exact components of the eigenvector on the model space is that

$$\mathbf{V} = \mathbf{P}\mathbf{\Delta}\mathbf{P}\mathbf{c} = \mathbf{\Delta}\mathbf{P}\mathbf{c} \quad (9)$$

or simply, restricted to the model space:

$$\mathbf{V} = \mathbf{\Delta}\mathbf{c}. \quad (10)$$

This equation is not sufficient to define $\mathbf{\Delta}$. Of course, $\mathbf{\Delta}$ might be a full square matrix as occurs in the partitioning technique.³⁹⁻⁴² However, it is sufficient to define the dressing through n matrix elements only since there are only n degrees of freedom (1 eigenenergy, $n-1$ coefficients).

B. Dressing formulations

One possibility is the diagonal dressing⁴³

$$\Delta_{ij} = \Delta_{ii} \delta_{ij}, \quad (11)$$

$$\Delta_{ii} = c_i^{-1} \sum_{\alpha \notin S} c_\alpha \langle \Phi_i | \mathcal{H} | \Phi_\alpha \rangle. \quad (12)$$

This solution has the drawback of the division by c_i which may lead to numerical instabilities and the possible appearance of physically meaningless eigenenergies of $\mathbf{P}(\mathbf{H} + \mathbf{\Delta})\mathbf{P}$ below the relevant root \mathcal{E}_0 .

Two of us have proposed elsewhere a first column dressing,⁴³ concerning the elements $\langle \Phi_i | \mathbf{\Delta} | \Phi_0 \rangle$

$$\langle \Phi_i | \mathbf{\Delta} | \Phi_0 \rangle = \sum_{\alpha \notin S} c_\alpha \langle \Phi_i | \mathcal{H} | \Phi_\alpha \rangle \quad (13)$$

which does not lead to such numerical troubles but which is non-hermitian.

We strongly recommend a new formulation of the dressing concerning the first column and first row only. The first column of $\mathbf{\Delta}$ is identical to the preceding except for $\langle \Phi_0 | \mathbf{\Delta} | \Phi_0 \rangle$, the dressing is made hermitian taking

$$\langle \Phi_0 | \mathbf{\Delta} | \Phi_i \rangle = \langle \Phi_i | \mathbf{\Delta} | \Phi_0 \rangle, \quad (14)$$

and Δ_{00} is then calculated as

$$\Delta_{00} = \langle \Phi_0 | \mathbf{\Delta} | \Phi_0 \rangle = \sum_{\alpha \notin S} c_\alpha \langle \Phi_0 | \mathcal{H} | \Phi_\alpha \rangle - \sum_{i \neq 0} c_i \langle \Phi_i | \mathbf{\Delta} | \Phi_0 \rangle. \quad (15)$$

These relations are easily obtained from the matrix multiplication of the first row of $\mathbf{\Delta}$ by \mathbf{c} .

In the case where the model space includes all the determinants interacting with Φ_0 the first summation is zero, and

$$\Delta_{00} = - \sum_{i \neq 0} c_i \langle \Phi_i | \mathbf{\Delta} | \Phi_0 \rangle. \quad (16)$$

This formulation is hermitian and has shown to be numerically stable.

C. Estimation of the coefficients of the outer space

All these formulations remain academic as far as the coefficients on the outer space are unknown. Of course, in practice they can be approximately evaluated from the knowledge of the c_i 's. Therefore these dressings require achieving a self-consistency condition. In the preceding paper³⁶ the coefficients c_α were evaluated perturbatively. This means that the coefficients for triple and quadruple excitations are obtained by a first order perturbation upon a vector built on all singly and doubly excited determinants.




FIG. 1. (a) Disconnected and (b) connected contributions to the coefficients of the quadruples.

The use of these c_α 's in Eq. (8) gives an energy which is correct to fourth order and includes important fifth order terms. The present work will try to improve such a scheme by systematically using the factorization theorem which is part of perturbation theory. This will transform the preceding order-by-order expansion into a self-consistent, all-order approach.

The following paragraphs will show how such an all-order scheme is obtained, first for the quadruple excitations. Then the triple excitations will be considered. A comparison with the coupled cluster approach will follow. Finally, we shall show how further improvements can be obtained by considering the nonadditivity of the denominators (this additivity is implicit in the Möller–Plesset perturbation theory) or by including higher-order terms in the CEPA spirit, but this time for triple and quadruple excitations. As far as we know this is the first time that EPV terms are included for excitations higher than single and double ones.

III. COUPLED CLUSTER FORMULATIONS OF SELF-CONSISTENT INTERMEDIATE HAMILTONIANS

A. Quadruple excitations

Up to third order in the wave function, two kinds of quadruple excitations have to be considered. At the second order disconnected quadruples appear, the connected ones appearing at the next order (see Fig. 1).

Regarding the energy, the connected quadruple amplitudes will contribute for the first time at fifth order, and furthermore the coefficient c_α^Q cannot be written as a function of the c_i 's. They have to be evaluated by perturbation, e.g., a

second order perturbation upon a singles and doubles vector. This is, of course, outside the scope of this method. The disconnected quadruples should be more important; their energy contribution appears at fourth order. By using the factorization theorem (see Ref. 44) one can show that the sum of the two terms of Fig. 1(a), can be rewritten as a product which by inspection yields the following second order relation

$$c_\alpha^Q = \sum_{(i,j)} c_i^{(1)} c_j^{(1)}, \quad (17)$$

where (i,j) stands for all couples of disjoint double excitations creating Φ_α from Φ_0

$$\Phi_\alpha = D_j^+ D_i^+ \Phi_0 = D_k^+ D_l^+ \Phi_0 = \dots \quad (18)$$

Equation (17), giving the coefficient of quadruples as products of double excitation coefficients, is generalized to an all-order relation by writing

$$c_\alpha^Q = \sum_{\substack{(i,j) \\ D_j^+ D_i^+ \Phi_0 = \Phi_\alpha}} c_i c_j, \quad (19)$$

where the c_i coefficients are obtained by diagonalization of the dressed Hamiltonian, giving thus a self-consistent iterative method.

From this relation it is very easy to derive approximate schemes such as “exact CEPA method,” called by us (SC)²MRCI and already presented in Ref. 29. These methods which considered EPV terms only will not be used here.




FIG. 2. (a) Connected and (b) disconnected contributions to the coefficients of the triples.

It is obvious that our scheme must have some relation with the coupled cluster method, that writes the amplitude of the quadruple excitations as product of double excitation amplitudes. A thorough comparison of both methods will be given later.

Comparison should also be done with our previous perturbative scheme.³⁶ Due to its perturbative character, one of the two double excitations forming the quadruple was described at all-order, whereas the other one was kept at first order only. We remedy, now, this unsymmetric way of describing quadruple excitations, and thus include further higher terms. It should be noted also that the fact that no denominators appear explicitly in this new scheme is certainly in favor of its convergence properties during the iterations.

B. Triple excitations

At variance with quadruple excitations, the connected amplitudes of the triples appear before the disconnected ones. These connected triples will contribute at fourth order to the energy whereas the disconnected ones at fifth order only (see Fig. 2).

The connected triples (T_c) coefficients cannot be rewritten in terms of the coefficients of the doubles and singles and thus have to be derived by perturbation upon a singly and doubly excited vector. They are single excitations of double-excitations.

$$c_{\alpha}^{T_c} = \frac{\sum_i' c_i^{(1)} \langle \Phi_{\alpha} | \mathcal{A} | D_i^+ \Phi_0 \rangle}{\Delta_{\alpha}} \quad (20)$$

where Σ' runs over all the double excitations for which there is no single excitation M_k^+ such that $M_k^+ D_i^+ \Phi_0 = \Phi_{\alpha}$ and Δ_{α} is an energy difference to be defined by the chosen perturbative scheme (MP, EN ...). The second order relation will be generalized to an all-order relation by writing

$$c_{\alpha}^{T_c} = \frac{\sum_i' c_i \langle \Phi_{\alpha} | \mathcal{A} | D_i^+ \Phi_0 \rangle}{\Delta_{\alpha}} \quad (21)$$

and we have again an iterative scheme which relates the double and triple excitations.

It should be noted that the generalization is perfectly well founded. If instead of a pure single reference perturbation theory we used a multireference method, the relation (21) would be obtained directly.

Comparison with coupled cluster will show, later, that this kind of Triple contributions is obtained only by CCSDT schemes or some of its approximations.

Concerning the comparison with our previous method there is no direct improvement in the treatment of the connected triples, however the double-excitation coefficients used in Eq. (21) are of better quality due to the improvement of the c_{α}^Q .

The disconnected triples (T_{dc}) part comes from the products of disjoint single and double excitations [see Fig. 2(b)]




FIG. 3. Fifth order energy diagram.

$$\Phi_{\alpha}^{T_{dc}} = M_i^+ D_j^+ \Phi_0 = M_k^+ D_l^+ \Phi_0 = \dots \quad (22)$$

Calling c_i' the coefficient of the singly excited determinant $\Phi_i = M_i^+ \Phi_0$, and using again the factorization theorem one can show that this sum can be rewritten as

$$c_{\alpha}^{T_{dc}} = \sum_{(i,j)} c_i^{(1)'} c_j^{(1)}, \quad (23)$$

where (i,j) stands here for the nine couples of single-double excitations such that

$$D_j^+ M_i^+ \Phi_0 = \Phi_{\alpha}. \quad (24)$$

This will be generalized to the obvious form

$$c_{\alpha}^{T_{dc}} = \sum_{(i,j)} c_i' c_j. \quad (25)$$

$$D_j^+ M_i^+ \Phi_0 = \Phi_{\alpha}$$

By including such terms in the dressing will include fifth order diagrams in the energy. One of them is shown in Fig. 3.

Such terms are automatically included if one uses natural or Brueckner orbitals instead of HF orbitals.^{45,46} Comparison with our previous method shows that the single excitation coefficient have changed from the first order to an all-order description. Since no denominators appear explicitly, better convergence properties can be expected during the iterations.

C. Comparison with the coupled cluster formalism

Let us expand Ψ_0 in Eq. (4) in a coupled cluster (CC) type way:

$$\Psi_0 = e^S \Phi_0. \quad (26)$$

If we take, for instance, the CCSD approximation, $S = T_1 + T_2$ and

$$\Psi_0 = \Phi_0 + (T_1 + T_2 + \frac{1}{2}T_1^2)\Phi_0 + (T_1T_2 + \frac{1}{6}T_1^3 + \frac{1}{2}T_1^2T_2 + \frac{1}{2}T_2^2 + \frac{1}{24}T_1^4)\Phi_0. \quad (27)$$

Written in this way, it is easy to see that each of the right hand terms in Eq. (27) has its counterpart in Eq. (4), for the case that the model space in Eq. (4) includes Φ_0 , all its Singles and all its Doubles.

Now, if we consider Ψ_0 in terms of the CI wave operator expansion and we group in the same way all excitations up to quadruples,

$$\Psi_0 = \Phi_0 + (C_1 + C_2)\Phi_0 + (C_3 + C_4)\Phi_0 \quad (28)$$

it follows that the procedure of taking into account the effects of the triples and quadruples on the SDCI matrix by a

proper dressing $\Delta_j = c_j^{-1} \sum_{\alpha \in S} c_{\alpha} h_{j\alpha}$ can be easily made equivalent to a full CCSD procedure. This is made clearer through the equivalence between the T_i and C_i operators

$$C_1 = T_1, \quad (29)$$

$$C_2 = T_2 + \frac{1}{2}T_1^2, \quad (30)$$

$$C_3 = \frac{1}{6}T_1^3 + T_1T_2, \quad (31)$$

$$C_4 = \frac{1}{24}T_1^4 + \frac{1}{2}T_2^2 + \frac{1}{2}T_1^2T_2. \quad (32)$$

Now taking T_1 and T_2 from Eqs. (29) and (30) and substituting them into Eqs. (31) and (32), we can write

$$C_3 = C_1C_2 - \frac{1}{3}C_1^3, \quad (33)$$

$$C_4 = \frac{1}{2}C_2^2 - \frac{1}{12}C_1^4, \quad (34)$$

i.e., we can rewrite the CCSD, originally based solely on T_1 and T_2 amplitudes, in terms of c_i model space wave function coefficients and c_{α} coefficients.

Comparison of Eqs. (33) and (34) with our $c_{\alpha}^{T_{dc}}$ and c_{α}^Q will reveal the close relation between both methods. The difference is given by third and fourth powers of C_1 operator. For the triple excitations these terms will contribute at 8th order of perturbation and for the quadruple ones at the 11th order! So the difference should be very tiny, indeed! These missing terms can, of course, be included easily in our scheme but this point is not obvious. Comparison with an all-order expansion will show that these terms are only some of the numerous terms appearing at those orders. There is no *a priori* way to say which scheme gives the best "arbitrary weight" to these powers of C_1 .

Anyway, in another work,⁴⁷ we have included these C_1 powers in order to show the strict equivalence between both schemes. Nevertheless, it remains a very important difference in the practicability. Our scheme yields a diagonalization of an effective Hamiltonian (containing no energy differences) whereas the CCSD scheme is the resolution of a system of equations (containing energy differences) of which convergence is often improbable.

So far, we have only considered the disconnected terms, it remains to see how coupled cluster includes connected triple excitations. We have already said that we will not include connected quadruple excitations but their effects have been studied by Bartlett.^{48,49} In order to include T_c one should use the CCSDT scheme. Such a procedure will, indeed, include the connected triples but also numerous higher-order terms yielding a method which goes as n^8 .^{48,49} For most of the chemical problems such a high cost is prohibitive and approximations should be derived. The most radical one would be to decouple the T_c amplitudes from the T_1 and T_2 ones and to calculate their contributions by a MP₄ formula, using however iterative T_1 and T_2 amplitudes: this is the CCSD(T).¹³⁻¹⁵ A similar method can be easily obtained with our procedure. More interesting is to keep the coupling between outer and model space as we do by having the $c_{\alpha}^{T_c}$ in the iterative procedure.

Rewritten in terms of coefficients but in a coupled cluster manner, we have the c_i of the model space and

$$C_3 = C_1 C_2 + C_\alpha^{Tc} + (-\frac{1}{3}C_1^3), \quad (35)$$

$$C_4 = \frac{1}{2}C_2^2 + (-\frac{1}{12}C_1^4), \quad (36)$$

the parenthesis meaning that, at convenience, the powers of C_1 should or not be included. Including all terms would give similar equations as CCSDT so the approximations are done, in fact, in the equation defining T_3 or C_α^{Tc} . Comparing our C_α^{Tc} equation [Eq. (21)] with the T_3 equation as given by Bartlett³⁷ we can see that we include only the first term of this complicated equation. In the nomenclature of Bartlett this scheme corresponds to CCSDT-1a.³⁸ However, in our approach we are free to define the denominator of Eq. (21) by taking MP or EN method or to further improve this denominator by using CEPA arguments.

So, in conclusion, we will say that both methods are quite similar but that we are working with wave function coefficients and thus diagonalizing, whereas the CC approach uses amplitudes of excitations and must solve quite complicated systems of equations. Due to the fact that, nowadays, the diagonalization algorithms are very efficient and can be made direct,^{50,51} we think that our formulation opens the road to very efficient Direct methods as we propose in the present work, or to very efficient CCSDT-*n*.

D. Proposed nomenclature

The acronyms used in this work, which have been designed in order to take into account the great flexibility of the dressed intermediate Hamiltonian method, first indicate the Hamiltonian matrix which is iteratively dressed and diagonalized, say SDCI. Then, in rectangular braces, the terms included in the dressing: T_{dc} and Q for disconnected triples and quadruples, respectively, and T for both connected and disconnected triples. The subindex f stands for “factorization” and p for “perturbation,” as the way of calculating disconnected terms. Inclusion of higher order EPV terms is also indicated by adding +EPV to subindex. Angular braces indicate that the mean value of the dressed Hamiltonians has been taken instead of an iterative diagonalization procedure.

E. Improvements of the SDCI[TQ]_f

Our procedure can be improved in two ways. First one should consider that the factorization theorem is only correct if the energies are additive which is implicit with Möller-Plesset theory. This is a very good approximation, however there are cases where the nonadditivity can be crucial, as for example in Solid State Physics. Examples are easily found in molecular problems also.⁵² Second, one can think of using CEPA arguments in order to improve the description of the triple and quadruple excitations.

1. Nonadditivity of the denominators

If one no longer assumes the energy denominators additivity, one may move back to the perturbative evaluation of c_α as a sum of contributions as in Eq. (17), which is translated diagrammatically in Fig. 1(a). Thus,

$$c_\alpha^Q = \frac{\sum_{(i,j)} (c_i \langle \Phi_\alpha | \mathcal{H} | \Phi_i \rangle + c_j \langle \Phi_\alpha | \mathcal{H} | \Phi_j \rangle)}{E_0 - E_\alpha^0}, \quad (37)$$

where (i,j) stands for the couples of double excitations such that

$$D_j^+ D_i^+ \Phi_0 = \Phi_\alpha. \quad (38)$$

Since

$$\langle \Phi_\alpha | \mathcal{H} | \Phi_i \rangle = \langle \Phi_j | \mathcal{H} | \Phi_0 \rangle, \quad (39)$$

$$\langle \Phi_\alpha | \mathcal{H} | \Phi_j \rangle = \langle \Phi_i | \mathcal{H} | \Phi_0 \rangle, \quad (40)$$

and since

$$c_i^{(1)} = \frac{\langle \Phi_i | \mathcal{H} | \Phi_0 \rangle}{E_0^0 - E_i^0}, \quad (41)$$

$$c_j^{(1)} = \frac{\langle \Phi_j | \mathcal{H} | \Phi_0 \rangle}{E_0^0 - E_j^0}, \quad (42)$$

replacing c_i for $c_i^{(1)}$ (i.e. a variational evaluation of the coefficients of the doubles for the perturbative one), Eq. (37) becomes

$$c_\alpha^Q = \frac{\sum_{(i,j)} c_i c_j [(E_0^0 - E_i^0) + (E_0^0 - E_j^0)]}{E_0^0 - E_\alpha^0}. \quad (43)$$

In a more compact notation, calling Δ the energy differences,

$$c_\alpha^Q = \frac{\sum_{(i,j)} c_i c_j (\Delta_i + \Delta_j)}{\Delta_\alpha} \quad (44)$$

and similarly for the triples

$$c_\alpha^{Tdc} = \frac{\sum_{(i,j)} c_i' c_j' (\Delta_i + \Delta_j)}{\Delta_\alpha}. \quad (45)$$

This formulation will of course use the Epstein–Nesbet definition of \mathcal{H}_0 as the diagonal of the Hamiltonian CI matrix.

2. Inclusion of the EPV terms

This proposal has already been formulated in the preceding paper for the perturbative dressing of the SDCI matrix by the triples and quadruples (SDCI[TQ]_p, previously labeled TD1). Its justification is based on an earlier work,²⁹ proposing a method, (SC)²CI, which makes size-consistent any selected CI. The coefficient c_α of any triple or quadruple Φ_α is supposed to be obtained by considering a CI matrix where Φ_α has been added to the SDCI space. Then the size consistency is obtained by dressing Φ_α by the quantity $E_{corr} + EPV_\alpha$ where E_{corr} is the correlation energy and EPV_α represents the effect of all double excitations which are impossible to perform on Φ_α

$$EPV_\alpha = - \sum_{D_k^+ \Phi_\alpha=0} c_k \langle \Phi_0 | \mathcal{H} | \Phi_k \rangle. \quad (46)$$

Then, the eigenequation for Φ_α leads to the equation

$$c_\alpha = \frac{\sum_i c_i \langle \Phi_\alpha | \mathcal{H} | \Phi_i \rangle}{\langle \Phi_0 | \mathcal{H} | \Phi_0 \rangle - (\langle \Phi_\alpha | \mathcal{H} | \Phi_\alpha \rangle + EPV_\alpha)}. \quad (47)$$

The denominator including the EPV corrections may be used for the linked part of triples only, i.e., the perturbative

residue in the here proposed SDCI[TQ]_f scheme. A direct derivation of the use of this denominator may be obtained as an infinite partial summation of EPV diagrams.⁵³

F. Further improvements

In his concluding remarks of a recent workshop devoted to the coupled cluster methods,⁵⁴ Kutzelnigg pointed out the necessity to imagine some flexible combinations of coupled cluster with lower cost approximations, in order to be able to treat problems with larger number of electrons and/or larger basis sets. Thinking in terms of dressing actually allows one to derive a whole series of approximations combining consistently different methods of different accuracies and costs.

1. Combination with poorer approximations

Suppose that approximate natural orbitals can be obtained by a low cost calculation of the density matrix (for instance, a MP2 calculation). Then, it is possible to retain the most active natural MOs (those of occupation number closer to one) in a class-1. The dressing of the SDCI matrix may be done as follows:

—by the (SC)²SDCI procedure (taking into account the effect of the unlinked triples and quadruples)

—and by treating as a full dressing the effects of the triples and quadruples which only involve holes and particles of the class-1. Then the n^7 time-limiting step runs over a smaller number of MOs.

One may also consider a three-class partition of the NOs, putting into a class-3 the NOs whose occupation numbers are close to 2 or 0. Then the SDCI is reduced to the determinants involving only the MOs of class-1 and class-2. The other doubly excited determinants may be treated through the self-consistently dressed independent pair excitation approximation.⁵⁵ A unique vector of all coefficients obtained at different levels, is used to define EPV terms and to dress in a consistent manner the energies of the large SDCI matrix and of the small 2×2 matrices, as already proposed and practiced.⁵⁶

2. Higher accuracy approximation

One may of course dress a SDTCI matrix by the quadruples and pentuples as was done here for the SDCI matrix, but this will be in general impracticable, except if a preselection of MOs have been done. It seems more reasonable to add the most important triples and quadruples (or more highly excited determinants) to singles and doubles. Then it is perfectly possible to dress the selected triples and quadruples by the unlinked effects of the pentuples and hexuples, as done in the (SC)²CI method. The effect of the other triples and quadruples on the singles and doubles would then be added, including their linked effect, as proposed in the present work. The method would be size-consistent and would include the effect of the leading T_3 and T_4 operators at a very moderate cost. This procedure would allow to combine a CAS main model space with a CCSDT-1a single reference formalism and would certainly be very efficient for the study of multiple bond(s) breaking, without entering the complexity of the MRCC methods.

G. Relation with other methods: Taking the mean value of dressed Hamiltonians

In CC methods, the information about triples and quadruples is stocked in the wave function by means of the e^S operator. However, effective Hamiltonians stock the information in the modified Hamiltonian matrix, and therefore we can obtain a good approximation to \mathcal{E}_0 by calculating the mean value of the $\mathbf{H} + \mathbf{\Delta}$ matrix, $\langle \tilde{\Psi}_0 | \mathbf{H} + \mathbf{\Delta} | \tilde{\Psi}_0 \rangle$, $\tilde{\Psi}_0$ being “good-enough” SD wave function. We can take, for instance, a (SC)²SDCI wave function as it has been made by some of us previously, and will be thereafter named as $\langle \text{SDCI}[\text{TQ}]_f \rangle$.³⁵ This involves no more than one iteration of the CC procedure and it is accurate enough to yield values of \mathcal{E}_0 which, at equilibrium geometries, are in general better than CCSD(T) values.

H. Practical implementation

This method has been implemented in the framework of the algorithm recently proposed by Maynau and Heully⁵⁷ to calculate the perturbation of the triples and quadruples over the SDCI space. This algorithm demonstrates that it is much more rapid to perform an outer loop over the triples and quadruples Φ_α than to generate the appropriate Φ_α 's from all couples (Φ_i, Φ_j) . Thus, for each Φ_α , we obtain all the couples (Φ_i, Φ_j) contributing to it, we determine the coefficient c_α from the coefficients c_i and c_j , and then the contribution of Φ_α to the dressing Δ_i and Δ_j of all the couples (Φ_i, Φ_j) .

The algorithm proceeds as follows:

- (1) Start: A conventional SDCI to get the initial c_i 's.
- (2) Loop over Triples and Quadruples:
 - Calculation of c_α ;
 - calculation of $\mathbf{\Delta}$.
- (3) Building and diagonalization of $\mathbf{H} + \mathbf{\Delta}$.
- (4) Comparison of the \mathcal{E}_0 with the previous value. If convergence is not achieved, one comes back to 2 with the new c_i 's.

If only the dressing of the doubles by the quadruples is taken into account the algorithm results in an iterative CCD diagonalization scheme. If the unlinked contributions to the triples are included too, the dressing of the singles by the triples is taken into account, and then the algorithm results in an iterative CCSD diagonalization scheme. The inclusion of the linked contributions to the triples (and the EPV's) by means of the Epstein–Nesbet partition results in a method similar to CCSDT-1a, which we name as SDCI[TQ]_f (or SDCI[TQ]_{f+EPV}) always in a form of iterative diagonalizations. EPV's have been implemented through the introduction of one-, two-, and three-dimensional arrays,²⁹ which is based on a previous work on the infinite summation of EPV diagrams.⁵³

The algorithm in its present form has the following advantages:

- (1) It converges even where conventional CCD, CCSD, and CCSD(T) does not, for instance, at large interatomic distances in a single or two single bond dissociation process.

TABLE I. Differences (millihartrees) with FCI energies in BH (DZ) and H₂O (DZ and DZP).

Method	BH (DZP)			H ₂ O (DZ)			H ₂ O (DZP)		
	R_e	1,5 R_e	2 R_e	R_e	1,5 R_e	2 R_e	R_e	1,5 R_e	2 R_e
CCSD ^a	1,78	2,64	5,04	1,79	5,59	9,33	4,12	10,16	21,4
SDCI[T _{dc} Q] _f	1,78	2,64	5,03	1,79	5,59	9,55	4,12	10,16	21,5
CCSD(T)	0,41 ^b	0,55 ^b	0,41 ^b	0,57	1,47	-7,70	0,72 ^b	2,00 ^b	-4,64 ^b
CCSDT-1a ^a	0,45	0,61	0,72	0,45	1,46	-5,58	0,60	1,99	-2,65
SDCI[TQ] _f	-0,15	-0,34	-1,22	0,06	-0,38	-15,11	-0,29	-1,04	-14,47
SDCI[TQ] _{f+EPV}	-0,07	-0,18	-0,68	0,11	0,11	-10,29	-0,14	-0,25	-9,70
SDCI[TQ] _{f(MP)}	0,46	0,63	1,02	0,60	2,01	-2,51
CCSDT ^a	0,06	0,02	0,02	0,44	1,52	-1,87	0,53	1,83	-2,17
CC5SD(TQ*) ^c	0,05	0,01	-0,59	0,19	0,13	-1,96
CCSDTQ ^d	0,00	0,00	+0,00	0,02	0,14	-0,02

^aReference 16.^bReference 14.^cReference 48.^dReference 49.

- (2) At equilibrium distances the number of iterations to converge within 1 μ H is typically of 4 to 6.
- (3) Its possibility to take benefit of the parallel machines. Slave processors can calculate independently the contributions of different triples or quadruples to Dressing matrix Δ .⁵⁸ Moreover, a parallel algorithm to diagonalize large matrices is in progress. Thus, the possibility to calculate very large systems, or to use very large basis sets is open.

The main disadvantage of this algorithm in its present implementation is that it grows as n^8 , although it could take benefit of the same reduction to n^6 or n^7 that can be achieved on CCSD or CCSDT-1a methods.¹⁹ Test calculations on H₂O with a DZ basis set shows that the present implementation of the SDCI[TQ]_f is three times faster than full CCSDT, but three times slower than CCSDT-1a one, the CC calculations being performed with the Scuseria's series of programs.⁵⁹

IV. NUMERICAL RESULTS AND DISCUSSION

We have applied our methodology to a few examples. Two of them are classic comparisons with benchmark calculations, the other two refer to 4-electron systems previously presented:³⁶ the square to rectangular deformation of the H₄ system, and the breaking of a single bond in presence of an electron pair, in the T-shaped Li₄ system. These systems represent situations where a degeneracy appears between Φ_0 and one of the doubles. We present also the results on another 4-electron system which dissociates into two closed shell atoms, the Be₂ molecule.

A. Benchmark calculations: BH and H₂O molecules

As a first application of the method we have considered the BH molecule. The reference calculations are those DZP FCI ones at R_e from Ref. 60, and at 1,5 R_e and 2 R_e from Ref. 16. The second application concerns to H₂O molecule with a double- ζ (DZ) and double- ζ plus polarization basis set (DZP). Comparisons are made with the benchmark calculations at DZ FCI level at R_e by Saxe *et al.*,⁶¹ and at 1,5 R_e

and 2 R_e by Harrison *et al.*,⁶⁰ and at DZP level the frozen core FCI calculations at the same three distances by Bauschlicher *et al.*⁶² The differences in mH relative to FCI values are shown in Table I. The values for conventional CC methods are taken from Refs. 14, 16, 48, and 49. The perturbative inclusion of the triples in an iterative manner, by means of the Epstein–Nesbet partition (results marked as SDCI[TQ]_f) shows the trends expected from this approach. Since higher order effects are included, the correlation energies lies under the FCI value in almost all the cases considered. However, the values at short distances are closer to FCI than the corresponding Möller–Plesset CCSDT-1a ones. Inclusion of the EPV higher order effects makes the results (named as SDCI[TQ]_{f+EPV}) closer to FCI. In fact, at short distances, the SDCI[TQ]_{f+EPV} results are comparable to full CCSDT ones. However, at 2 R_e , both SDCI[TQ]_f and SDCI[TQ]_{f+EPV} methods give values far from FCI results. Table I shows also the DZP results for BH and H₂O molecules when the linked contributions to triples are taken into account within the Möller–Plesset partition, marked as SDCI[TQ]_{f(MP)}. It is to note that the results of SDCI[TQ]_{f(MP)} calculations are almost identical to CCSDT-1a ones near the equilibrium distances. At long distances the differences can be related to the different treatment of the T_1 terms. When one takes into account only the effects of the nonlinked contributions to triples and quadruples (named as SDCI[T_{dc}Q]_f), the values obtained at R_e and 1,5 R_e are the same as those obtained with CCSD, but small differences appear at 2 R_e in the water molecule. These differences could be related with the absence of terms including powers of T_1 , which are important when polar single bonds are simultaneously broken.

B. The rectangular to square H₄ problem

This model problem has been a critical test for many methods, especially for CC methods.^{63–65} Along the rectangular to square transformation for H₄ system a degeneration appears between the Φ_0 determinant and another doubly excited determinant. Geometry has been explicated elsewhere.³⁶ The calculations have been performed with a double- ζ basis set. Table II shows the energy differences relatives to FCI

TABLE II. Rectangular H_4 system in a double zeta basis set as a function of R (bohr) for $r=1.40158$ bohr. FCI energies in hartrees and differences to FCI in millihartrees.

R	$E(\text{FCI})$	$\text{SDCI}[\text{TQ}]_f$	$\text{SDCI}[\text{TQ}]_{f+\text{EPV}}$	$\langle \text{SDCI}[\text{T}]_p[\text{Q}]_f \rangle$	$\text{CCSD}(\text{T})$	$\text{QSDCI}(\text{T})$
1,20	-1,793989	0,00	0,01	0,05	0,16	0,17
1,24	-1,802647	-0,02	0,00	0,07	0,23	0,24
1,28	-1,810103	-0,06	-0,02	0,11	0,34	0,35
1,32	-1,817367	-0,18	-0,11	0,16	0,48	0,51
1,36	-1,826043	-0,54	-0,39	0,17	0,45	0,54
1,40	-1,838209	-1,42	-1,11	-0,13	-1,04	-0,77
1,40158	-1,838784	-1,47	-1,15	-0,16	-1,17	-0,89
1,42	-1,846002	-1,01	-0,77	0,03	-0,13	0,05
1,44	-1,854827	-0,65	-0,49	0,11	0,32	0,43
1,46	-1,864440	-0,42	-0,31	0,13	0,43	0,50
1,48	-1,874579	-0,28	-0,19	0,12	0,42	0,47
1,50	-1,885016	-0,19	-0,12	0,11	0,38	0,41
1,52	-1,895580	-0,13	-0,08	0,09	0,33	0,35
1,54	-1,906151	-0,09	-0,05	0,08	0,28	0,29
1,56	-1,916645	-0,06	-0,03	0,07	0,24	0,25
1,58	-1,927007	-0,04	-0,02	0,06	0,21	0,21
1,60	-1,937201	-0,03	-0,01	0,06	0,18	0,18
1,80	-2,027364	0,00	0,01	0,03	0,07	0,07
2,00	-2,096821	0,00	0,01	0,03	0,04	0,04
2,20	-2,149662	0,00	0,01	0,02	0,03	0,03
2,40	-2,189660	0,00	0,01	0,02	0,02	0,02
2,60	-2,219768	0,00	0,00	0,02	0,02	0,02
2,80	-2,242283	0,00	0,00	0,02	0,01	0,02
20,00	-2,303002	0,01	0,01	0,00	0,00	0,00

values in millihartrees calculated at $\text{SDCI}[\text{TQ}]_f$, $\text{SDCI}[\text{TQ}]_{f+\text{EPV}}$, and $\langle \text{SDCI}[\text{T}]_p[\text{Q}]_f \rangle$ levels of theory and compared with the $\text{CCSD}(\text{T})$ and $\text{QSDCI}(\text{T})$ values for different intermolecular distances R . All these results are also

displayed in Fig. 4. The behaviour of the $\text{SDCI}[\text{TQ}]_f$ and $\text{SDCI}[\text{TQ}]_{f+\text{EPV}}$ methods is very similar, with the $\text{SDCI}[\text{TQ}]_{f+\text{EPV}}$ values slightly closer to FCI results. At the degeneracy point, all the iterative methods give nearly the


FIG. 4. Energy differences in mH relative to FCI in the H_4 problem in a double- ζ basis set, as a function of R for $r=1.40158$ a.u.

TABLE III. T-shaped Li_4 system in an ANO $[2s, 1p]$ basis set as a function of the axial bond length, R (in bohr). FCI energies in hartrees and differences to FCI in millihartrees.

R	$E(\text{FCI})$	$\text{SDCI}[\text{TQ}]_f$	$\text{SDCI}[\text{TQ}]_{f+\text{EPV}}$	$\text{CCSD}(\text{T})$	$\text{QSDCI}(\text{T})$	$\langle \text{SDCI}[\text{T}]_p[\text{Q}]_f \rangle$
4,00	-29,772243	-0,31	-0,11	0,16
4,40	-29,783253	-0,30	-0,11	0,13
4,80	-29,789148	-0,31	-0,10	0,49	...	0,10
5,00	-29,790673	-0,31	-0,11	0,51	0,49	0,09
5,20	-29,791469	-0,32	-0,10	0,53	0,51	0,09
5,40	-29,791674	-0,33	-0,10	0,55	0,53	0,08
5,60	-29,791404	-0,34	-0,10	0,57	0,55	0,07
5,80	-29,790761	-0,35	-0,10	0,60	0,57	0,07
6,00	-29,789830	-0,36	-0,10	0,63	0,60	0,07
6,20	-29,788681	-0,38	-0,10	0,66	0,63	0,07
6,40	-29,787375	-0,40	-0,10	0,69	0,67	0,08
6,60	-29,785964	-0,42	-0,10	0,73	0,71	0,08
6,80	-29,784489	-0,44	-0,10	0,77	0,75	0,08
7,00	-29,782986	-0,48	-0,10	0,81	0,79	0,09
7,20	-29,781484	-0,51	-0,10	0,86	0,84	0,09
7,40	-29,780007	-0,56	-0,11	0,91	0,89	0,09
7,60	-29,778575	-0,61	-0,11	0,96	0,94	0,09
7,80	-29,777203	-0,68	-0,12	1,01	0,99	0,09
8,00	-29,775901	-0,75	-0,13	2,52	1,04	0,08
9,00	-29,770660	-1,32	-0,25	3,31	1,24	-0,05
10,00	-29,767460	-2,17	-0,46	1,19	1,22	-0,35
20,00	-29,763831	-7,94	-1,67	a	a	-2,97
40,00	-29,763798	-8,92	-1,73	a	a	-3,32
100,00	-29,763798	-9,68	-1,76	a	a	-3,54

^aMethod does not converge.

same error (0,9 to 1,5 mH), but for all other distances, out of the $R=r$ point, the $\text{SDCI}[\text{TQ}]_f$ and $\text{SDCI}[\text{TQ}]_{f+\text{EPV}}$ methods remain closer to FCI values than $\text{CCSD}(\text{T})$ and $\text{QSDCI}(\text{T})$ ones. Moreover, their behaviour is also more regular, since the deviation from FCI has the same sign for all values of R . The shape of the error curve for the mean value $\text{SDCI}[\text{TQ}]_f$ method ($\langle \text{SDCI}[\text{T}]_p[\text{Q}]_f \rangle$),³⁵ is similar to that of $\text{CCSD}(\text{T})$ or $\text{QSDCI}(\text{T})$ ones, but its mean error is smaller. We point out that, for this system the errors reported in the present work, in the degeneracy region, are larger than those obtained from the $\text{SDCI}[\text{TQ}]_p$ methods,³⁶ where the coefficients of the triples and quadruples are estimated perturbatively. At the degeneracy point ($r=R$) these errors were 0,5 mH for the method $\text{SDCI}[\text{TQ}]_p$ and 0,09 mH for the method $\text{SDCI}[\text{TQ}]_{p+\text{EPV}}$. The small errors in $\text{SDCI}[\text{TQ}]_{p+\text{EPV}}$ method could be explained because in it, higher-order EPV corrections are included for all contributions to triples and quadruples.

C. The T-shaped Li_4 cluster

The geometric parameters describing the breaking of a single bond in the T-shaped Li_4 cluster four electron system have been described in the previous work.³⁶ Calculations have been carried out with a DZP $[2s, 1p]$ ANO's basis set.⁶⁶ The values for the four electron FCI total energy and the differences from the FCI values are shown in Table III, and in Fig. 5 the potential energy curve for the dissociation process is displayed. The errors of the $\text{SDCI}[\text{TQ}]_f$ method with relation to FCI are of the same order or somewhat lesser than those of $\text{CCSD}(\text{T})$ or $\text{QSDCI}(\text{T})$ methods, although of opposite sign, due to the Epstein-Nesbet choice of the zeroth

order Hamiltonian. When higher order EPV corrections are included in the linked contributions to triples, the errors fall to $-0,1$ mH in almost all the range of values of R . Note here the convergence of our algorithm, based on iteratively diagonalizing a dressed SDCI matrix, by comparison with CC or QCI methods which fail to converge for large values of R . $\text{SDCI}[\text{TQ}]_f$ curve (Fig. 5) shows the typical behaviour of the CC dissociation curves, where dissociation limit lies far below the FCI dissociation limit. The EPV terms avoid the most of this deficiency, since they increase the resistance to the degeneracies occurring at long bond distances.

D. The Be_2 molecule

Be_2 molecule is a very challenging system for theoretical methods.^{37,67} Despite the closed shell character of the Be atom, a true chemical bond is formed between two Be atoms at short distances.⁶⁸ Most of the theoretical methods fail to represent this bond, and a good basis set is necessary.⁶⁷ Figure 6 shows the potential energy curves calculated with four electrons and a $[3s, 2p, 1d]$ ANO basis set,⁶⁶ and FCI, $\text{CCSD}(\text{T})$ and the $\text{SDCI}[\text{TQ}]_f$ methods. The FCI energies and the differences to FCI energies for these methods are shown in Table IV. The most striking feature is the inability of the $\text{CCSD}(\text{T})$ method to merely reproduce a bond between the two atoms with this basis set. The importance of the linked triples is very large for this system, and they are included by means of a MP single calculation in $\text{CCSD}(\text{T})$ method. If the linked triples are taken into account by an iterative EN procedure, as in the $\text{SDCI}[\text{TQ}]_f$ method, the main well at short lengths appears, but the bond is overestimated, and the minimum well is too deep. When higher-




FIG. 5. Evolution of the energy in the T-shaped Li_4 problem in a $[2s, 1p]$ ANO basis set, as a function of the length R of the axial bond.

order EPV's are included, as in the SDCI[TQ]_f+EPV method, the resulting curve is almost identical to the FCI one, and the errors, near R_e , are equal to 0,02 mH or smaller, 2 orders of magnitude less than CCSD(T) ones. At long distances, the

three methods converge towards the FCI dissociation limit. It is to note the $\langle \text{SDCI}[\text{T}]_p[\text{Q}]_f \rangle$ results, which are fairly parallel to the FCI curve, although the errors at the dissociation limit are somehow larger than in the minimum region.




FIG. 6. Evolution of the energy in the Be_2 molecule in a $[3s, 2p, 1d]$ ANO basis set, as a function of the bond length, R .

TABLE IV. Be₂ molecule in an ANO [3*s*,2*p*,1*d*] basis set as a function of the internuclear distance, *R* (in bohr). FCI energies in hartrees and differences to FCI in millihartrees.

<i>R</i>	<i>E</i> (FCI)	SDCI[TQ] _{<i>f</i>}	SDCI[TQ] _{<i>f</i>} +EPV	CCSD(T)	⟨SDCI[T] _{<i>p</i>} [Q] _{<i>f</i>} ⟩
4	-29,229824	-0,92	-0,12	1,51	0,23
4,4	-29,235853	-0,73	-0,08	1,36	0,30
4,8	-29,237368	-0,54	-0,04	1,15	0,39
5	-29,237467	-0,46	-0,02	1,03	0,43
5,2	-29,237414	-0,39	-0,01	0,92	0,46
5,6	-29,237227	-0,28	0,00	0,71	0,50
6	-29,237140	-0,21	0,00	0,53	0,52
7	-29,237231	-0,10	0,00	0,25	0,54
8	-29,237255	-0,05	0,00	0,12	0,55
9	-29,237158	-0,02	0,00	0,06	0,55
10	-29,237044	-0,01	0,00	0,03	0,55
12	-29,236905	0,00	0,00	0,01	0,54
14	-29,236853	0,00	0,00	0,00	0,54
16	-29,236834	0,00	0,00	0,00	0,54
20	-29,236824	0,00	0,00	0,00	0,54

V. CONCLUDING REMARKS

Within the intermediate Hamiltonian framework, we have formulated a total dressing of a SDCI Hamiltonian matrix for a closed-shell single reference. The present formulation, more general than CI or CC methods, includes these as particular cases, and allows an easy extension of the CC series. Also, the flexibility in the definition of the dressing matrix allows for the definition of mixed approximated methods or selected CC-like methods. Moreover, the possibility of dealing with MRCI matrices open the door for MR-CC-like formalisms. Since the method proceeds by diagonalization of the SDCI matrix, the effect of the quadruples and the triples on all the single and doubly excited determinants includes an infinite partial summation of higher order diagrams. An estimation of the coefficients of the non-linked triples and quadruples in a coupled cluster fashion results in an iterative method which includes all the fourth order, and which differs from the CCSD method in the lack of the terms including third and fourth powers of the T_1 (or C_1) operator. If the linked contributions to triples are included perturbatively, the resulting method is similar to the previous CCSDT-1a, but differing in the Epstein–Nesbet choice of the unperturbed Hamiltonian. This difference is evidenced in the slight overestimation of the correlation energy, which is a characteristic of this choice. Inclusion of higher order EPV's correction is trivial in EN framework, and the resulting method compares well with CCSDT except for long distances when two single bonds are broken simultaneously, as in the H₂O dissociation process. The numerical stability of the algorithm is also remarkable, based in an hermitian row and column dressing and the diagonalization of a SDCI dressed matrix, which allows it to converge even when traditional CC algorithms cannot. The present implementation of this algorithm involves a loop over all the triples and quadruples, and therefore the time of calculation grows as n^8 . However, it has been completely parallelized, and it could take benefit of the same intermediate summations which make CCSD a n^6 algorithm. Numerical applications on some model systems show that the combination of

EN zeroth order Hamiltonian and EPV higher order corrections give very good estimations of the FCI correlation energy.

ACKNOWLEDGMENTS

The research reported in this paper has been supported by the DGICYT Project No. PB91-0634, the SEUI Grant No. OP90-0042, and a grant from Conselleria de Cultura, Educació i Ciència de la Generalitat Valenciana. Authors are indebted to Professor G. Scuseria and B.O. Roos who kindly has sent to us their CC and MOLCAS-2 series of programs, respectively. Computational support from the C.I.U.V. of Universitat de València is also acknowledged.

¹K. A. Brueckner, Phys. Rev. **97**, 1353 (1953).

²K. A. Brueckner, Phys. Rev. **100**, 36 (1955).

³H. Primas, in *Modern Quantum Chemistry*, edited by O. Sinanoglu (Academic, New York, 1961), Vol. 2, p. 45.

⁴W. Kutzelnigg, in *Modern Theoretical Chemistry*, edited by H. F. Schaefer (Plenum, New York, 1977), Vol. 2.

⁵A. Szabo and N. S. Ostlund, *Modern Quantum Chemistry. Introduction to Advanced Electronic Structure Theory*, (McGraw-Hill, New York, 1989).

⁶F. Coestler, Nucl. Phys. **1**, 421 (1958).

⁷F. Coestler and H. Kummel, Nucl. Phys. **17**, 477 (1960).

⁸J. Cizek, Adv. Chem. Phys. **14**, 35 (1969).

⁹J. Cizek and J. Paldus, Phys. Scr. **21**, 251 (1980).

¹⁰R. J. Bartlett, Ann. Rev. Phys. Chem. **32**, 359 (1981).

¹¹R. J. Bartlett, J. Phys. Chem. **93**, 1697 (1989).

¹²J. Paldus, in *Methods in Computational Molecular Physics*, edited by S. Wilson and G. H. F. Dierksen (Plenum, New York, 1992), p. 99.

¹³K. Raghavachari, G. W. Trucks, J. A. Pople, and M. Head-Gordon, Chem. Phys. Lett. **157**, 479 (1989).

¹⁴R. J. Bartlett, J. D. Watts, S. A. Kucharski, and J. Noga, Chem. Phys. Lett. **165**, 513 (1990).

¹⁵R. J. Bartlett, J. D. Watts, S. A. Kucharski, and J. Noga, Chem. Phys. Lett. **167**, 609 (1990).

¹⁶J. Noga and R. J. Bartlett, J. Chem. Phys. **86**, 7041 (1987).

¹⁷J. Noga and R. J. Bartlett, J. Chem. Phys. **89**, 3401 (1988).

¹⁸G. E. Scuseria and H. F. Schaefer III, Chem. Phys. Lett. **152**, 382 (1988).

¹⁹G. E. Scuseria, C. L. Janssen, and H. F. Schaefer III, J. Chem. Phys. **89**, 7382 (1988).

²⁰P. Piecuch and L. Adamowich, J. Chem. Phys. **100**, 5857 (1994).

²¹H. P. Kelly and M. A. Sessler, Phys. Rev. **132**, 2091 (1963).

²²H. P. Kelly and M. A. Sessler, Phys. Rev. A **134**, 1450 (1964).

²³W. Meyer, Int. J. Quantum Chem. Symp. **5**, 341 (1971).

- ²⁴W. Meyer, *J. Chem. Phys.* **58**, 1017 (1973).
- ²⁵W. Meyer, *Theor. Chim. Acta* **35**, 277 (1974).
- ²⁶R. Ahlrichs, H. Lishka, V. Staemmler, and W. Kutzelnigg, *J. Chem. Phys.* **62**, 1225 (1975).
- ²⁷R. Ahlrichs, P. Scharf, and C. Ehrhardt, *J. Chem. Phys.* **82**, 890 (1985).
- ²⁸D. P. Chang and S. R. Langhoff, *J. Chem. Phys.* **84** (1986).
- ²⁹J. P. Daudey, J. L. Heully, and J. P. Malrieu, *J. Chem. Phys.* **99**, 1240 (1993).
- ³⁰J. Rubio, A. Povill, F. Illas, and J. P. Malrieu, *Chem. Phys. Lett.* **200**, 559 (1992).
- ³¹A. Povill, R. Caballol, J. Rubio, and J. P. Malrieu, *Chem. Phys. Lett.* **209**, 126 (1993).
- ³²R. Caballol and J. P. Malrieu, *Chem. Phys. Lett.* **188**, 543 (1992).
- ³³J. P. Malrieu, P. Durand, and J. P. Daudey, *J. Phys. A* **18**, 809 (1985).
- ³⁴P. Durand and J. P. Malrieu, in *Ab initio Methods in Quantum Chemistry*, edited by K. P. Lawley (Wiley, New York, 1986), Vol. 1, p. 321.
- ³⁵J. Sánchez-Marín, D. Maynau, and J. P. Malrieu, *Theor. Chim. Acta* **87**, 107 (1993).
- ³⁶J. P. Malrieu, I. Nebot-Gil, and J. Sánchez-Marín, *J. Chem. Phys.* **100**, 1440 (1994).
- ³⁷Y. S. Lee, S. A. Kucharski, and R. J. Bartlett, *J. Chem. Phys.* **81**, 5906 (1984).
- ³⁸M. Urban, J. Noga, S. J. Cole, and R. J. Bartlett, *J. Chem. Phys.* **83**, 4041 (1985).
- ³⁹H. Feschbach, *Ann. Rev. Nucl. Sci.* **8**, 49 (1958).
- ⁴⁰H. Feschbach, *Ann. Phys. (N.Y.)* **5**, 357 (1958).
- ⁴¹H. Feschbach, *Ann. Phys. (N.Y.)* **19**, 287 (1962).
- ⁴²P. O. Lowdin, *J. Math. Phys.* **3**, 969 (1962).
- ⁴³J. L. Heully and J. P. Malrieu, *Chem. Phys. Lett.* **199**, 545 (1992).
- ⁴⁴I. Lindgren and J. Morrison, *Atomic Many-Body Theory* (Springer, Berlin, 1986).
- ⁴⁵R. A. Chiles and C. A. Dykstra, *J. Chem. Phys.* **74**, 4544 (1981).
- ⁴⁶N. C. Handy, J. A. Pople, M. Head-Gordon, K. Raghavachari, and G. W. Trucks, *Chem. Phys. Lett.* **164**, 185 (1989).
- ⁴⁷I. Nebot-Gil, J. Sánchez-Marín, J. L. Heully, and J. P. Malrieu, *Chem. Phys. Lett.* **234**, 45 (1995).
- ⁴⁸S. A. Kucharski and R. J. Bartlett, *J. Chem. Phys.* **97**, 4282 (1992).
- ⁴⁹S. A. Kucharski and R. J. Bartlett, *Chem. Phys. Lett.* **206**, 574 (1993).
- ⁵⁰B. O. Roos and P. E. M. Siegbahn, in *Methods of Electronic Structure Theory*, edited by H. F. Schaefer III (Plenum, New York, 1977), p. 277.
- ⁵¹J. Olsen, B. O. Roos, P. Jørgensen, and H. J. Hensen, *J. Chem. Phys.* **89**, 2185 (1988).
- ⁵²J. P. Malrieu, J. P. Daudey, and R. Caballol, *J. Chem. Phys.* **101**, 8908 (1994).
- ⁵³M. B. Lepetit and J. P. Malrieu, *J. Chem. Phys.* **87**, 5937 (1987).
- ⁵⁴W. Kutzelnigg, Bad Honef Workshop on Coupled Cluster methods, Bad Honef (R.F.A.), October 1994.
- ⁵⁵M. B. Lepetit and J. P. Malrieu, *Chem. Phys. Lett.* **208**, 503 (1993).
- ⁵⁶J. Meller, J. L. Heully, and J. P. Malrieu, *Chem. Phys. Lett.* **218**, 276 (1994).
- ⁵⁷D. Maynau and J. L. Heully, *Chem. Phys. Lett.* **211**, 625 (1993).
- ⁵⁸W. Diaz, I. Nebot-Gil, D. Maynau, and J. Sanchez-Marín (unpublished).
- ⁵⁹G. E. Scuseria and T. J. Lee, *J. Chem. Phys.* **93**, 5851 (1990).
- ⁶⁰R. J. Harrison and N. C. Handy, *Chem. Phys. Lett.* **95**, 386 (1983).
- ⁶¹P. Saxe, H. F. Schaefer III, and N. C. Handy, *Chem. Phys. Lett.* **79**, 202 (1981).
- ⁶²C. W. Bauschlicher, Jr. and P. R. Taylor, *J. Chem. Phys.* **85**, 2779 (1986).
- ⁶³K. Jankovski and J. Paldus, *Int. J. Quantum Chem.* **18**, 1243 (1980).
- ⁶⁴K. Jankovski, J. Paldus, and J. Wasilewski, *J. Chem. Phys.* **95**, 3549 (1991).
- ⁶⁵K. Jankovski, J. Paldus, I. Grabowski, and K. Kowalski, *J. Chem. Phys.* **97**, 1600 (1992).
- ⁶⁶P.-O. Widmark, P.-A. Malmqvist, and B. O. Roos, *Theor. Chim. Acta* **77**, 291 (1990).
- ⁶⁷S. Evangelisti, G. L. Bendazzoli, and L. Gagliardi, *Chem. Phys.* **185**, 47 (1994).
- ⁶⁸V. E. Bondybey, *Chem. Phys. Lett.* **109**, 436 (1984).