

High Incidence of Chromosomal Abnormalities in Large-Headed and Multiple-Tailed Spermatozoa

Case Report

EMILIA MATEU,* LORENA RODRIGO,*
NICOLAS PRADOS,† MANUEL GIL-SALOM,*
JOSÉ REMOHÍ,* ANTONIO PELLICER,*
AND CARMEN RUBIO*

From the **Instituto Valenciano de Infertilidad, University of Valencia, Valencia, Spain; and †Instituto Valenciano de Infertilidad, Sevilla, Spain.*

Case Report

To date, several studies of aneuploidy rate in spermatozoa from infertile men have been published. Most of them found an increase incidence of aneuploid spermatozoa in infertile patients compared with normal donors (Moosani et al, 1995; Finkelstein et al, 1998; McInnes et al, 1998; Pang et al, 1999; Rives et al, 1999; Vegetti et al, 2000). However, in other studies, no differences in aneuploidy rates between fertile and infertile men were reported (Miharu et al, 1994; Guttenbach et al, 1997). The evaluated patients often showed different types of sperm parameter impairments, including oligo-, astheno-, oligoastheno-, oligoterato-, oligoastenoterato-, asthenoteratozoospermia, and unexplained infertility, and this may be responsible, at least in part, for the discrepancies observed in the results.

Studies taking into account an abnormality for an isolated sperm defect have shown an inverse correlation between sperm aneuploidy and concentration (Bernardini et al, 1998, 2000; Pang et al, 1999; Pfeffer et al, 1999; Rives et al, 1999; Nishikawa et al, 2000; Ushijima et al, 2000; Vegetti et al, 2000; Calogero et al, 2001a; Ohashi et al, 2001; Rubio et al, 2001; Martin et al, 2003). Furthermore, some authors have reported an association between the aneuploidy rate and the presence of abnormal head morphology and particularly with enlarged heads and multiple tails (Yurov et al, 1996; In't Veld et al, 1997; Bernardini et al, 1998; Viville et al, 2000; Devillard et al, 2002).

Because intracytoplasmic sperm injection (ICSI) is widely acknowledged to be the most effective therapeutic approach for severe male-factor infertility, including ter-

atozoospermia, it is important to counsel the couple about the risk of aneuploidy in their offspring.

We describe the case of a patient with a high incidence of sperm chromosomal abnormalities associated with morphology with large-headed and multiple-tailed spermatozoa.

Patient—We studied a 38-year-old infertile man with a clinical background of left varicocelectomy 4 years ago. Blood karyotype of the patient, as well as molecular analysis of Y chromosome microdeletions, were normal. He worked in a petroleum refinery and he was exposed daily to chemical agents such as naphtha, sulfhidric acid, asphalt, and ammonium. Four semen analyses during 10 months (Table 1) revealed a moderate oligozoospermia, severe asthenozoospermia, and total teratozoospermia according to the World Health Organization (WHO, 1999) criteria. Large-headed and multiple-tailed spermatozoa were observed in most of them (the Figure). The female partner was 37 years old and without any infertility problem.

The couple underwent a previous ICSI cycle in which the microinjection was technically difficult because of the size of the sperm heads and the absence of spermatozoa with normal morphology. In this cycle, 15 oocytes were microinjected, 11 fertilized, but embryo quality was impaired on day 3 of development (3 of them were blocked and the remaining embryos were of poor quality according to Alikani et al, 1999). A total of 4 embryos were transferred after assisted hatching and fragment removal and pregnancy was not achieved. Due to the poor embryo quality, the couple was offered to undergo a sperm analysis by fluorescence in situ hybridization (FISH).

FISH—The sperm sample was fixed with methanol: acetic acid (Merck, Darmstadt, Germany) and spread in superfrost/plus slides (O. Kindler, GmbH, Freiburg, Germany). For FISH analysis, sperm nuclei were decondensed by slide incubation for 5 to 7 minutes at 37°C in 5 mmol/L 1,4-Dithiothreitol (DTT, Roche Diagnostics, GmbH Mannheim, Germany) and 1% Triton X-100. DNA was denatured for 5 minutes at 73°C ± 1°C in a water bath in 70% formamide (Roche Diagnostics GmbH, Mannheim, Germany). Triple FISH was performed for chromosomes 18 (locus D18Z1, CEP 18 Spectrum Aqua; Vysis Inc Downers Grove, Ill), X (locus DXZ1, CEP X Spectrum Green; Vysis Inc), and Y (locus DYZ1, CEP Y Spectrum Orange; Vysis Inc). Double FISH was performed for chromosomes 13 (locus RB, LSI 13 Spectrum

Correspondence to: Dr Emilia Mateu, Instituto Valenciano de Infertilidad, Plaza Policía Local 3, 46015 Valencia, Spain (e-mail: emateu@ivi.es).

Received for publication May 12, 2005; accepted for publication August 19, 2005.

DOI: 10.2164/jandrol.05033

Table 1. Results of patient's spermograms

Sample	1	2	3	4
Volume (mL)	4	4.5	3	4
Concentration (10 ⁶ /mL)	15	10	10	12
% motile sperm	26	10	7	5
% live sperm	30	30	10	5
Morphology (% normal)	0	0	0	0

Green; Vysis Inc) and chromosome 21 (loci D21S259, D21S341, D21S342, LSI 21 Spectrum Orange; Vysis Inc) on a different slide. FISH incubation and detection were performed according to the manufacturer's instructions.

Analysis was carried out using an Olympus AX70 epi-fluorescence microscope equipped with a triple band-pass filter for 4'6-diamidino-2-phenylindole (DAPI)/Texas Red/fluorescein isothiocyanate (FITC) and single band-pass filters for FITC, Texas Red, and Aqua Blue.

Ejaculated spermatozoa from 5 normozoospermic fertile donors, classified according to WHO (1999) criteria and processed in the same manner as the samples of the patient were used as control group (Rodrigo et al, 2004).

Statistical Analysis—The incidence of disomy and diploidy for the analyzed chromosomes in the patient and in a control group of normozoospermic individuals were compared using χ^2 test (with Yates correction when necessary) and Fisher's exact test (Graphpad InStat v. 2.05a, Graphpad Software, San Diego, Calif).

Results


The results of triple- and double-color FISH are reported in Tables 2 and 3. Table 2 shows the distribution of the different types of abnormalities for chromosomes X, Y, and 18. A total of 215 sperm heads were scored for these chromosomes and 196 were abnormal (91.2%). In our control group, only 179 out of 50 404 (0.36%) were abnormal (Rodrigo et al, 2004), showing clear statistical differences ($P < .0001$).

Table 3 shows the distribution of the different types of

Table 2. FISH results for chromosomes X, Y, and 18

	n	%
18X (normal)	11	5.12
18Y (normal)	8	3.72
18XY (disomy)	24	11.16
18YY (disomy)	1	0.47
18XXY (trisomy)	8	3.72
18XYY (trisomy)	4	1.86
18XXYY (tetrasomy)	5	2.33
1818X (disomy)	8	3.72
1818Y (disomy)	4	1.86
181818Y (trisomy)	2	0.93
1818XX (diploidy)	4	1.86
1818YY (diploidy)	2	0.93
1818XY (diploidy)	22	10.23
181818XYY (triploidy)	6	2.79
181818XXY (triploidy)	3	1.40
18181818XXYY	4	1.86
18.	4	1.86
._X	1	0.47
._Y	1	0.47
._XY	2	0.93
._XXY	1	0.47
._XYY	1	0.47
1818.	1	0.47
1818XXYY	25	11.63
1818XXY	11	5.12
1818XYY	10	4.65
181818XY	12	5.58
181818YY	1	0.47
181818XX	3	1.40
181818XXYY	12	5.58
18181818XY	5	2.33
18181818XXY	3	1.40
18181818XYY	6	2.79
Number of cells scored	215	...
Total abnormal cells	196	91.2

abnormalities for chromosomes 13 and 21. In this analysis, a total of 102 sperm heads were evaluated and 96 were abnormal (94.1%). In the control group, 50 373 spermatozoa were analyzed and 186 were abnormal (0.37%), again with marked statistical differences ($P < .0001$).



Arrows indicate multiple tails.

Table 3. FISH results for chromosomes 13 and 21

	n	%
1321 (normal)	6	5.88
131321 (disomy)	4	3.92
13131321 (trisomy)	1	0.98
1313131321 (tetrasomy)	1	0.98
132121 (disomy)	5	4.90
13212121 (trisomy)	1	0.98
1321212121 (tetrasomy)	2	1.96
13132121 (diploidy)	10	9.80
131313212121 (triploidy)	17	16.67
1313131321212121 (tetraploidy)	18	17.65
..212121	2	1.96
..2121	1	0.98
..21	2	1.96
13..	4	3.92
1313..	1	0.98
131321212121	4	3.92
1313212121	2	1.96
13131321212121	6	5.88
1313132121	9	8.82
13131313212121	3	2.94
131313132121	3	2.94
Number of cells scored	102	...
Total abnormal cells	96	94.1

Discussion

The available literature on FISH analysis of human sperm confirms higher rates of sperm aneuploidy in infertile men as compared with fertile men, despite a normal blood karyotype (Moosani et al, 1995; Lähdetie et al, 1997; Bernardini et al, 1998; Arán et al, 1999; Pang et al, 1999; Pfeffer et al, 1999; Rives et al, 1999; Ushijima et al, 2000; Vegetti et al, 2000; Martin et al, 2003). An inverse correlation between sperm quality and sperm aneuploidy rates has been reported (Bernardini et al, 2000; Nishikawa et al, 2000; Ushijima et al, 2000; Vegetti et al, 2000; Rubio et al, 2001), but contradictory results have been published concerning the relationship of sperm aneuploidy with specific sperm defects.

An inverse relationship has been reported between sperm aneuploidy and sperm concentration (Pang et al, 1999; Vegetti et al, 2000; Calogero et al, 2001b; Rubio et al, 2001) and between sperm aneuploidy and sperm motility (Vegetti et al, 2000). With respect to the sperm morphology, there are conflicting results, mainly due to the variability of sperm morphological types between patients and within the same sperm sample.

There have been several reports regarding the relationship of specific morphological types to the incidence of aneuploidies. Significant differences in aneuploidy rates have not been found between normal controls and patients with shortened flagella syndrome and spermatozoa with irregular acrosomes (Viville et al, 2000). Regarding globozoospermia, there are few studies and they yield controversial results: some of them found an increased inci-

dence of sperm chromosomal abnormalities (Carrell et al, 1999, 2004; Martin et al, 2003; Morel et al, 2004; Ditzel et al, 2005), whereas other authors did not find such an increase (Viville et al, 2000; Vicari et al, 2002). However, there is an agreement in patients with large-headed spermatozoa with significantly higher incidence of sperm chromosomal abnormalities (In't Veld et al, 1997; Viville et al, 2000; Lewis-Jones et al, 2003; Vicari et al, 2003).

Our patient had a total teratozoospermia with almost all spermatozoa showing large head and multiple tails. There was clearly identified an isolated type of morphological abnormality, and FISH results for chromosomes X, Y, and 18 showed that 91.2% of the spermatozoa had chromosomal abnormalities. A high incidence of aneuploidy for these chromosomes and mainly hyperhaploid spermatozoa were also obtained by In't Veld (1997) and more recently by Lewis-Jones et al (2003). In our study, we also evaluated chromosomes 13 and 21, and with these results, we confirmed that the predominant abnormality found in the macrocephalic spermatozoa would be hyperhaploidy with aneuploidy for all the tested chromosomes. A low number of spermatozoa were evaluated due to the difficulty to discriminate between sperm heads and immature cells and only tailed spermatozoa were taken into account. Despite this consideration, we believe our results are as conclusive as other FISH studies in severe oligozoospermic, teratozoospermic, or azoospermic patients in whom a low number of spermatozoa were also evaluated and the results were statistically analyzed in the same manner (Levron et al, 2001; Lewis-Jones et al, 2003; Mateizel et al, 2002; Gianaroli et al, 2005).

To our knowledge, only 1 study reports the effect of the exposure to the chemicals in workers of a petroleum refinery on the sperm morphology (Rosemberg et al, 1985). They find a lack of association between abnormal morphology and the exposure to chemicals in this environment, without any mention of sperm chromosomal abnormalities.

The clinical consequences of using sperm samples with an abnormal FISH result in ICSI programs have been evaluated by several authors. It seems that sperm chromosomal abnormalities may adversely affect ICSI outcome in oligoasthenoteratozoospermic patients and in epididymal and testicular spermatozoa from azoospermic patients, decreasing fertilization (Pfeffer et al, 1999) and pregnancy rates (Pang et al, 1999; Pfeffer et al, 1999; Bernardini et al, 2000; Calogero et al, 2001b; Rubio et al, 2001) and increasing miscarriage rates (Rubio et al, 2001), at least in some cases. Moreover, Gianaroli et al (2000) and Silber et al (2003) analyzed the incidence of chromosomal abnormalities in embryos originated from azoospermic patients participating in a preimplantation genetic diagnosis program. Embryos from these patients suffered higher rates of abnormalities than those obtained

from normozoospermic or oligozoospermic patients, with high incidences of embryos with aneuploidies for sex chromosomes (Gianaroli et al, 2000) and mosaic embryos (Silber et al, 2003). Furthermore, Burrello et al (2004) have described that, in oligoasthenoteratozoospermic patients, there is an increased incidence of sperm chromosomal abnormalities in both types of spermatozoa, those with normal and abnormal head shape. Therefore, the selection of normal-shaped spermatozoa for ICSI would not prevent the risk of chromosomal abnormalities in the offspring.

Preimplantation genetic diagnosis has been offered by several groups as an alternative in patients at risk of chromosomal abnormalities in their spermatozoa (Arán et al, 1999; Gianaroli et al, 2000; Silber et al, 2003). However, in our patient, with nearly all spermatozoa being abnormal with only 5 out of 23 chromosomes analyzed, we discouraged a new ICSI attempt and therefore sperm donation was advised. The couple accepted and is currently in our donor insemination program.

References

- Alikani M, Cohen J, Tomkin G, Garrisi J, Mack C, Scot RT. Human embryo fragmentation in vitro and its implications for pregnancy and implantation. *Fertil Steril*. 1999;71:836–842.
- Arán B, Blanco J, Vidal F, Vendrell JM, Egozcue S, Barri PN, Egozcue J, Veiga A. Screening for abnormalities of chromosomes X, Y, and 18 and for diploidy in spermatozoa from infertile men participating in an in vitro fertilization-intracytoplasmic sperm injection program. *Fertil Steril*. 1999;72:696–701.
- Bernardini L, Borini A, Preti S, Conte N, Flamigni C, Capitanio GL, Venturini PL. Study of aneuploidy in normal and abnormal germ cells from semen of fertile and infertile men. *Hum Reprod*. 1998;13:3406–3413.
- Bernardini L, Gianaroli L, Fortini D, Conte N, Magli C, Cavani S, Gaggero G, Tindiglia C, Ragni N, Venturini PL. Frequency of hyper-, hypohaploidy and diploidy in ejaculate, epididymal and testicular germ cells of infertile patients. *Hum Reprod*. 2000;15:2165–2172.
- Burrello N, Arcidiacono G, Vicari E, Asero P, Di Benedetto D, De Palma A, Romeo R, D'Agata R, Calogero AE. Morphologically normal spermatozoa of patients with secretory oligo-astheno-teratozoospermia have an increased aneuploidy rate. *Hum Reprod*. 2004;19:2298–2302.
- Calogero AE, De Palma A, Grazioso C, Barone N, Romeo R, Rappazzo G, D'Agata R. High sperm aneuploidy rate in unselected infertile patients and its relationship with intracytoplasmic sperm injection outcome. *Hum Reprod*. 2001a;16:1433–1439.
- Calogero AE, De Palma A, Grazioso C, Barone N, Romeo R, Rappazzo G, D'Agata R. Aneuploidy rate in spermatozoa of selected men with abnormal semen parameters. *Hum Reprod*. 2001b;16:1172–1179.
- Carrell DT, Emery BR, Liu L. Characterization of aneuploidy rates, protamine levels, ultrastructure, and functional ability of round-headed sperm from two siblings and implications for intracytoplasmic sperm injection. *Fertil Steril*. 1999;71:511–516.
- Carrell DT, Emery BR, Wilcox AL, Campbell B, Erickson L, Hatasaka HH, Jones KP, Peterson CM. Sperm chromosome aneuploidy as related to male factor infertility and some ultrastructure defects. *Arch Androl*. 2004;50:181–185.
- Devillard F, Metzler-Guillemain C, Pelletier R, DeRobertis C, Bergues U, Hennebicq S, Guichaoua M, Sele B, Rousseaux S. Polyploidy in large-headed sperm: FISH study of three cases. *Hum Reprod*. 2002;17:1292–1298.
- Ditzel N, El-Danasouri I, Just W, Sterzik K. Higher aneuploidy rates of chromosomes 13, 16, and 21 in a patient with globozoospermia. *Fertil Steril*. 2005;84:217–218.
- Finkelstein S, Mukamel E, Yavetz H, Paz G, Avivi L. Increased rate of nondisjunction in sex cells derived from low-quality semen. *Hum Genet*. 1998;102:129–137.
- Gianaroli L, Magli MC, Cavallini G, Crippa A, Nadalini M, Bernardini L, Fabris GFM, Voliani S, Ferraretti AP. Frequency of aneuploidy in sperm from patients with extremely severe male factor infertility. *Hum Reprod*. 2005;20:2140–2152.
- Gianaroli L, Magli MC, Ferraretti AP, Iammarrone E. Preimplantation diagnosis after assisted reproduction techniques for genetically-determined male infertility. *J Endocrinol Invest*. 2000;23:711–716.
- Guttenbach M, Martinez-Exposito MJ, Michelmann HW, Engel W, Schmid M. Incidence of diploid and disomic sperm nuclei in 45 infertile men. *Hum Reprod*. 1997;12:468–473.
- In't Veld PA, Broekmans FJ, de France HF, Pearson PL, Pieters MH, van Kooij RJ. Intracytoplasmic sperm injection (ICSI) and chromosomally abnormal spermatozoa. *Hum Reprod*. 1997;12:752–754.
- Lähdele J, Saari N, Ajosenpää-Saari M, Mykkanen J. Incidence of aneuploid spermatozoa among infertile men studied by multicolour fluorescence in situ hybridization. *Am J Med Genet*. 1997;71:115–121.
- Lewis-Jones I, Aziz N, Seshadri S, Douglas A, Howard P. Sperm chromosomal abnormalities are linked to sperm morphologic deformities. *Fertil Steril*. 2003;79:212–215.
- Levron J, Aviram-Goldring A, Madgar I, Raviv G, Barkai G, Dor J. Sperm chromosome abnormalities in men with severe male factor infertility who are undergoing in vitro fertilization with intracytoplasmic sperm injection. *Fertil Steril*. 2001;76:479–484.
- Martin RH, Rademaker AW, Greene C, Ko E, Hoang T, Barclay L, Chernos J. A comparison of the frequency of sperm chromosome abnormalities in men with mild, moderate, and severe oligozoospermia. *Biol Reprod*. 2003;69:535–539.
- Mateizel I, Verheyen G, Van Assche E, Tournaye H, Liebaers I, Van Steirteghem A. FISH analysis of chromosome X, Y and 18 abnormalities in testicular sperm from azoospermic patients. *Hum Reprod*. 2002;17:2249–2257.
- McInnes B, Rademaker A, Greene CA, Ko E, Barclay L, Martin RH. Abnormalities for chromosomes 13 and 21 detected in spermatozoa from infertile men. *Hum Reprod*. 1998;13:2787–2790.
- Miharu N, Best RG, Young SR. Numerical chromosome abnormalities in spermatozoa of fertile and infertile men detected by fluorescence in situ hybridization. *Hum Genet*. 1994;93:502–506.
- Moosani N, Pattinson HA, Carter MD, Cox DM, Rademaker AW, Martin RH. Chromosomal analysis of sperm from men with idiopathic infertility using sperm karyotyping and fluorescence in situ hybridization. *Fertil Steril*. 1995;64:811–817.
- Morel F, Douet-Guilbert N, Moerman A, Duban B, Marchetti C, Delobel B, Le Bris M-J, Amice V, De Braekeleer M. Chromosome aneuploidy in the spermatozoa of two men with globozoospermia. *Mol Hum Reprod*. 2004;10:835–838.
- Nishikawa N, Murakami I, Ikuta K, Suzumori K. Sex chromosomal analysis of spermatozoa from infertile men using fluorescence in situ hybridization. *J Assist Reprod Genet*. 2000;17:97–102.
- Ohashi Y, Miharu N, Honda H, Samura O, Ohama K. High frequency of XY disomy in spermatozoa of severe oligozoospermic men. *Hum Reprod*. 2001;16:703–708.
- Pang MG, Hoegerman SF, Cuticchia AJ, Moon SY, Doncel GF, Acosta AA, Kearns WG. Detection of aneuploidy for chromosomes 4, 6, 7, 8, 9, 10, 11, 12, 13, 17, 18, 21, X and Y by fluorescence in-situ hybridization in spermatozoa from nine patients with oligoastheno-

- teratozoospermia undergoing intracytoplasmic sperm injection. *Hum Reprod.* 1999;14:1266–1273.
- Pfeffer J, Pang MG, Hoegerman SF, Osgood CJ, Stacey MW, Mayer J, Oehninger S, Kearns WG. Aneuploidy frequencies in semen fractions from ten oligoasthenoteratozoospermic patients donating sperm for intracytoplasmic sperm injection. *Fertil Steril.* 1999;72:472–478.
- Rives N, Saint Clair A, Mazurier S, Sibert L, Simeon N, Joly G, Mace B. Relationship between clinical phenotype, semen parameters and aneuploidy frequency in sperm nuclei of 50 infertile males. *Hum Genet.* 1999;105:266–272.
- Rodrigo L, Rubio C, Mateu E, Simon C, Remohi J, Pellicer A, Gil-Salom M. Analysis of chromosomal abnormalities in testicular and epididymal spermatozoa from azoospermic ICSI patients by fluorescence in-situ hybridization. *Hum Reprod.* 2004;19:118–123.
- Rosemberg MJ, Wyrobek AJ, Ratcliffe J, Gordon LA, Watchmaker G, Fox SH, Moore DH, Hornung RW. Sperm as an indicator of reproductive risk among petroleum refinery workers. *Br J Ind Med.* 1985;42:123–127.
- Rubio C, Gil-Salom M, Simon C, Vidal F, Rodrigo L, Minguez Y, Remohi J, Pellicer A. Incidence of sperm chromosomal abnormalities in a risk population: relationship with sperm quality and ICSI outcome. *Hum Reprod.* 2001;16:2084–2092.
- Silber S, Escudero T, Lenahan K, Abdelhadi I, Kilani Z, Munne S. Chromosomal abnormalities in embryos derived from testicular sperm extraction. *Fertil Steril.* 2003;79:30–38.
- Ushijima C, Kumasako Y, Kihai PE, Hirotsuru K, Utsunomiya T. Analysis of chromosomal abnormalities in human spermatozoa using multi-colour fluorescence in-situ hybridization. *Hum Reprod.* 2000;15:1107–1111.
- Vegetti W, Van Assche E, Frias A, Verheyen G, Bianchi MM, Bonduelle M, Liebaers I, Van Steirteghem A. Correlation between semen parameters and sperm aneuploidy rates investigated by fluorescence in-situ hybridization in infertile men. *Hum Reprod.* 2000;15:351–365.
- Vicari E, de Palma A, Burrello N, Longo G, Grazioso C, Barone N, Zahi M, D'Agata R, Calogero AE. Absolute polymorphic teratozoospermia in patients with oligo-asthenozoospermia is associated with an elevated sperm aneuploidy rate. *J Androl.* 2003;24:598–603.
- Vicari E, Perdichizzi A, De Palma A, Burrello N, D'Agata R, Calogero AE. Globozoospermia is associated with chromatin structure abnormalities: case report. *Hum Reprod.* 2002;17:2128–2133.
- Viville S, Mollard R, Bach ML, Falquet C, Gerlinger P, Warter S. Do morphological anomalies reflect chromosomal aneuploidies?: case report. *Hum Reprod.* 2000;15:2563–2566.
- World Health Organization. *WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction*, 4th ed. World Health Organization. Cambridge, United Kingdom: Cambridge University Press; 1999.
- Yurov YB, Saia MJ, Vorsanoya SG, Emy R, Soloviev IV, Sharonin VO, Guichaoua MR, Luciani JM. Rapid chromosomal analysis of germline cells by FISH: an investigation of an infertile male with large-headed spermatozoa. *Mol Hum Reprod.* 1996;2:665–658.