



Search for the decay $B_s^0 \rightarrow D^{*\mp} \pi^\pm$

The LHCb collaboration[†]

Abstract

A search for the decay $B_s^0 \rightarrow D^{*\mp} \pi^\pm$ is presented using a data sample corresponding to an integrated luminosity of 1.0 fb^{-1} of pp collisions collected by LHCb. This decay is expected to be mediated by a W -exchange diagram, with little contribution from rescattering processes, and therefore a measurement of the branching fraction will help to understand the mechanism behind related decays such as $B_s^0 \rightarrow \pi^+ \pi^-$ and $B_s^0 \rightarrow D \bar{D}$. Systematic uncertainties are minimised by using $B^0 \rightarrow D^{*\mp} \pi^\pm$ as a normalisation channel. We find no evidence for a signal, and set an upper limit on the branching fraction of $\mathcal{B}(B_s^0 \rightarrow D^{*\mp} \pi^\pm) < 6.1 (7.8) \times 10^{-6}$ at 90 % (95 %) confidence level.

Submitted to Phys. Rev. D (R)

[†]Authors are listed on the following pages.

LHCb collaboration

R. Aaij⁴⁰, C. Abellan Beteta^{35,n}, B. Adeva³⁶, M. Adinolfi⁴⁵, C. Adrover⁶, A. Affolder⁵¹, Z. Ajaltouni⁵, J. Albrecht⁹, F. Alessio³⁷, M. Alexander⁵⁰, S. Ali⁴⁰, G. Alkhazov²⁹, P. Alvarez Cartelle³⁶, A.A. Alves Jr^{24,37}, S. Amato², S. Amerio²¹, Y. Amhis⁷, L. Anderlini^{17,f}, J. Anderson³⁹, R. Andreassen⁵⁹, R.B. Appleby⁵³, O. Aquines Gutierrez¹⁰, F. Archilli¹⁸, A. Artamonov³⁴, M. Artuso⁵⁶, E. Aslanides⁶, G. Auriemma^{24,m}, S. Bachmann¹¹, J.J. Back⁴⁷, C. Baesso⁵⁷, V. Balagura³⁰, W. Baldini¹⁶, R.J. Barlow⁵³, C. Barschel³⁷, S. Barsuk⁷, W. Barter⁴⁶, Th. Bauer⁴⁰, A. Bay³⁸, J. Beddow⁵⁰, F. Bedeschi²², I. Bediaga¹, S. Belogurov³⁰, K. Belous³⁴, I. Belyaev³⁰, E. Ben-Haim⁸, M. Benayoun⁸, G. Bencivenni¹⁸, S. Benson⁴⁹, J. Benton⁴⁵, A. Berezhnoy³¹, R. Bernet³⁹, M.-O. Bettler⁴⁶, M. van Beuzekom⁴⁰, A. Bien¹¹, S. Bifani¹², T. Bird⁵³, A. Bizzeti^{17,h}, P.M. Bjørnstad⁵³, T. Blake³⁷, F. Blanc³⁸, J. Blouw¹¹, S. Blusk⁵⁶, V. Bocci²⁴, A. Bondar³³, N. Bondar²⁹, W. Bonivento¹⁵, S. Borghi⁵³, A. Borgia⁵⁶, T.J.V. Bowcock⁵¹, E. Bowen³⁹, C. Bozzi¹⁶, T. Brambach⁹, J. van den Brand⁴¹, J. Bressieux³⁸, D. Brett⁵³, M. Britsch¹⁰, T. Britton⁵⁶, N.H. Brook⁴⁵, H. Brown⁵¹, I. Burducea²⁸, A. Bursche³⁹, G. Busetto^{21,q}, J. Buytaert³⁷, S. Cadeddu¹⁵, O. Callot⁷, M. Calvi^{20,j}, M. Calvo Gomez^{35,n}, A. Camboni³⁵, P. Campana^{18,37}, A. Carbone^{14,c}, G. Carboni^{23,k}, R. Cardinale^{19,i}, A. Cardini¹⁵, H. Carranza-Mejia⁴⁹, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵¹, M. Cattaneo³⁷, Ch. Cauet⁹, M. Charles⁵⁴, Ph. Charpentier³⁷, P. Chen^{3,38}, N. Chiapolini³⁹, M. Chrzaszcz²⁵, K. Ciba³⁷, X. Cid Vidal³⁶, G. Ciezarek⁵², P.E.L. Clarke⁴⁹, M. Clemencic³⁷, H.V. Cliff⁴⁶, J. Closier³⁷, C. Coca²⁸, V. Coco⁴⁰, J. Cogan⁶, E. Cogneras⁵, P. Collins³⁷, A. Comerma-Montells³⁵, A. Contu¹⁵, A. Cook⁴⁵, M. Coombes⁴⁵, S. Coquereau⁸, G. Corti³⁷, B. Couturier³⁷, G.A. Cowan³⁸, D. Craik⁴⁷, S. Cunliffe⁵², R. Currie⁴⁹, C. D'Ambrosio³⁷, P. David⁸, P.N.Y. David⁴⁰, I. De Bonis⁴, K. De Bruyn⁴⁰, S. De Capua⁵³, M. De Cian³⁹, J.M. De Miranda¹, M. De Oyanguren Campos^{35,o}, L. De Paula², W. De Silva⁵⁹, P. De Simone¹⁸, D. Decamp⁴, M. Deckenhoff⁹, L. Del Buono⁸, D. Derkach¹⁴, O. Deschamps⁵, F. Dettori⁴¹, A. Di Canto¹¹, H. Dijkstra³⁷, M. Dogaru²⁸, S. Donleavy⁵¹, F. Dordei¹¹, A. Dosil Suárez³⁶, D. Dossett⁴⁷, A. Dovbnya⁴², F. Dupertuis³⁸, R. Dzhelyadin³⁴, A. Dziurda²⁵, A. Dzyuba²⁹, S. Easo^{48,37}, U. Egede⁵², V. Egorychev³⁰, S. Eidelman³³, D. van Eijk⁴⁰, S. Eisenhardt⁴⁹, U. Eitschberger⁹, R. Ekelhof⁹, L. Eklund⁵⁰, I. El Rifai⁵, Ch. Elsasser³⁹, D. Elsby⁴⁴, A. Falabella^{14,e}, C. Färber¹¹, G. Fardell⁴⁹, C. Farinelli⁴⁰, S. Farry¹², V. Fave³⁸, D. Ferguson⁴⁹, V. Fernandez Albor³⁶, F. Ferreira Rodrigues¹, M. Ferro-Luzzi³⁷, S. Filippov³², C. Fitzpatrick³⁷, M. Fontana¹⁰, F. Fontanelli^{19,i}, R. Forty³⁷, O. Francisco², M. Frank³⁷, C. Frei³⁷, M. Frosini^{17,f}, S. Furcas²⁰, E. Furfaro²³, A. Gallas Torreira³⁶, D. Galli^{14,c}, M. Gandelman², P. Gandini⁵⁴, Y. Gao³, J. Garofoli⁵⁶, P. Garosi⁵³, J. Garra Tico⁴⁶, L. Garrido³⁵, C. Gaspar³⁷, R. Gauld⁵⁴, E. Gersabeck¹¹, M. Gersabeck⁵³, T. Gershon^{47,37}, Ph. Ghez⁴, V. Gibson⁴⁶, V.V. Gligorov³⁷, C. Göbel⁵⁷, D. Golubkov³⁰, A. Golutvin^{52,30,37}, A. Gomes², H. Gordon⁵⁴, M. Grabalosa Gándara⁵, R. Graciani Diaz³⁵, L.A. Granado Cardoso³⁷, E. Graugés³⁵, G. Graziani¹⁷, A. Grecu²⁸, E. Greening⁵⁴, S. Gregson⁴⁶, O. Grünberg⁵⁸, B. Gui⁵⁶, E. Gushchin³², Yu. Guz³⁴, T. Gys³⁷, C. Hadjivasiliou⁵⁶, G. Haefeli³⁸, C. Haen³⁷, S.C. Haines⁴⁶, S. Hall⁵², T. Hampson⁴⁵, S. Hansmann-Menzemer¹¹, N. Harnew⁵⁴, S.T. Harnew⁴⁵, J. Harrison⁵³, T. Hartmann⁵⁸, J. He⁷, V. Heijne⁴⁰, K. Hennessy⁵¹, P. Henrard⁵, J.A. Hernando Morata³⁶, E. van Herwijnen³⁷, E. Hicks⁵¹, D. Hill⁵⁴, M. Hoballah⁵, C. Hombach⁵³, P. Hopchev⁴, W. Hulsbergen⁴⁰, P. Hunt⁵⁴, T. Huse⁵¹, N. Hussain⁵⁴, D. Hutchcroft⁵¹, D. Hynds⁵⁰, V. Iakovenko⁴³, M. Idzik²⁶, P. Ilten¹², R. Jacobsson³⁷, A. Jaeger¹¹, E. Jans⁴⁰, P. Jaton³⁸, F. Jing³, M. John⁵⁴, D. Johnson⁵⁴,

C.R. Jones⁴⁶, B. Jost³⁷, M. Kabbalo⁹, S. Kandybei⁴², M. Karacson³⁷, T.M. Karbach³⁷,
 I.R. Kenyon⁴⁴, U. Kerzel³⁷, T. Ketel⁴¹, A. Keune³⁸, B. Khanji²⁰, O. Kochebina⁷,
 I. Komarov^{38,31}, R.F. Koopman⁴¹, P. Koppenburg⁴⁰, M. Korolev³¹, A. Kozlinskiy⁴⁰,
 L. Kravchuk³², K. Kreplin¹¹, M. Kreps⁴⁷, G. Krocker¹¹, P. Krokovny³³, F. Kruse⁹,
 M. Kucharczyk^{20,25,j}, V. Kudryavtsev³³, T. Kvaratskheliya^{30,37}, V.N. La Thi³⁸, D. Lacarrere³⁷,
 G. Lafferty⁵³, A. Lai¹⁵, D. Lambert⁴⁹, R.W. Lambert⁴¹, E. Lanciotti³⁷, G. Lanfranchi^{18,37},
 C. Langenbruch³⁷, T. Latham⁴⁷, C. Lazzeroni⁴⁴, R. Le Gac⁶, J. van Leerdam⁴⁰, J.-P. Lees⁴,
 R. Lefèvre⁵, A. Leflat^{31,37}, J. Lefrançois⁷, S. Leo²², O. Leroy⁶, B. Leverington¹¹, Y. Li³,
 L. Li Gioi⁵, M. Liles⁵¹, R. Lindner³⁷, C. Linn¹¹, B. Liu³, G. Liu³⁷, J. von Loeben²⁰, S. Lohn³⁷,
 J.H. Lopes², E. Lopez Asamar³⁵, N. Lopez-March³⁸, H. Lu³, D. Lucchesi^{21,q}, J. Luisier³⁸,
 H. Luo⁴⁹, F. Machefert⁷, I.V. Machikhiliyan^{4,30}, F. Maciuc²⁸, O. Maev^{29,37}, S. Malde⁵⁴,
 G. Manca^{15,d}, G. Mancinelli⁶, U. Marconi¹⁴, R. Märki³⁸, J. Marks¹¹, G. Martellotti²⁴,
 A. Martens⁸, L. Martin⁵⁴, A. Martín Sánchez⁷, M. Martinelli⁴⁰, D. Martinez Santos⁴¹,
 D. Martins Tostes², A. Massafferri¹, R. Matev³⁷, Z. Mathe³⁷, C. Matteuzzi²⁰, E. Maurice⁶,
 A. Mazurov^{16,32,37,e}, J. McCarthy⁴⁴, R. McNulty¹², A. McNab⁵³, B. Meadows^{59,54}, F. Meier⁹,
 M. Meissner¹¹, M. Merk⁴⁰, D.A. Milanes⁸, M.-N. Minard⁴, J. Molina Rodriguez⁵⁷, S. Monteil⁵,
 D. Moran⁵³, P. Morawski²⁵, M.J. Morello^{22,s}, R. Mountain⁵⁶, I. Mous⁴⁰, F. Muheim⁴⁹,
 K. Müller³⁹, R. Muresan²⁸, B. Muryn²⁶, B. Muster³⁸, P. Naik⁴⁵, T. Nakada³⁸,
 R. Nandakumar⁴⁸, I. Nasteva¹, M. Needham⁴⁹, N. Neufeld³⁷, A.D. Nguyen³⁸, T.D. Nguyen³⁸,
 C. Nguyen-Mau^{38,p}, M. Nicol⁷, V. Niess⁵, R. Niet⁹, N. Nikitin³¹, T. Nikodem¹¹,
 A. Nomerotski⁵⁴, A. Novoselov³⁴, A. Oblakowska-Mucha²⁶, V. Obraztsov³⁴, S. Oggero⁴⁰,
 S. Ogilvy⁵⁰, O. Okhrimenko⁴³, R. Oldeman^{15,d,37}, M. Orlandea²⁸, J.M. Otalora Goicochea²,
 P. Owen⁵², B.K. Pal⁵⁶, A. Palano^{13,b}, M. Palutan¹⁸, J. Panman³⁷, A. Papanestis⁴⁸,
 M. Pappagallo⁵⁰, C. Parkes⁵³, C.J. Parkinson⁵², G. Passaleva¹⁷, G.D. Patel⁵¹, M. Patel⁵²,
 G.N. Patrick⁴⁸, C. Patrignani^{19,i}, C. Pavel-Nicorescu²⁸, A. Pazos Alvarez³⁶, A. Pellegrino⁴⁰,
 G. Penso^{24,l}, M. Pepe Altarelli³⁷, S. Perazzini^{14,c}, D.L. Perego^{20,j}, E. Perez Trigo³⁶,
 A. Pérez-Calero Yzquierdo³⁵, P. Perret⁵, M. Perrin-Terrin⁶, G. Pessina²⁰, K. Petridis⁵²,
 A. Petrolini^{19,i}, A. Phan⁵⁶, E. Picatoste Olloqui³⁵, B. Pietrzyk⁴, T. Pilař⁴⁷, D. Pinci²⁴,
 S. Playfer⁴⁹, M. Plo Casasus³⁶, F. Polci⁸, G. Polok²⁵, A. Poluektov^{47,33}, E. Polcarpo²,
 D. Popov¹⁰, B. Popovici²⁸, C. Potterat³⁵, A. Powell⁵⁴, J. Prisciandaro³⁸, V. Pugatch⁴³,
 A. Puig Navarro³⁸, G. Punzi^{22,r}, W. Qian⁴, J.H. Rademacker⁴⁵, B. Rakotomiamanana³⁸,
 M.S. Rangel², I. Raniuk⁴², N. Rauschmayr³⁷, G. Raven⁴¹, S. Redford⁵⁴, M.M. Reid⁴⁷,
 A.C. dos Reis¹, S. Ricciardi⁴⁸, A. Richards⁵², K. Rinnert⁵¹, V. Rives Molina³⁵,
 D.A. Roa Romero⁵, P. Robbe⁷, E. Rodrigues⁵³, P. Rodriguez Perez³⁶, S. Roiser³⁷,
 V. Romanovsky³⁴, A. Romero Vidal³⁶, J. Rouvinet³⁸, T. Ruf³⁷, F. Ruffini²², H. Ruiz³⁵,
 P. Ruiz Valls^{35,o}, G. Sabatino^{24,k}, J.J. Saborido Silva³⁶, N. Sagidova²⁹, P. Sail⁵⁰, B. Saitta^{15,d},
 C. Salzmann³⁹, B. Sanmartin Sedes³⁶, M. Sannino^{19,i}, R. Santacesaria²⁴,
 C. Santamarina Rios³⁶, E. Santovetti^{23,k}, M. Sapunov⁶, A. Sarti^{18,l}, C. Satriano^{24,m},
 A. Satta²³, M. Savrie^{16,e}, D. Savrina^{30,31}, P. Schaack⁵², M. Schiller⁴¹, H. Schindler³⁷,
 M. Schlupp⁹, M. Schmelling¹⁰, B. Schmidt³⁷, O. Schneider³⁸, A. Schopper³⁷, M.-H. Schune⁷,
 R. Schwemmer³⁷, B. Sciascia¹⁸, A. Sciubba²⁴, M. Seco³⁶, A. Semennikov³⁰, K. Senderowska²⁶,
 I. Sepp⁵², N. Serra³⁹, J. Serrano⁶, P. Seyfert¹¹, M. Shapkin³⁴, I. Shapoval^{42,37}, P. Shatalov³⁰,
 Y. Shcheglov²⁹, T. Shears^{51,37}, L. Shekhtman³³, O. Shevchenko⁴², V. Shevchenko³⁰,
 A. Shires⁵², R. Silva Coutinho⁴⁷, T. Skwarnicki⁵⁶, N.A. Smith⁵¹, E. Smith^{54,48}, M. Smith⁵³,
 M.D. Sokoloff⁵⁹, F.J.P. Soler⁵⁰, F. Soomro^{18,37}, D. Souza⁴⁵, B. Souza De Paula², B. Spaan⁹,
 A. Sparkes⁴⁹, P. Spradlin⁵⁰, F. Stagni³⁷, S. Stahl¹¹, O. Steinkamp³⁹, S. Stoica²⁸, S. Stone⁵⁶,

B. Storaci³⁹, M. Straticiu²⁸, U. Straumann³⁹, V.K. Subbiah³⁷, S. Swientek⁹, V. Syropoulos⁴¹, M. Szczekowski²⁷, P. Szczypka^{38,37}, T. Szumlak²⁶, S. T'Jampens⁴, M. Teklishyn⁷, E. Teodorescu²⁸, F. Teubert³⁷, C. Thomas⁵⁴, E. Thomas³⁷, J. van Tilburg¹¹, V. Tisserand⁴, M. Tobin³⁹, S. Tolk⁴¹, D. Tonelli³⁷, S. Topp-Joergensen⁵⁴, N. Torr⁵⁴, E. Tournefier^{4,52}, S. Tourneur³⁸, M.T. Tran³⁸, M. Tresch³⁹, A. Tsaregorodtsev⁶, P. Tsopelas⁴⁰, N. Tuning⁴⁰, M. Ubeda Garcia³⁷, A. Ukleja²⁷, D. Urner⁵³, U. Uwer¹¹, V. Vagnoni¹⁴, G. Valenti¹⁴, R. Vazquez Gomez³⁵, P. Vazquez Regueiro³⁶, S. Vecchi¹⁶, J.J. Velthuis⁴⁵, M. Veltri^{17,g}, G. Veneziano³⁸, M. Vesterinen³⁷, B. Viaud⁷, D. Vieira², X. Vilasis-Cardona^{35,n}, A. Vollhardt³⁹, D. Volyanskyy¹⁰, D. Voong⁴⁵, A. Vorobyev²⁹, V. Vorobyev³³, C. Voß⁵⁸, H. Voss¹⁰, R. Waldi⁵⁸, R. Wallace¹², S. Wandernoth¹¹, J. Wang⁵⁶, D.R. Ward⁴⁶, N.K. Watson⁴⁴, A.D. Webber⁵³, D. Websdale⁵², M. Whitehead⁴⁷, J. Wicht³⁷, J. Wiechczynski²⁵, D. Wiedner¹¹, L. Wiggers⁴⁰, G. Wilkinson⁵⁴, M.P. Williams^{47,48}, M. Williams⁵⁵, F.F. Wilson⁴⁸, J. Wishahi⁹, M. Witek²⁵, S.A. Wotton⁴⁶, S. Wright⁴⁶, S. Wu³, K. Wyllie³⁷, Y. Xie^{49,37}, F. Xing⁵⁴, Z. Xing⁵⁶, Z. Yang³, R. Young⁴⁹, X. Yuan³, O. Yushchenko³⁴, M. Zangoli¹⁴, M. Zavertyaev^{10,a}, F. Zhang³, L. Zhang⁵⁶, W.C. Zhang¹², Y. Zhang³, A. Zhelezov¹¹, A. Zhokhov³⁰, L. Zhong³, A. Zvyagin³⁷.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

⁷LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹⁰Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹¹Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹²School of Physics, University College Dublin, Dublin, Ireland

¹³Sezione INFN di Bari, Bari, Italy

¹⁴Sezione INFN di Bologna, Bologna, Italy

¹⁵Sezione INFN di Cagliari, Cagliari, Italy

¹⁶Sezione INFN di Ferrara, Ferrara, Italy

¹⁷Sezione INFN di Firenze, Firenze, Italy

¹⁸Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

¹⁹Sezione INFN di Genova, Genova, Italy

²⁰Sezione INFN di Milano Bicocca, Milano, Italy

²¹Sezione INFN di Padova, Padova, Italy

²²Sezione INFN di Pisa, Pisa, Italy

²³Sezione INFN di Roma Tor Vergata, Roma, Italy

²⁴Sezione INFN di Roma La Sapienza, Roma, Italy

²⁵Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

²⁶AGH University of Science and Technology, Kraków, Poland

²⁷National Center for Nuclear Research (NCBJ), Warsaw, Poland

²⁸Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

²⁹Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

³⁰Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia

³¹Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia

³²Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia

³³Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia

³⁴Institute for High Energy Physics (IHEP), Protvino, Russia

- ³⁵ *Universitat de Barcelona, Barcelona, Spain*
- ³⁶ *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- ³⁷ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ³⁸ *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ³⁹ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴⁰ *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- ⁴¹ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- ⁴² *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴³ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁴ *University of Birmingham, Birmingham, United Kingdom*
- ⁴⁵ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁶ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁴⁷ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁴⁸ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁴⁹ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵⁰ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵¹ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵² *Imperial College London, London, United Kingdom*
- ⁵³ *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁴ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁵ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁵⁶ *Syracuse University, Syracuse, NY, United States*
- ⁵⁷ *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁵⁸ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹¹*
- ⁵⁹ *University of Cincinnati, Cincinnati, OH, United States, associated to ⁵⁶*

^a *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*

^b *Università di Bari, Bari, Italy*

^c *Università di Bologna, Bologna, Italy*

^d *Università di Cagliari, Cagliari, Italy*

^e *Università di Ferrara, Ferrara, Italy*

^f *Università di Firenze, Firenze, Italy*

^g *Università di Urbino, Urbino, Italy*

^h *Università di Modena e Reggio Emilia, Modena, Italy*

ⁱ *Università di Genova, Genova, Italy*

^j *Università di Milano Bicocca, Milano, Italy*

^k *Università di Roma Tor Vergata, Roma, Italy*

^l *Università di Roma La Sapienza, Roma, Italy*

^m *Università della Basilicata, Potenza, Italy*

ⁿ *LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*

^o *IFIC, Universitat de Valencia-CSIC, Valencia, Spain*

^p *Hanoi University of Science, Hanoi, Viet Nam*

^q *Università di Padova, Padova, Italy*

^r *Università di Pisa, Pisa, Italy*

^s *Scuola Normale Superiore, Pisa, Italy*

Decays of B_s^0 mesons to final states such as D^+D^- , $D^0\bar{D}^0$ [1] and $\pi^+\pi^-$ [2] have been recently observed by LHCb. Such decays can proceed, at short distance, by two types of amplitudes, referred to as weak exchange and penguin annihilation. Example diagrams are shown in Fig. 1(a) and (b). There is also a potential long distance contribution from rescattering. For example, the D^+D^- final state can be obtained from a $b \rightarrow c\bar{c}s$ decay to $D_s^+D_s^-$ followed by the $s\bar{s}$ pair rearranging to $d\bar{d}$. Understanding rescattering effects in hadronic B meson decays is important in order to interpret various CP -violating observables.




Figure 1: Decay diagrams for (a) $B_s^0 \rightarrow D^{(*)+}D^{(*)-}$ via weak exchange, (b) $B_s^0 \rightarrow D^{(*)+}D^{(*)-}$ via penguin annihilation, (c) $B_s^0 \rightarrow D^{(*)-}\pi^+$ via weak exchange.

A measurement of the branching fraction of the decay $B_s^0 \rightarrow D^{*-}\pi^+$ can be used to disentangle the contributions from different decay diagrams and from rescattering [3, 4]. This decay has only weak exchange contributions, as shown in Fig. 1(c). (The suppressed diagram for $B_s^0 \rightarrow D^{*+}\pi^-$ is not shown.) Moreover, rescattering contributions to the $B_s^0 \rightarrow D^{(*)\mp}\pi^\pm$ decay are expected to be small [5]. Therefore, if the observed branching fraction for the decay $B_s^0 \rightarrow \pi^+\pi^-$ is explained by rescattering, a low value of $\mathcal{B}(B_s^0 \rightarrow D^{*-}\pi^+) = (1.2 \pm 0.2) \times 10^{-6}$ is predicted [5]. However, if short-distance amplitudes are the dominant effect in $B_s^0 \rightarrow \pi^+\pi^-$ and related decays, $\mathcal{B}(B_s^0 \rightarrow D^{*-}\pi^+)$ could be much larger. The measured $B_s^0 \rightarrow D\bar{D}$ [1] and $B^+ \rightarrow D_s^+\phi$ [6] rates are at the upper end of the expected range in the rescattering-based model, but further measurements are needed to establish whether long-distance processes are dominant in these hadronic B decays.

In this paper, the result of a search for the decay $B_s^0 \rightarrow D^{*\mp}\pi^\pm$ is presented. No previous measurements of this decay have been made. The inclusion of charge conjugated processes is implied throughout the paper. Since the flavour of the B_s^0 meson at production is not tagged, the $D^{*-}\pi^+$ and $D^{*+}\pi^-$ final states are combined. The analysis is based on a data sample corresponding to an integrated luminosity of 1.0fb^{-1} of LHC pp collision data, at a centre-of-mass energy of 7 TeV, collected with the LHCb detector during 2011. In high energy pp collisions all b hadron species are produced, so the $B^0 \rightarrow D^{*\mp}\pi^\pm$ decay, with branching fraction $\mathcal{B}(B^0 \rightarrow D^{*-}\pi^+) = (2.76 \pm 0.13) \times 10^{-3}$ [7, 8], is both a potentially serious background channel as well as the ideal normalisation mode for the measurement of the B_s^0 branching fraction.

The LHCb detector [9] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region, a large-area silicon-strip detector

located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes placed downstream. The combined tracking system has momentum resolution $\Delta p/p$ that varies from 0.4 % at 5 GeV/c to 0.6 % at 100 GeV/c, and impact parameter (IP) resolution of 20 μm for tracks with high transverse momentum (p_{T}). Charged hadrons are identified using two ring-imaging Cherenkov detectors. Photon, electron and hadron candidates are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers.

The trigger [10] consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage which applies a full event reconstruction. In this analysis, signal candidates are accepted if one of the final state particles created a cluster in the calorimeter with sufficient transverse energy to fire the hardware trigger. Events that are triggered at the hardware level by another particle in the $pp \rightarrow b\bar{b}X$ event are also retained. The software trigger requires characteristic signatures of b -hadron decays: at least one track, with $p_{\text{T}} > 1.7$ GeV/c and χ_{IP}^2 with respect to any primary interaction vertex (PV) greater than 16, that subsequently forms a two-, three- or four-track secondary vertex with a high sum of the p_{T} of the tracks and significant displacement from the PV. The χ_{IP}^2 is the difference between the χ^2 of the PV reconstruction with and without the considered track. In the offline analysis, the software trigger decision is required to be due to the candidate signal decay.

Candidates that are consistent with the decay chain $B_{(s)}^0 \rightarrow D^{*\mp}\pi^{\pm}$, $D^{*-} \rightarrow \bar{D}^0\pi^{-}$, $\bar{D}^0 \rightarrow K^+\pi^{-}$ are selected. The \bar{D}^0 and D^{*-} candidate invariant masses are required to satisfy $1814 < m_{K^+\pi^-} < 1914$ MeV/c² and $2008.78 < m_{\bar{D}^0\pi^-} < 2011.78$ MeV/c², respectively, where a D^0 mass constraint is applied in the evaluation of $m_{\bar{D}^0\pi^-}$. The bachelor pion, from the $B_{(s)}^0$ decay, is required to be consistent with the pion mass hypothesis, based on particle identification (PID) information from the RICH detectors [11]. All other selection criteria were tuned on the $B^0 \rightarrow D^{*\mp}\pi^{\pm}$ control channel in a similar manner to that used in another recent LHCb publication [12]. The large yield in the normalisation sample allows the selection to be based on data, though the efficiencies are determined using Monte Carlo (MC) simulated events in which pp collisions are generated using PYTHIA 6.4 [13] with a specific LHCb configuration [14]. Decays of hadronic particles are described by EVTGEN [15]. The interaction of the generated particles with the detector and its response are implemented using the GEANT4 toolkit [16] as described in Ref. [17].

The selection requirements include criteria on the quality of the tracks forming the signal candidate, their p , p_{T} and inconsistency with the hypothesis of originating from the PV (χ_{IP}^2). Requirements are also placed on the corresponding variables for candidate composite particles (\bar{D}^0 , $B_{(s)}^0$) together with restrictions of the decay fit (χ_{vertex}^2), the flight distance (χ_{flight}^2), and the cosine of the angle between the momentum vector and the line joining the PV to the $B_{(s)}^0$ vertex ($\cos\theta_{\text{dir}}$) [18].

Further discrimination between signal and background categories is achieved by calculating weights for the remaining B^0 candidates [19]. The weights are based on a simplified

fit to the B candidate invariant mass distribution, where the B_s^0 region is neither examined nor included in the fit. The weights are used to train a neural network [20] in order to maximise the separation between categories. To retain sufficient background events for the network training, the requirement on $m_{\bar{D}^0\pi^-}$ is not applied. A total of fifteen variables are used as input to the network. They include the χ_{IP}^2 of the four candidate tracks, the χ_{IP}^2 , χ_{vertex}^2 , χ_{flight}^2 and $\cos\theta_{\text{dir}}$ of the \bar{D}^0 and $B_{(s)}^0$ candidates, and the $B_{(s)}^0$ candidate p_{T} . The p_{T} asymmetry and track multiplicity in a cone with half-angle of 1.5 units in the plane of pseudorapidity and azimuthal angle (measured in radians) [21] around the $B_{(s)}^0$ candidate flight direction are also used. The input quantities to the neural network depend only weakly on the kinematics of the $B_{(s)}^0$ decay. A requirement on the network output is imposed that reduces the combinatorial background by an order of magnitude while retaining about 75% of the signal. Potential biases from this data-driven method are investigated by training the neural network with different fractions of the data sample. The same results are obtained using a neural network trained on 30, 40, 50, 60 and 70% of the total data sample.

After all selection requirements are applied, approximately 50 000 candidates are selected in the invariant mass range $5150 < m_{D^{*-}\pi^+} < 5600 \text{ MeV}/c^2$. About 1% of events with at least one candidate also contain a second candidate. Such multiple candidates are retained and treated the same as other candidates.

In addition to combinatorial background, candidates may be formed from misidentified or partially reconstructed $B_{(s)}^0$ decays. Contributions from partially reconstructed decays are reduced by requiring the invariant mass of the $B_{(s)}^0$ candidate to be above $5150 \text{ MeV}/c^2$. The contribution from $B_{(s)}^0$ decays to identical final states but without intermediate charmed mesons is negligible due to the requirement on the D^{*-} candidate invariant mass. A small but significant number of background events are expected from $B^0 \rightarrow D^{*-}K^+$ decays with the K^+ misidentified as a pion. The branching fractions of $\bar{B}_s^0 \rightarrow D^{*-}K^+$ and $A_b^0 \rightarrow D^{*-}p$ are expected to be small due to CKM suppression, so that these potential backgrounds are negligible.

Since the B^0 decay mode is several orders of magnitude more abundant than the B_s^0 decay, it is critical to understand precisely the shape of the B^0 signal peak. The dependence of the width of the peak on different kinematic variables of the B^0 decay was investigated. The strongest correlation was found to be with the angle between the momenta of the D^{*-} candidate and the bachelor π^+ in the lab frame, denoted as θ_{bach} . Simulated pseudo-experiments were used to find an optimal number of θ_{bach} bins to be used in a simultaneous fit. The outcome is that five bins are used, with ranges 0–0.046, 0.046–0.067, 0.067–0.092, 0.092–0.128 and 0.128–0.4 rad, chosen to have approximately equal numbers of B^0 decays in each. The peak width in the highest bin is approximately 60% of that in the lowest bin. The pseudo-experiments show that the simultaneous fit in bins of θ_{bach} is approximately 20% more sensitive to a potential B_s^0 signal than the fit without binning.

The signal yields are obtained from a maximum likelihood fit to the $D^{*-}\pi^+$ invariant mass distribution in the range 5150–5600 MeV/c^2 . The fit is performed simultaneously in

the five θ_{bach} bins. The fit includes double Gaussian shapes, where the two Gaussian functions share a common mean, for B^0 and B_s^0 signals, together with an exponential component for the partially reconstructed background, a linear component for the combinatorial background and a non-parametric function, derived from simulation, for $B^0 \rightarrow D^{*-}K^+$ decays. The probability density function (PDF) for the $B^0 \rightarrow D^{*-}K^+$ background is shifted by the mass difference between data and simulation for each bin of θ_{bach} .

The parameters of the double Gaussian shapes are constrained to be identical for B^0 and B_s^0 signals, with an offset in their mean values fixed to the known B^0 - B_s^0 mass difference [8]. Additionally, the relative normalisation of the two Gaussian functions and the ratio of their widths are constrained within uncertainties to the value obtained in simulation. A total of thirty-three parameters are allowed to vary in the fit: the ratio of yields $N(B_s^0)/N(B^0)$, the linear slope of the combinatorial background and the exponential parameter of the partially reconstructed background, plus separate parameters in each of the θ_{bach} bins to describe the peak position and core Gaussian width of the signal PDF, and the yields of the B^0 peak, the combinatorial background, the partially reconstructed background, and the background from $B^0 \rightarrow D^{*-}K^+$.

The results of the fit are shown in Fig. 2. The total number of $B^0 \rightarrow D^{*\mp}\pi^\pm$ decays is found to be $30\,000 \pm 400$, and the ratio of yields is determined to be $N(B_s^0)/N(B^0) = (1.4 \pm 3.5) \times 10^{-4}$, where the uncertainty is statistical only. The number of $B^0 \rightarrow D^{*-}K^+$ decays found is $1\,200 \pm 200$, with a correlation of 7% to the ratio of signal yields.

The ratio of yields is converted to a branching fraction following

$$\mathcal{B}(B_s^0 \rightarrow D^{*\mp}\pi^\pm) = \frac{N(B_s^0)}{N(B^0)} \times \frac{\epsilon(B^0)}{\epsilon(B_s^0)} \times \frac{f_d}{f_s} \times \mathcal{B}(B^0 \rightarrow D^{*-}\pi^+), \quad (1)$$

where $\epsilon(B^0)$ and $\epsilon(B_s^0)$ are the efficiencies for the B^0 and B_s^0 decay modes respectively, while f_d (f_s) is the probability that a b quark produced in the acceptance results in a B^0 (B_s^0) meson. Their ratio has been determined to be $f_s/f_d = 0.256 \pm 0.020$ [22].

The total efficiencies are $(0.165 \pm 0.002)\%$ and $(0.162 \pm 0.002)\%$ for the B^0 and B_s^0 decay modes respectively, including contributions from detector acceptance, selection criteria, PID and trigger effects. The ratio is consistent with unity, as expected. The PID efficiency is measured using a control sample of $D^{*-} \rightarrow \bar{D}^0\pi^-$, $\bar{D}^0 \rightarrow K^+\pi^-$ decays to obtain background-subtracted efficiency tables for kaons and pions as functions of their p and p_T [2]. The kinematic properties of the tracks in signal decays are obtained from simulation, allowing the PID efficiency for each event to be obtained from the tables. Note that this calibration sample is dominated by promptly produced D^* mesons. The remaining contributions to the total efficiency are determined from simulation, and validated using data.

Systematic uncertainties on $\mathcal{B}(B_s^0 \rightarrow D^{*\mp}\pi^\pm)$ are assigned due to the following sources, given in units of 1×10^{-6} , summarised in Table 1. Event selection efficiencies for both modes are found to be consistent in simulation to within 2%, yielding a systematic uncertainty of 0.02. The fit model is varied by replacing the double Gaussian signal shapes with double Crystal Ball [23] functions (with both upper and lower tails), changing the linear combinatorial background shape to quadratic and including a possible contribution




Figure 2: Simultaneous fit to the full data sample in five bins of θ_{bach} : (a) 0–0.046, (b) 0.046–0.067, (c) 0.067–0.092, (d) 0.092–0.128 and (e) 0.128–0.4 rad. Note the y -axis scale is logarithmic and is the same for each bin. Data points are shown in black, the full PDF as a solid blue line and the component PDFs as: (red dot-dashed) partially reconstructed background, (magenta dashed) combinatorial background, (blue dashed) B^0 signal, (black dot-dashed) B_s^0 signal and (green 3 dot-dashed) $B^0 \rightarrow D^*K$ background.

Table 1: Systematic uncertainties on $\mathcal{B}(B_s^0 \rightarrow D^{*\mp}\pi^\pm)$.

Source	Uncertainty (10^{-6})
Efficiency	0.02
Fit model	1.44
Fit bias	0.12
Multiple candidates	0.22
f_s/f_d	0.12
$\mathcal{B}(B^0 \rightarrow D^{*-}\pi^+)$	0.08
Total	1.47

from $\overline{B}_s^0 \rightarrow D^{*-}K^+$. The non-parametric function for the $B^0 \rightarrow D^{*-}K^+$ background was scaled in each bin to account for the change in the width of the B^0 signal. Combined in quadrature these sources contribute 1.44 to the systematic uncertainty. Possible biases in the determination of the fit parameters are investigated by simulated pseudo-experiments, leading to an uncertainty of 0.12. Events with multiple candidates are investigated by performing a fit having chosen one candidate at random. This fit is performed 100 times, with different seeds, and the spread of the results, 0.22, is taken as the systematic uncertainty. The uncertainty on the quantity f_s/f_d contributes 0.12, while that on $\mathcal{B}(B^0 \rightarrow D^{*-}\pi^+)$ gives 0.08. Combining all sources in quadrature, the total absolute systematic uncertainty is 1.47×10^{-6} , and the B_s^0 branching fraction is determined to be $\mathcal{B}(B_s^0 \rightarrow D^{*\mp}\pi^\pm) = (1.5 \pm 3.8 \pm 1.5) \times 10^{-6}$, where the first uncertainty is statistical and the second is systematic.

A number of cross-checks are performed to test the stability of the result. Candidates are divided based upon the hardware trigger decision into three groups; events in which a particle from the signal decay created a large enough cluster in the calorimeter to fire the trigger, events that were triggered independently of the signal decay and those events that were triggered by both the signal decay and the rest of the event. The neural network and PID requirements are tightened and loosened. The non-parametric PDF used to describe the background from D^*K decays was smoothed to eliminate potential statistical fluctuations. All cross-checks give consistent results.

Since no significant signal is observed, upper limits are set, at both 90% and 95% confidence level (CL), using a Bayesian approach. The statistical likelihood curve from the fit is convolved with a Gaussian function of width given by the systematic uncertainty, and the upper limits are taken as the values containing 90% (95%) of the integral of the likelihood in the physical region. The obtained limits are

$$\mathcal{B}(B_s^0 \rightarrow D^{*\mp}\pi^\pm) < 6.1 \text{ (7.8)} \times 10^{-6} \text{ at 90\% (95\%) CL.}$$

In summary, the decay $B_s^0 \rightarrow D^{*\mp}\pi^\pm$ is searched for in a data sample of 1.0 fb^{-1} of data collected with the LHCb detector during 2011. No significant signal is observed and upper limits on the branching fraction are set. The absence of a detectable signal indicates that rescattering effects may make significant contributions to other hadronic decays, such as $B_s^0 \rightarrow \pi^+\pi^-$ and $B_s^0 \rightarrow D\overline{D}$, as recently suggested [5].

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 and Region Auvergne (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (The Netherlands); SCSR (Poland); ANCS/IFA (Romania); MinES, Rosatom, RFBR and NRC “Kurchatov Institute” (Russia); MinECo, XuntaGal and GENCAT (Spain); SNSF and SER (Switzerland); NAS Ukraine (Ukraine); STFC (United Kingdom); NSF (USA). We also acknowledge the support received from the ERC under FP7. The Tier1 computing centres are supported by IN2P3 (France), KIT and BMBF (Germany), INFN (Italy), NWO and SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom). We are thankful for the computing resources put at our disposal by Yandex LLC (Russia), as well as to the communities behind the multiple open source software packages that we depend on.

References

- [1] LHCb collaboration, R. Aaij *et al.*, *First observation of $B^0 \rightarrow D^+D^-$, $D_s^+D^-$ and $D^0\bar{D}^0$ decays*, arXiv:1302.5854, submitted to Phys. Rev. D.
- [2] LHCb collaboration, R. Aaij *et al.*, *Measurement of b -hadron branching fractions for two-body decays into charmless charged hadrons*, JHEP **10** (2012) 037, arXiv:1206.2794.
- [3] M. Gronau, O. F. Hernandez, D. London, and J. L. Rosner, *Broken $SU(3)$ symmetry in two-body B decays*, Phys. Rev. **D52** (1995) 6356, arXiv:hep-ph/9504326.
- [4] R. Fleischer, *New strategies to obtain insights into CP violation through $B_s \rightarrow D_s^\pm K^\mp$, $D_s^{*\pm} K^\mp$, ... and $B_d \rightarrow D^\pm \pi^\mp$, $D^{*\pm} \pi^\mp$, ... decays*, Nucl. Phys. **B671** (2003) 459, arXiv:hep-ph/0304027.
- [5] M. Gronau, D. London, and J. L. Rosner, *Rescattering contributions to rare B -meson decays*, Phys. Rev. **D87** (2013) 036008, arXiv:1211.5785.
- [6] LHCb collaboration, R. Aaij *et al.*, *First evidence for the annihilation decay mode $B^+ \rightarrow D_s^+ \phi$* , JHEP **02** (2013) 043, arXiv:1210.1089.
- [7] BaBar collaboration, B. Aubert *et al.*, *Branching fraction measurement of $B^0 \rightarrow D^{(*)+} \pi^-$, $B^- \rightarrow D^{(*)0} \pi^-$ and isospin analysis of $\bar{B} \rightarrow D^{(*)} \pi$ decays*, Phys. Rev. **D75** (2007) 031101, arXiv:hep-ex/0610027.
- [8] Particle Data Group, J. Beringer *et al.*, *Review of particle physics*, Phys. Rev. **D86** (2012) 010001.

- [9] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [10] R. Aaij *et al.*, *The LHCb trigger and its performance*, arXiv:1211.3055, submitted to JINST.
- [11] M. Adinolfi *et al.*, *Performance of the LHCb RICH detector at the LHC*, arXiv:1211.6759, submitted to Eur. Phys. J. C.
- [12] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay $B^0 \rightarrow \bar{D}^0 K^+ K^-$ and evidence of $B_s^0 \rightarrow \bar{D}^0 K^+ K^-$* , Phys. Rev. Lett. **109** (2012) 131801, arXiv:1207.5991.
- [13] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, arXiv:hep-ph/0603175.
- [14] I. Belyaev *et al.*, *Handling of the generation of primary events in GAUSS, the LHCb simulation framework*, Nuclear Science Symposium Conference Record (NSS/MIC) **IEEE** (2010) 1155.
- [15] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [16] GEANT4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; GEANT4 collaboration, S. Agostinelli *et al.*, *GEANT4: a simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [17] M. Clemencic *et al.*, *The LHCb simulation application, GAUSS: design, evolution and experience*, J. of Phys. : Conf. Ser. **331** (2011) 032023.
- [18] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay $\bar{B}_s^0 \rightarrow D^0 K^{*0}$ and a measurement of the ratio of branching fractions $\frac{\mathcal{B}(\bar{B}_s^0 \rightarrow D^0 K^{*0})}{\mathcal{B}(\bar{B}^0 \rightarrow D^0 \rho^0)}$* , Phys. Lett. **B706** (2011) 32, arXiv:1110.3676.
- [19] M. Pivk and F. R. Le Diberder, *sPlot: a statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, arXiv:physics/0402083.
- [20] M. Feindt and U. Kerzel, *The NeuroBayes neural network package*, Nucl. Instrum. Meth. **A559** (2006) 190.
- [21] LHCb collaboration, R. Aaij *et al.*, *Observation of CP violation in $B^\pm \rightarrow DK^\pm$ decays*, Phys. Lett. **B712** (2012) 203, arXiv:1203.3662.
- [22] LHCb collaboration, R. Aaij *et al.*, *Measurement of the ratio of fragmentation functions f_s/f_d and the dependence on B meson kinematics*, arXiv:1301.5286, submitted to JHEP.
- [23] T. Skwarnicki, *A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances*, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.