

Measurement of the Mass of the D^0 Meson

J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ E. Grauges,² A. Palano^{ab,3} G. Eigen,⁴ B. Stugu,⁴ D. N. Brown,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ M. J. Lee,⁵ G. Lynch,⁵ H. Koch,⁶ T. Schroeder,⁶ C. Hearty,⁷ T. S. Mattison,⁷ J. A. McKenna,⁷ R. Y. So,⁷ A. Khan,⁸ V. E. Blinov^{ac,9} A. R. Buzykaev^{a,9} V. P. Druzhinin^{ab,9} V. B. Golubev^{ab,9} E. A. Kravchenko^{ab,9} A. P. Onuchin^{ac,9} S. I. Serednyakov^{ab,9} Yu. I. Skovpen^{ab,9} E. P. Solodov^{ab,9} K. Yu. Todyshev^{ab,9} A. N. Yushkov^{a,9} D. Kirkby,¹⁰ A. J. Lankford,¹⁰ M. Mandelkern,¹⁰ B. Dey,¹¹ J. W. Gary,¹¹ O. Long,¹¹ G. M. Vitug,¹¹ C. Campagnari,¹² M. Franco Sevilla,¹² T. M. Hong,¹² D. Kovalskiy,¹² J. D. Richman,¹² C. A. West,¹² A. M. Eisner,¹³ W. S. Lockman,¹³ B. A. Schumm,¹³ A. Seiden,¹³ D. S. Chao,¹⁴ C. H. Cheng,¹⁴ B. Echenard,¹⁴ K. T. Flood,¹⁴ D. G. Hitlin,¹⁴ P. Ongmongkolkul,¹⁴ F. C. Porter,¹⁴ R. Andreassen,¹⁵ Z. Huard,¹⁵ B. T. Meadows,¹⁵ C. Pappenheimer,¹⁵ B. G. Pushpawela,¹⁵ M. D. Sokoloff,¹⁵ L. Sun,¹⁵ P. C. Bloom,¹⁶ W. T. Ford,¹⁶ A. Gaz,¹⁶ U. Nauenberg,¹⁶ J. G. Smith,¹⁶ S. R. Wagner,¹⁶ R. Ayad,^{17,*} W. H. Toki,¹⁷ B. Spaan,¹⁸ R. Schwierz,¹⁹ D. Bernard,²⁰ M. Verderi,²⁰ S. Playfer,²¹ D. Bettoni^{a,22} C. Bozzi^{a,22} R. Calabrese^{ab,22} G. Cibinetto^{ab,22} E. Fioravanti^{ab,22} I. Garzia^{ab,22} E. Luppi^{ab,22} L. Piemontese^{a,22} V. Santoro^{a,22} R. Baldini-Ferrolì,²³ A. Calcaterra,²³ R. de Sangro,²³ G. Finocchiaro,²³ S. Martellotti,²³ P. Patteri,²³ I. M. Peruzzi,^{23,†} M. Piccolo,²³ M. Rama,²³ A. Zallo,²³ R. Contri^{ab,24} E. Guido^{ab,24} M. Lo Vetere^{ab,24} M. R. Monge^{ab,24} S. Passaggio^{a,24} C. Patrignani^{ab,24} E. Robutti^{a,24} B. Bhuyan,²⁵ V. Prasad,²⁵ M. Morii,²⁶ A. Adametz,²⁷ U. Uwer,²⁷ H. M. Lacker,²⁸ P. D. Dauncey,²⁹ U. Mallik,³⁰ C. Chen,³¹ J. Cochran,³¹ W. T. Meyer,³¹ S. Prell,³¹ A. V. Gritsan,³² N. Arnaud,³³ M. Davier,³³ D. Derkach,³³ G. Grosdidier,³³ F. Le Diberder,³³ A. M. Lutz,³³ B. Malaescu,^{33,‡} P. Roudeau,³³ A. Stocchi,³³ G. Wormser,³³ D. J. Lange,³⁴ D. M. Wright,³⁴ J. P. Coleman,³⁵ J. R. Fry,³⁵ E. Gabathuler,³⁵ D. E. Hutchcroft,³⁵ D. J. Payne,³⁵ C. Touramanis,³⁵ A. J. Bevan,³⁶ F. Di Lodovico,³⁶ R. Sacco,³⁶ G. Cowan,³⁷ J. Bougher,³⁸ D. N. Brown,³⁸ C. L. Davis,³⁸ A. G. Denig,³⁹ M. Fritsch,³⁹ W. Gradl,³⁹ K. Griessinger,³⁹ A. Hafner,³⁹ E. Prencipe,³⁹ K. R. Schubert,³⁹ R. J. Barlow,^{40,§} G. D. Lafferty,⁴⁰ E. Behn,⁴¹ R. Cenci,⁴¹ B. Hamilton,⁴¹ A. Jawahery,⁴¹ D. A. Roberts,⁴¹ R. Cowan,⁴² D. Dujmic,⁴² G. Sciolla,⁴² R. Cheaib,⁴³ P. M. Patel,^{43,¶} S. H. Robertson,⁴³ P. Biassoni^{ab,44} N. Neri^{a,44} F. Palombo^{ab,44} L. Cremaldi,⁴⁵ R. Godang,^{45,**} P. Sonnek,⁴⁵ D. J. Summers,⁴⁵ M. Simard,⁴⁶ P. Taras,⁴⁶ G. De Nardo^{ab,47} D. Monorchio^{ab,47} G. Onorato^{ab,47} C. Sciacca^{ab,47} M. Martinelli,⁴⁸ G. Raven,⁴⁸ C. P. Jessop,⁴⁹ J. M. LoSecco,⁴⁹ K. Honscheid,⁵⁰ R. Kass,⁵⁰ J. Brau,⁵¹ R. Frey,⁵¹ N. B. Sinev,⁵¹ D. Strom,⁵¹ E. Torrence,⁵¹ E. Feltresi^{ab,52} M. Margoni^{ab,52} M. Morandin^{a,52} M. Posocco^{a,52} M. Rotondo^{a,52} G. Simi^{a,52} F. Simonetto^{ab,52} R. Stroili^{ab,52} S. Akar,⁵³ E. Ben-Haim,⁵³ M. Bomben,⁵³ G. R. Bonneaud,⁵³ H. Briand,⁵³ G. Calderini,⁵³ J. Chauveau,⁵³ Ph. Leruste,⁵³ G. Marchiori,⁵³ J. Ocariz,⁵³ S. Sitt,⁵³ M. Biasini^{ab,54} E. Manoni^{a,54} S. Pacetti^{ab,54} A. Rossi^{a,54} C. Angelini^{ab,55} G. Batignani^{ab,55} S. Bettarini^{ab,55} M. Carpinelli^{ab,55,††} G. Casarosa^{ab,55} A. Cervelli^{ab,55} F. Forti^{ab,55} M. A. Giorgi^{ab,55} A. Lusiani^{ac,55} B. Oberholzer^{ab,55} E. Paoloni^{ab,55} A. Perez^{a,55} G. Rizzo^{ab,55} J. J. Walsh^{a,55} D. Lopes Pegna,⁵⁶ J. Olsen,⁵⁶ A. J. S. Smith,⁵⁶ R. Faccini^{ab,57} F. Ferrarotto^{a,57} F. Ferroni^{ab,57} M. Gaspero^{ab,57} L. Li Gioi^{a,57} G. Piredda^{a,57} C. Büniger,⁵⁸ O. Grünberg,⁵⁸ T. Hartmann,⁵⁸ T. Leddig,⁵⁸ C. Voß,⁵⁸ R. Waldi,⁵⁸ T. Adye,⁵⁹ E. O. Olaiya,⁵⁹ F. F. Wilson,⁵⁹ S. Emery,⁶⁰ G. Hamel de Monchenault,⁶⁰ G. Vasseur,⁶⁰ Ch. Yèche,⁶⁰ F. Anulli,^{61,‡‡} D. Aston,⁶¹ D. J. Bard,⁶¹ J. F. Benitez,⁶¹ C. Cartaro,⁶¹ M. R. Convery,⁶¹ J. Dorfan,⁶¹ G. P. Dubois-Felsmann,⁶¹ W. Dunwoodie,⁶¹ M. Ebert,⁶¹ R. C. Field,⁶¹ B. G. Fulsom,⁶¹ A. M. Gabareen,⁶¹ M. T. Graham,⁶¹ C. Hast,⁶¹ W. R. Innes,⁶¹ P. Kim,⁶¹ M. L. Kocian,⁶¹ D. W. G. S. Leith,⁶¹ P. Lewis,⁶¹ D. Lindemann,⁶¹ B. Lindquist,⁶¹ S. Luitz,⁶¹ V. Luth,⁶¹ H. L. Lynch,⁶¹ D. B. MacFarlane,⁶¹ D. R. Muller,⁶¹ H. Neal,⁶¹ S. Nelson,⁶¹ M. Perl,⁶¹ T. Pulliam,⁶¹ B. N. Ratcliff,⁶¹ A. Roodman,⁶¹ A. A. Salnikov,⁶¹ R. H. Schindler,⁶¹ A. Snyder,⁶¹ D. Su,⁶¹ M. K. Sullivan,⁶¹ J. Va'vra,⁶¹ A. P. Wagner,⁶¹ W. F. Wang,⁶¹ W. J. Wisniewski,⁶¹ M. Wittgen,⁶¹ D. H. Wright,⁶¹ H. W. Wulsin,⁶¹ V. Ziegler,⁶¹ W. Park,⁶² M. V. Purohit,⁶² R. M. White,^{62,§§} J. R. Wilson,⁶² A. Randle-Conde,⁶³ S. J. Sekula,⁶³ M. Bellis,⁶⁴ P. R. Burchat,⁶⁴ T. S. Miyashita,⁶⁴ E. M. T. Puccio,⁶⁴ M. S. Alam,⁶⁵ J. A. Ernst,⁶⁵ R. Gorodeisky,⁶⁶ N. Guttman,⁶⁶ D. R. Peimer,⁶⁶ A. Soffer,⁶⁶ S. M. Spanier,⁶⁷ J. L. Ritchie,⁶⁸ A. M. Ruland,⁶⁸ R. F. Schwitters,⁶⁸ B. C. Wray,⁶⁸ J. M. Izen,⁶⁹ X. C. Lou,⁶⁹ F. Bianchi^{ab,70} F. De Mori^{ab,70} A. Filippi^{a,70} D. Gamba^{ab,70} S. Zambito^{ab,70} L. Lanceri^{ab,71} L. Vitale^{ab,71} F. Martinez-Vidal,⁷² A. Oyanguren,⁷² P. Villanueva-Perez,⁷²

H. Ahmed,⁷³ J. Albert,⁷³ Sw. Banerjee,⁷³ F. U. Bernlochner,⁷³ H. H. F. Choi,⁷³ G. J. King,⁷³ R. Kowalewski,⁷³
M. J. Lewczuk,⁷³ T. Lueck,⁷³ I. M. Nugent,⁷³ J. M. Roney,⁷³ R. J. Sobie,⁷³ N. Tasneem,⁷³ T. J. Gershon,⁷⁴
P. F. Harrison,⁷⁴ T. E. Latham,⁷⁴ H. R. Band,⁷⁵ S. Dasu,⁷⁵ Y. Pan,⁷⁵ R. Prepost,⁷⁵ and S. L. Wu⁷⁵

(The BABAR Collaboration)

¹Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP),
Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France

²Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

³INFN Sezione di Bari^a; Dipartimento di Fisica, Università di Bari^b, I-70126 Bari, Italy

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁷University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

⁸Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

⁹Budker Institute of Nuclear Physics SB RAS, Novosibirsk 630090^a,
Novosibirsk State University, Novosibirsk 630090^b,

Novosibirsk State Technical University, Novosibirsk 630092^c, Russia

¹⁰University of California at Irvine, Irvine, California 92697, USA

¹¹University of California at Riverside, Riverside, California 92521, USA

¹²University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹³University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁴California Institute of Technology, Pasadena, California 91125, USA

¹⁵University of Cincinnati, Cincinnati, Ohio 45221, USA

¹⁶University of Colorado, Boulder, Colorado 80309, USA

¹⁷Colorado State University, Fort Collins, Colorado 80523, USA

¹⁸Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany

¹⁹Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²⁰Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS/IN2P3, F-91128 Palaiseau, France

²¹University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²²INFN Sezione di Ferrara^a; Dipartimento di Fisica e Scienze della Terra, Università di Ferrara^b, I-44122 Ferrara, Italy

²³INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy

²⁴INFN Sezione di Genova^a; Dipartimento di Fisica, Università di Genova^b, I-16146 Genova, Italy

²⁵Indian Institute of Technology Guwahati, Guwahati, Assam, 781 039, India

²⁶Harvard University, Cambridge, Massachusetts 02138, USA

²⁷Universität Heidelberg, Physikalisches Institut, D-69120 Heidelberg, Germany

²⁸Humboldt-Universität zu Berlin, Institut für Physik, D-12489 Berlin, Germany

²⁹Imperial College London, London, SW7 2AZ, United Kingdom

³⁰University of Iowa, Iowa City, Iowa 52242, USA

³¹Iowa State University, Ames, Iowa 50011-3160, USA

³²Johns Hopkins University, Baltimore, Maryland 21218, USA

³³Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11,
Centre Scientifique d'Orsay, F-91898 Orsay Cedex, France

³⁴Lawrence Livermore National Laboratory, Livermore, California 94550, USA

³⁵University of Liverpool, Liverpool L69 7ZE, United Kingdom

³⁶Queen Mary, University of London, London, E1 4NS, United Kingdom

³⁷University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom

³⁸University of Louisville, Louisville, Kentucky 40292, USA

³⁹Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany

⁴⁰University of Manchester, Manchester M13 9PL, United Kingdom

⁴¹University of Maryland, College Park, Maryland 20742, USA

⁴²Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA

⁴³McGill University, Montréal, Québec, Canada H3A 2T8

⁴⁴INFN Sezione di Milano^a; Dipartimento di Fisica, Università di Milano^b, I-20133 Milano, Italy

⁴⁵University of Mississippi, University, Mississippi 38677, USA

⁴⁶Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7

⁴⁷INFN Sezione di Napoli^a; Dipartimento di Scienze Fisiche,
Università di Napoli Federico II^b, I-80126 Napoli, Italy

⁴⁸NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands

⁴⁹University of Notre Dame, Notre Dame, Indiana 46556, USA

⁵⁰Ohio State University, Columbus, Ohio 43210, USA

⁵¹University of Oregon, Eugene, Oregon 97403, USA

⁵²INFN Sezione di Padova^a; Dipartimento di Fisica, Università di Padova^b, I-35131 Padova, Italy

- ⁵³Laboratoire de Physique Nucléaire et de Hautes Energies,
IN2P3/CNRS, Université Pierre et Marie Curie-Paris6,
Université Denis Diderot-Paris7, F-75252 Paris, France
- ⁵⁴INFN Sezione di Perugia^a; Dipartimento di Fisica, Università di Perugia^b, I-06123 Perugia, Italy
- ⁵⁵INFN Sezione di Pisa^a; Dipartimento di Fisica,
Università di Pisa^b; Scuola Normale Superiore di Pisa^c, I-56127 Pisa, Italy
- ⁵⁶Princeton University, Princeton, New Jersey 08544, USA
- ⁵⁷INFN Sezione di Roma^a; Dipartimento di Fisica,
Università di Roma La Sapienza^b, I-00185 Roma, Italy
- ⁵⁸Universität Rostock, D-18051 Rostock, Germany
- ⁵⁹Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶⁰CEA, Irfu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France
- ⁶¹SLAC National Accelerator Laboratory, Stanford, California 94309 USA
- ⁶²University of South Carolina, Columbia, South Carolina 29208, USA
- ⁶³Southern Methodist University, Dallas, Texas 75275, USA
- ⁶⁴Stanford University, Stanford, California 94305-4060, USA
- ⁶⁵State University of New York, Albany, New York 12222, USA
- ⁶⁶Tel Aviv University, School of Physics and Astronomy, Tel Aviv, 69978, Israel
- ⁶⁷University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁶⁸University of Texas at Austin, Austin, Texas 78712, USA
- ⁶⁹University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁷⁰INFN Sezione di Torino^a; Dipartimento di Fisica, Università di Torino^b, I-10125 Torino, Italy
- ⁷¹INFN Sezione di Trieste^a; Dipartimento di Fisica, Università di Trieste^b, I-34127 Trieste, Italy
- ⁷²IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
- ⁷³University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- ⁷⁴Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
- ⁷⁵University of Wisconsin, Madison, Wisconsin 53706, USA

We report a measurement of the D^0 meson mass using the decay chain $D^*(2010)^+ \rightarrow D^0\pi^+$ with $D^0 \rightarrow K^-K^-K^+\pi^+$. The data were recorded with the BABAR detector at center-of-mass energies at and near the $\Upsilon(4S)$ resonance, and correspond to an integrated luminosity of approximately 477 fb^{-1} . We obtain $m(D^0) = (1864.841 \pm 0.048 \pm 0.063) \text{ MeV}$, where the quoted errors are statistical and systematic, respectively. The uncertainty of this measurement is half that of the best previous measurement.

PACS numbers: 13.20.Fc, 13.25.Ft, 14.40.Lb, 12.38.Gc, 12.38.Qk, 12.39.Ki, 12.39.Pn

INTRODUCTION

The D^0 is one of the ground states of the charm mesons, and its mass sets the mass scale for the heavier excited states. As such, the D^0 mass is directly relevant to several measurements. For example, the reported $D^*(2010)^+$ mass is the sum of the nominal D^0 mass and the measured difference, Δm , between the masses of the D^{*+} and D^0 mesons. Mixing parameters in the $D^0 - \bar{D}^0$ system can be extracted from a time-dependent amplitude analysis of the mass-constrained Dalitz plot, as in the analyses of the $K_S^0\pi^+\pi^-$ and $K_S^0K^+K^-$ final states [1–3]. A new value of the D^0 mass would shift an event's position in the Dalitz plot if the final state is constrained to the D^0 mass value. A precise D^0 mass measurement can also serve as a reference point in magnetic field calibration studies. The D^0 mass is also important for the determination of the relationship of the $D^0\bar{D}^{*0}$ threshold to the mass of $X(3872)$. In this regard, a recent analysis by LHCb [4] reported $J^{PC} = 1^{++}$ for $X(3872)$, which favors an exotic model such as one in which the $X(3872)$ is a $D^0\bar{D}^{*0}$ molecule [5].

We measure the D^0 mass using the decay chain

$D^*(2010)^+ \rightarrow D^0\pi^+$, $D^0 \rightarrow K^-K^-K^+\pi^+$. The use of charge conjugate reactions is implied here and throughout this paper. We chose this mode specifically for the relatively low Q-value, $m(K^-K^-K^+\pi^+) - 3m(K^\pm) - m(\pi^+) \approx 250 \text{ MeV}$, which produces small backgrounds, yields good resolution, and minimizes systematic uncertainties. The previous most precise measurements were made by the CLEO collaboration [6], which reported $m(D^0) = (1864.847 \pm 0.150 \pm 0.095) \text{ MeV}$, and by the LHCb collaboration [7], which reported $m(D^0) = (1864.75 \pm 0.15 \pm 0.11) \text{ MeV}$, where the errors are statistical and systematic, respectively. The integrated luminosity for this analysis is about 60 times that of the CLEO analysis. After selection criteria are chosen to minimize systematic uncertainties, our sample is about 15 times larger than the CLEO sample [6]. The LHCb analysis [7] uses the same decay channel we use, but their signal has about half the number of events, and our signal-to-background ratio is much higher.

DATA SAMPLE AND DETECTOR

This analysis is based on a data sample corresponding to an integrated luminosity of approximately 477 fb^{-1} [8] recorded at, and 40 MeV below, the $\Upsilon(4S)$ resonance by the *BABAR* detector at the PEP-II asymmetric energy e^+e^- collider at the SLAC National Accelerator Laboratory. The *BABAR* detector is described in detail elsewhere [9, 10]. Here, we summarize the most relevant components. The momenta of charged particles are measured with a combination of a cylindrical drift chamber (DCH) and a 5-layer silicon vertex tracker (SVT), both operating within the 1.5 T magnetic field of a superconducting solenoid. Information from a ring-imaging Cherenkov detector is combined with specific ionization (dE/dx) measurements from the SVT and DCH to identify charged-kaon and pion candidates. Electrons are identified, and photons measured, with a CsI(Tl) electromagnetic calorimeter. The return yoke of the superconducting coil is instrumented with tracking chambers for the identification of muons.

EVENT SELECTION

We expect our measurement to be dominated by systematic uncertainties, so we focus on choosing a balanced set of selection criteria that produces a very clean signal, to control systematic uncertainties, while still preserving a relatively large number of signal events. To avoid potential bias, the final selection criteria are chosen based on extensive studies of Monte Carlo (MC) simulation events and on 5% of the real data; the latter are not used in the mass measurement. With the final selection criteria determined from these preliminary studies, systematic uncertainties are determined from a blind analysis of the remaining 95% of the data, in which the measured mass of each event is shifted by an unknown, common offset. After completion of all the systematic uncertainty studies, we removed the offset and performed a fit to the measured mass spectrum.

We reconstruct the decay chain $D^*(2010)^+ \rightarrow D^0\pi^+$, $D^0 \rightarrow K^-K^+K^+\pi^+$ using a kinematic fit requiring that the D^0 daughters emerge from a common vertex and that the D^0 and the $D^*(2010)^+$ (D^{*+}) daughter pion momenta point back to the primary vertex, with the additional constraint that the D^{*+} candidates originate from the luminous region [11]. The χ^2/ν (ν denotes the number of degrees of freedom) reported by the fit was required to be less than 20/9, corresponding to a fit probability greater than 1.8%. For events with multiple candidates, we choose the candidate with the largest confidence level.

To produce a good balance between expected statistical precision and systematic uncertainty we apply a variety of selection criteria. We consider the momentum of the D^{*+} in the e^+e^- center-of-mass frame, $p^*(D^{*+})$,

and Δm . In addition, we require kaon and pion tracks to pass particle identification (PID) selections. We look at the statistical significance and purity of data sets with varying $p^*(D^{*+})$, Δm , and PID cuts. We choose a reasonable set of selection criteria, $p^*(D^{*+}) > 2.5 \text{ GeV}$, $|\Delta m - \Delta m_{\text{PDG}}| < 1.5 \text{ MeV}$, where the PDG subscript indicates the value listed by the PDG [12], before the remaining 95% of the data sample is studied, thus avoiding potential analyst bias.

In approximately 1% of events, we mis-reconstruct the decay and swap the pion from the D^{*+} decay (referred to as the slow pion, π_s) with the pion from the D^0 decay. This exchange could potentially shift the measured mass value, as these events tend to concentrate at lower values of reconstructed D^0 mass. To eliminate these events, we define the variables $m'(D^0) = m(K^-K^+K^+\pi_s^+)$ and $\Delta m' = m(K^-K^+K^+\pi_s^+) - m(K^-K^+K^+\pi_s^+)$. In this $(m', \Delta m')$ system, correctly reconstructed events are shifted away from the $(m', \Delta m')$ signal region, and events with the exchanged pion mis-reconstruction are shifted into the $(m', \Delta m')$ signal region. The requirement $\Delta m' > 0.15 \text{ GeV}$ eliminates 80% of the exchanged events that survive as D^0 candidates with essentially no loss of correctly reconstructed signal. According to MC simulation, the surviving 20% of mis-reconstructed events have daughter-pion and slow-pion momenta so similar that the wrong mass values have a negligible effect on our D^0 mass determination.

In a prior study of the line shape and relativistic Breit-Wigner pole position in $D^{*+} \rightarrow D^0\pi_s^+$ decays [13], we observed strong mass and mass-difference dependence on track momentum. We also observed that the reconstructed K_S^0 mass value was systematically low and varied with momentum. This effect was not replicated in MC.

We determined that increasing the magnitude of the laboratory momentum by a factor of 1.0003 and increasing the energy loss reported by the Kalman fit by 1.8% and 5.9% for the beam pipe and SVT, respectively, removes the momentum dependence and offset of the reconstructed K_S^0 mass. The increase in the magnitude of momentum corresponds to increasing the magnetic field by 0.45 mT. The corrections also eliminated the momentum dependence of the mass difference, which was the subject of the $D^{*+} \rightarrow D^0\pi_s^+$ study. The procedure for determining these parameters and those used for systematic uncertainty corrections is detailed in Ref. [13]. We apply the same magnetic-field and material-model-corrections in this analysis.

In studying the K_S^0 signal, we observed that even after the corrections described above, the K_S^0 mass dropped dramatically when either of its daughter tracks had $\cos\theta > 0.89$. We therefore reject events for which any of the daughter tracks of the D^0 has $\cos\theta > 0.89$. This criterion reduces the final data sample by approximately 10%. We additionally require that the momentum of the

slow pion is at least 150 MeV in the laboratory frame in order that the MC laboratory momentum distribution accurately replicates that observed for data.

FIT TO DATA

To determine the D^0 mass, we perform an extended, unbinned maximum likelihood fit over the candidate mass range 1.75 – 1.98 GeV, using a Voigtian distribution to describe the signal. This function provides better agreement between the model and the $m(K^-K^-K^+\pi^+)$ signal distribution than a multi-Gaussian model. The background is described by an exponential function. The Voigtian distribution is the convolution of a Cauchy and a Gaussian distribution, defined as

$$V(m) = \int_{-\infty}^{\infty} d\mu \exp\left[\frac{-(\mu - m_D)^2}{2\sigma^2}\right] \frac{1}{(m - \mu)^2 + \frac{\gamma^2}{4}}, \quad (1)$$

where m is the $K^-K^-K^+\pi^+$ invariant mass, and the three parameters are m_D , the D^0 mass, and γ and σ , two ad-hoc resolution parameters. The fit to the data is shown in Fig. 1, and the results of this fit are summarized in Table I. At the peak, the signal-to-background ratio is approximately 175:1.

TABLE I: The results of the fits to data for the $K^-K^-K^+\pi^+$ channel (statistical uncertainties only).

| Fit Parameters | Values |
|-------------------------|----------------------|
| Number of signal events | 4345 ± 70 |
| $m(D^0)$ (MeV) | 1864.841 ± 0.048 |
| γ (MeV) | 2.596 ± 0.152 |
| σ (MeV) | 1.762 ± 0.086 |

SYSTEMATIC UNCERTAINTIES

Systematic uncertainties arise from a variety of sources. The dominant systematic uncertainty is the effect of the uncertainty in the charged-kaon mass on the $K^-K^-K^+\pi^+$ invariant mass. The corrections to the momentum scale and dE/dx energy loss in detector material are also varied to account for the uncertainty in the K_S^0 mass [12, 13]. Additionally, we vary the fit model to estimate the systematic uncertainty associated with our choice of signal and background shape, and we study the systematic uncertainty associated with inner bremsstrahlung. We then divide the data into disjoint subsets corresponding to intervals of azimuthal angle, ϕ , the laboratory momentum of the D^0 , p_{lab} , and Δm . We assign systematic uncertainties using a method similar




FIG. 1: (color online) Results of the extended, unbinned maximum likelihood fit to the reconstructed D^0 mass distribution. Normalized residuals are defined as $(N_o - N_p)/\sqrt{N_p}$, where N_o and N_p are observed and predicted numbers of events, respectively.

TABLE II: Systematic uncertainties for the D^0 mass measurement.

| Source | σ_{sys} (MeV) |
|--|-----------------------------|
| K^\pm mass uncertainty | 0.046 |
| Magnetic field and material model | 0.031 |
| Signal shape | 0.009 |
| Background model | 0.005 |
| Bremsstrahlung | < 0.001 |
| Disjoint ϕ interval variation | 0.000 |
| Disjoint p_{lab} interval variation | 0.000 |
| Disjoint Δm interval variation | 0.028 |
| Sum in quadrature | 0.063 |

to that used in obtaining the PDG scale factor [12], as described below. The systematic uncertainties are summarized in Table II.

To estimate the effect of the charged-kaon mass uncertainty on the reconstructed D^0 mass, we vary the charged-kaon mass value by $\pm 1\sigma_{\text{PDG}} = \pm 16$ keV [12], re-determine $m(K^-K^-K^+\pi^+)$ of each event, and refit each new $m(K^-K^-K^+\pi^+)$ distribution. We take the average of the magnitude of the differences from the nominal fit

results as the systematic uncertainty. We find an average variation of 46 keV, which proves to be the largest source of systematic uncertainty. The charged pion mass is known to a much higher precision, so its mass uncertainty has a negligible effect on the reconstructed D^0 mass value.

We estimate the uncertainty associated with the choice of our correction parameters for the magnetic field and detector material model by examining the variation between the fit results using the nominal parameter values and those obtained by tuning to the mass values $m_{\text{PDG}}(K_S^0) \pm 1\sigma_{\text{PDG}}$ [12, 13]. We compare the fitted mass value from the nominal fit to the mass values extracted using the $\pm 1\sigma_{\text{PDG}}$ correction parameters and take the largest difference between those fits and the nominal fit, 31 keV, as a measure of the systematic uncertainty.

We alternatively fit the real data with a double Gaussian signal shape and see a shift of 9 keV in the central mass value. We also change the background model used in the fit procedure from the nominal exponential distribution to a second degree polynomial. Although it provides a relatively poor description, using a quadratic background changes the central value by only 5 keV. We report the magnitudes of these shifts as systematic uncertainties in Table II.

We study the systematic uncertainty associated with inner bremsstrahlung using PHOTOS [14]. About 3.5% of the generated $D^0 \rightarrow K^- K^- K^+ \pi^+ n \gamma$ decays had at least one photon, and in these events the $K^- K^- K^+ \pi^+$ invariant mass averaged about 15 keV below the nominal D^0 mass, leading to a shift in the full sample average of less than 1 keV; we therefore neglect this as a source of systematic uncertainty.

We study the D^0 mass dependence on $p_{\text{lab}}(D^0)$, on Δm , and on ϕ , based on dependences observed in previous *BABAR* analyses. We divide the data into 10 subsets for p_{lab} and Δm , and 12 subsets for ϕ (to reflect the 6-fold symmetry of the detector); the intervals in the p_{lab} and Δm plots have variable widths selected to include roughly equal numbers of signal events. If the fit results from the disjoint subsets are compatible with a constant value, in the sense that $\chi^2/\nu < 1$, we assign no systematic uncertainty. However, if we find $\chi^2/\nu > 1$ we ascribe an uncertainty using a variation on the scale factor method used by the PDG (see the discussion of unconstrained averaging [12]). In our version of this procedure, we determine the systematic uncertainty for unknown detector issues to be the value that, when summed in quadrature with the statistical error, produces $\chi^2/\nu = 1$, so that

$$\sigma_{\text{sys}} = \sigma_{\text{stat}} \sqrt{S^2 - 1}, \quad (2)$$

where $S^2 = \chi^2/\nu$. The χ^2 statistic gives a measure of fluctuations, including those expected from statistics,

and those from systematic effects. Once we remove statistical fluctuations, we associate what remains with a possible systematic uncertainty. Fig. 2 does not show any obvious patterns of variations, but using this procedure we assign a 28 keV systematic uncertainty due to the variations seen in the Δm subsets. We find that the variation of the D^0 mass with azimuthal angle and with laboratory momentum is consistent with statistical fluctuations, and thus assign no associated systematic error.




FIG. 2: The value of $m(D^0)$ obtained from fits to data divided into disjoint subsets in (a) ϕ , (b) $p_{\text{lab}}(D^0)$, and (c) Δm . Each point represents an individual fit, and the horizontal line indicates the nominal fit result; the points are plotted at the mean value in the range, and the uncertainty in the mean values is smaller than the marker size.

CONCLUSION

We have measured the D^0 mass with more than twice the precision of the previous best measurement [6] by analyzing a high-purity sample of D^{*+} continuum events produced in e^+e^- collisions near 10.6 GeV. The corresponding integrated luminosity is approximately 477 fb^{-1} . Reconstructing the decay $D^0 \rightarrow K^- K^- K^+ \pi^+$, we measure $m(D^0) = (1864.841 \pm 0.048 \pm 0.063)\text{ MeV}$. The dominant source of systematic uncertainty is the uncertainty in the charged-kaon mass. This measurement will significantly improve the precision of the current world average, $(1864.86 \pm 0.13)\text{ MeV}$ [12].

The largest source of uncertainty in the present measurement comes from the charged-kaon mass uncertainty, reported to be $\pm 16\text{ keV}$ by the PDG [12]. The two most precise measurements of the charged-kaon mass [15, 16] are both more than one standard deviation from the PDG central value, and they differ by 60 keV with reported uncertainties of only 7 keV and 11 keV. Therefore, we also give the dependence of our final result on $m(K^\pm)$, $m(D^0) = (1864.841 \pm 0.048 \pm 0.043 + 3[m(K^\pm) - 493.677])\text{ MeV}$, where the uncertainties are statistical and instrumental, respectively. Using this value, the D^0 mass can readily be obtained from an improved kaon mass value.

We may use our result to estimate the binding energy of the $X(3872)$, interpreting it as a $D^0 \bar{D}^{*0}$ molecule. Combining it with the current PDG average [12] for the $D^*(2007)^0 - D^0$ mass difference, $(142.12 \pm 0.07)\text{ MeV}$, and the PDG average mass of the $X(3872)$, $(3871.68 \pm 0.17)\text{ MeV}$, yields

$$\begin{aligned} E_b &= m(D^0) + m(D^{*0}) - m(X(3872)) \\ &= 2m(D^0) + [m(D^{*0}) - m(D^0)] - m(X(3872)) \\ &= (0.12 \pm 0.24)\text{ MeV}. \end{aligned}$$

ACKNOWLEDGEMENTS

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), the Commissariat à l’Energie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemein-

schaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Education and Science of the Russian Federation, Ministerio de Ciencia e Innovación (Spain), and the Science and Technology Facilities Council (United Kingdom). Individuals have received support from the Marie-Curie IEF program (European Union) and the A. P. Sloan Foundation (USA). The University of Cincinnati is gratefully acknowledged for its support of this research through a WISE (Women in Science and Engineering) fellowship to C. Pappeneheimer.

* Now at the University of Tabuk, Tabuk 71491, Saudi Arabia

† Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

‡ Now at Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Paris, France

§ Now at the University of Huddersfield, Huddersfield HD1 3DH, UK

¶ Deceased

** Now at University of South Alabama, Mobile, Alabama 36688, USA

†† Also with Università di Sassari, Sassari, Italy

‡‡ Also with INFN Sezione di Roma, Roma, Italy

§§ Now at Universidad Técnica Federico Santa María, Valparaíso, Chile 2390123

- [1] P. del Amo Sanchez *et al.* (*BABAR* Collaboration), *Phys. Rev. Lett.* **105**, 081803 (2010).
- [2] D. M. Asner *et al.* (*CLEO* Collaboration), *Phys. Rev. D* **72**, 012001 (2005).
- [3] L. M. Zhang *et al.* (*Belle* Collaboration), *Phys. Rev. Lett.* **99**, 131803 (2007).
- [4] R. Aaij *et al.* (*LHCb* Collaboration), *Phys. Rev. Lett.* **110**, 222001 (2013).
- [5] N. A. Tornqvist, *Phys. Lett. B* **590**, 209 (2004).
- [6] C. Cawlfeld *et al.* (*CLEO* Collaboration), *Phys. Rev. Lett.* **98** 092002 (2007).
- [7] R. Aaij *et al.* (*LHCb* Collaboration), *JHEP* **1306**, 065 (2013).
- [8] J. P. Lees *et al.* (*BABAR* Collaboration), *Nucl. Instr. Methods Phys. Res., Sect. A* **726**, 203 (2013).
- [9] B. Aubert *et al.* (*BABAR* Collaboration), *Nucl. Instr. Methods Phys. Res., Sect. A* **479** 1 (2002).
- [10] B. Aubert *et al.* (*BABAR* Collaboration), in press in *Nucl. Instr. Methods Phys. Res., Sect. A* arXiv:1305.3560 (2013).
- [11] D. Asner, Ph.D. thesis, University of California at Santa Barbara 2000.
- [12] J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86** 010001 (2012).
- [13] J. P. Lees *et al.* (*BABAR* Collaboration), arXiv:1304.5009.
- [14] E. Barberio and Z. Was, *Comput. Phys. Commun.* **79**, 291 (1994).
- [15] A. Denisov, A. Zhelamkov, Y. Ivanov, L. Lapina, P. Levchenko, *et al.*, *JETP Lett.* **54** 558 (1991).
- [16] K. Gall, E. Austin, J. Miller, F. O’Brien, B. Roberts *et al.*, *Phys. Rev. Lett.* **60**, 186 (1988).