

NO ν A and T2K: The race for the neutrino mass hierarchy

Olga Mena¹, Hiroshi Nunokawa² and Stephen Parke¹

¹ *Theoretical Physics Department, Fermi National Accelerator Laboratory
P.O.Box 500, Batavia, IL 60510, USA and*

² *Departamento de Física, Pontifícia Universidade Católica do Rio de Janeiro,
C. P. 39071, 22452-9790, Rio de Janeiro, Brazil*

(Dated: February 2, 2008)

Abstract

The determination of the ordering of the neutrino masses (the hierarchy) is probably a crucial prerequisite to understand the origin of lepton masses and mixings and to establish their relationship to the analogous properties in the quark sector. Here, we follow an alternative strategy to the usual neutrino–antineutrino comparison in long baseline neutrino oscillation experiments: we exploit the combination of the neutrino-only data from the NO ν A and the T2K experiments by performing these two off-axis experiments at different distances but at the same $\langle E \rangle / L$, where $\langle E \rangle$ is the mean neutrino energy and L is the baseline. This would require a minor adjustment to the proposed off-axis angle for one or both of the proposed experiments.

PACS numbers: 14.60Pq

arXiv:hep-ph/0609011v1 1 Sep 2006

I. INTRODUCTION

During the last several years the physics of neutrinos has achieved a remarkable progress. The experiments with solar [1–6], atmospheric [7], reactor [8], and also long-baseline accelerator [9] neutrinos (improved with the recent MINOS [10] results [11]) have provided compelling evidence for the existence of neutrino oscillations, implying non zero neutrino masses. The present data¹ require two large (θ_{12} and θ_{23}) and one small (θ_{13}) angles in the neutrino mixing matrix [14], and at least two mass squared differences, $\Delta m_{ji}^2 \equiv m_j^2 - m_i^2$ (where m_j 's are the neutrino masses), one driving the atmospheric (Δm_{31}^2) and the other one the solar (Δm_{21}^2) neutrino oscillations. The mixing angles θ_{12} and θ_{23} control the solar and the dominant atmospheric neutrino oscillations, while θ_{13} is the angle limited by the data from the CHOOZ and Palo Verde reactor experiments [15, 16].

The Super-Kamiokande (SK) [7] and K2K [9] data are well described in terms of dominant $\nu_\mu \rightarrow \nu_\tau$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau$) vacuum oscillations. The MINOS Collaboration has recently reported the first neutrino oscillation results from 1.27×10^{20} protons on target exposure of the MINOS far detector [11]. The value of the oscillation parameters from MINOS are consistent with the ones from K2K, as well as from SK data. A recent global fit [17] (see also Ref. [18]) provides the following 3σ allowed ranges for the atmospheric mixing parameters

$$|\Delta m_{31}^2| = (1.9 - 3.2) \times 10^{-3} \text{eV}^2, \quad 0.34 < \sin^2 \theta_{23} < 0.68 . \quad (1)$$

The sign of Δm_{31}^2 , $\text{sign}(\Delta m_{31}^2)$, cannot be determined with the existing data. The two possibilities, $\Delta m_{31}^2 > 0$ or $\Delta m_{31}^2 < 0$, correspond to two different types of neutrino mass ordering: normal hierarchy and inverted hierarchy. Both possibilities are illustrated in Fig. 1, extracted from Ref. [19]. In addition, information on the octant in which θ_{23} lies, if $\sin^2 2\theta_{23} \neq 1$, is beyond the reach of present experiments.

The 2-neutrino oscillation analysis of the solar neutrino data, including the results from the complete salt phase of the Sudbury Neutrino Observatory (SNO) experiment [6], in combination with the KamLAND spectrum data [20], shows that the solar neutrino oscilla-

¹ We restrict ourselves to a three-family neutrino analysis. The unconfirmed LSND signal [12] cannot be explained within this context and might require additional light sterile neutrinos or more exotic explanations. The ongoing MiniBooNE experiment [13] is going to test the oscillation explanation of the LSND result.

tion parameters lie in the low-LMA (Large Mixing Angle) region, with best fit values [17] $\Delta m_{21}^2 = 7.9 \times 10^{-5} \text{ eV}^2$ and $\sin^2 \theta_{12} = 0.30$.

A combined 3-neutrino oscillation analysis of the solar, atmospheric, reactor and long-baseline neutrino data [17] constrains the third mixing angle to be $\sin^2 \theta_{13} < 0.041$ at the 3σ C.L. However, the bound on $\sin^2 \theta_{13}$ is dependent on the precise value of Δm_{31}^2 .

The future goals for the study of neutrino properties is to precisely determine the already measured oscillation parameters and to obtain information on the unknown ones: namely θ_{13} , the CP-violating phase δ and the type of neutrino mass hierarchy (or equivalently $\text{sign}(\Delta m_{31}^2)$).

In this paper we concentrate on the extraction of the neutrino mass hierarchy by combining the Phase I (neutrino-data only) of the long-baseline ν_e appearance experiments T2K [21] and NO ν A [22], both exploiting the off-axis technique². We start by presenting the general formalism in Sec. II. In Sec. III we describe the strategy and the experimental setups. In Sec. IV the experimental configuration is optimized in order to maximize the sensitivity to the mass hierarchy extraction and we present our results for different possible experimental combinations. Finally, in Sec. V, we make our final remarks. For our analysis, unless otherwise stated, we will use a representative value of $|\Delta m_{31}^2| = 2.4 \times 10^{-3} \text{ eV}^2$ and $\sin^2 2\theta_{23} = 1$. For the solar oscillation parameters Δm_{21}^2 and θ_{12} , we will use the best fit values quoted earlier in this section.

II. THE OSCILLATION PROBABILITY $\nu_\mu \rightarrow \nu_e$

The mixing angle θ_{13} controls $\nu_\mu \rightarrow \nu_e$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ conversions in long-baseline ν_e appearance experiments and the $\bar{\nu}_e$ disappearance in short-baseline reactor experiments. Present and future conventional beams [10], super-beams with an upgraded proton source and a more intense neutrino flux and future reactor neutrino experiments have the possibility

² A neutrino beam with narrow energy spectrum can be produced by placing the detector off-axis, i. e., at some angle with respect to the forward direction. The resulting neutrino spectrum is very narrow in energy (nearly monochromatic, $\Delta E/E \sim 15 - 25\%$) and peaked at lower energies with respect to the on-axis one. The off-axis technique allows a discrimination between the peaked ν_e oscillation signal and the intrinsic ν_e background which has a broad energy spectrum [23].




FIG. 1: The range of probability of finding the α -flavor in the i -th mass eigenstate as indicated for the two different mass hierarchies for the best fit values of the solar and atmospheric mixing parameters, $\sin^2 2\theta_{13} = 0.1$ as the CP-violating phase, δ is varied.

to measure³, or set a stronger limit on, θ_{13} . Therefore, with the possibility of the first measurement of θ_{13} being made by a 1-to 2-km baseline reactor experiment [30], the long-baseline off-axis ν_e appearance experiments, T2K [21] and NO ν A [22], need to adjust their focus to emphasize other physics topics. The most important of these questions is the form of the mass hierarchy, normal ($\Delta m_{31}^2 > 0$) versus inverted ($\Delta m_{31}^2 < 0$), and whether or not leptonic CP violation occurs.

Consider the probability $P(\nu_\mu \rightarrow \nu_e)$ in the context of three-neutrino mixing in the presence of matter [31], represented by the matter parameter a , defined as $a \equiv G_F n_e / \sqrt{2}$, where n_e is the average electron number density over the baseline, taken to be constant throughout the present study. Defining $\Delta_{ij} \equiv \frac{\Delta m_{ij}^2 L}{4E}$, a convenient and precise approximation is obtained by expanding to second order in the following small parameters: θ_{13} , Δ_{21}/Δ_{32} , Δ_{21}/aL and Δ_{21} . The result is (details of the calculation can be found in Ref. [25], see also

³ Smaller values of this mixing angle could be accessed by very long baseline experiments such as neutrino factories [24, 25], or by β -beams [26–29] which exploit neutrinos from boosted-ion decays.

Ref. [32]):

$$\begin{aligned}
P(\nu_\mu \rightarrow \nu_e) &\simeq \\
&\left| \sin \theta_{23} \sin 2\theta_{13} \left(\frac{\Delta_{31}}{\Delta_{31} - aL} \right) \sin(\Delta_{31} - aL) e^{-i(\Delta_{32} + \delta)} + \cos \theta_{23} \sin 2\theta_{12} \left(\frac{\Delta_{21}}{aL} \right) \sin(aL) \right|^2 \\
&= \sin^2 \theta_{23} \sin^2 2\theta_{13} \left(\frac{\Delta_{31}}{\Delta_{31} - aL} \right)^2 \sin^2(\Delta_{31} - aL) \\
&+ \sin 2\theta_{13} \sin 2\theta_{23} \sin 2\theta_{12} \left(\frac{\Delta_{21}}{aL} \right) \left(\frac{\Delta_{31}}{\Delta_{31} - aL} \right) \sin(aL) \sin(\Delta_{31} - aL) \cos(\Delta_{32} + \delta) \\
&+ \cos^2 \theta_{23} \sin^2 2\theta_{12} \left(\frac{\Delta_{21}}{aL} \right)^2 \sin^2(aL), \tag{2}
\end{aligned}$$

where L is the baseline and $a \rightarrow -a$, $\delta \rightarrow -\delta$ for $P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e)$.

Suppose $P(\nu_\mu \rightarrow \nu_e) < P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e)$ for given energy and baseline, then in vacuum this implies CP violation. However, in matter, this implies CP violation only for the normal hierarchy but not necessarily for the inverted hierarchy around the first oscillation maximum. The different index of refraction for neutrinos and antineutrinos induces differences in the ν , $\bar{\nu}$ propagation that could be misinterpreted as CP violation [33]. Typically, the proposed long baseline neutrino oscillation experiments have a single far detector and plan to run with the beam in two different modes, neutrinos and antineutrinos. In principle, by comparing the probability of neutrino and antineutrino flavor conversion, the values of the CP-violating phase δ and of $\text{sign}(\Delta m_{31}^2)$ could be extracted. However, different sets of values of CP-conserving and violating parameters, $(\theta_{13}, \theta_{23}, \delta, \text{sign}(\Delta m_{31}^2))$, lead to the same probabilities of neutrino and antineutrino conversion and provide a good description of the data at the same confidence level. This problem is known as the problem of degeneracies in the neutrino parameter space [34–38] and severely affects the sensitivities to these parameters in future long-baseline experiments. Many strategies have been advocated to resolve this issue. Some of the degeneracies might be eliminated with sufficient energy or baseline spectral information [25]. In practice, statistical errors and realistic efficiencies and backgrounds limit considerably the capabilities of this method. Another detector [35, 39–42] or the combination with another experiment [43–51] would, thus, be necessary⁴.

The use of only a neutrino beam could help in resolving the type of hierarchy when

⁴ New approaches which exploit other neutrino oscillations channels such as muon neutrino disappearance have been proposed [52] for determining the type of hierarchy. They require very precise measurements of the neutrino oscillation disappearance oscillation parameters though.

two different long-baselines are considered [42–44, 53, 54]. It was shown in ref. [44] that if the $\langle E \rangle/L$ for the two different experiments is approximately the same then the allowed regions for the two hierarchies are disconnected and thus this method for determining the hierarchy is free of degeneracies.⁵ Naively, we can understand this method in the following way for $\sin^2 2\theta_{13} > 0.01$: assume that matter effects are negligible for the short baseline, then at the same $\langle E \rangle/L$, if the oscillation probability at the long baseline is larger than the oscillation probability at the short baseline, one can conclude that the hierarchy is normal, since matter effects enhance the neutrino oscillation probabilities for the normal hierarchy. For the inverted hierarchy the oscillation probability for the long baseline is suppressed relative to the short baseline. This will be explained in some detail in the next section.

III. OUR STRATEGY: ONLY NEUTRINO RUNNING AND TWO DETECTORS

Following the line of thought developed by Minakata, Nunokawa and Parke [44], we exploit the neutrino data from two experiments at different distances and at different off-axis locations. The off-axis location of the detectors and the baseline must be chosen such that the $\langle E \rangle/L$ is the same for the two experiments. Here we explain the advantages of such a strategy versus the commonly exploited neutrino-antineutrino comparison.

Suppose we compute the oscillation probabilities $P(\nu_\mu \rightarrow \nu_e)$ and $P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e)$ for a given set of oscillation parameters and the CP-phase δ is varied between 0 and 2π : we obtain a closed CP trajectory (an ellipse) in the bi-probability space of neutrino and antineutrino conversion [36]. Matter effects are responsible for the departure of the center of the ellipses from the diagonal line in the bi-probability plane for the both cases of normal and inverted hierarchy, see Fig. 2 (a), where we have illustrated the case for $E = 2.0$ GeV and $L = 810$ km. The distance between the center of the ellipse for the normal hierarchy (lower blue) and that for the inverted hierarchy (upper red) is governed by the size of the matter effects. Notice that the ellipses overlap for a significant fraction of values of the CP-phase δ for every allowed value of $\sin^2 2\theta_{13}$. This indicates that, generically, a measurement of the probability of conversion for neutrinos and antineutrinos cannot uniquely determine the type of hierarchy in a single experiment. This makes the determination of $\text{sign}(\Delta m_{31}^2)$ extremely difficult, i. e.,

⁵ This method, while optimal for extracting the neutrino mass hierarchy, is not the most powerful one for extracting the CP violating phase δ for which antineutrino running would be, in general, necessary.




FIG. 2: (a) Bi-probability neutrino-antineutrino ellipses at the far distances (810 km) for normal (lower blue) and inverted (upper red) hierarchies. (b) Same as (a) but in the bi-neutrino plane at the short (295 km) and far distances (810 km).

the $\text{sign}(\Delta m_{31}^2)$ -extraction is not free of degeneracies.

In the case of bi-probability plots for neutrino-neutrino modes at different distances (which will be referred as near (N) and far (F)), the CP-trajectory is also elliptical (see Fig. 2 (b)). The overlap of the two bands, which implies the presence of a degeneracy of the type of hierarchy with other parameters, is controlled by the difference in the slopes and the width of the bands. Using the fact that matter effects are small ($aL \ll \Delta_{31}$), we can perform a perturbative expansion and assuming that the $\langle E \rangle / L$ of the near and far experiments is the same⁶, at first order, the ratio of the slopes reads [44]

$$\frac{\alpha_+}{\alpha_-} \simeq 1 + 4(a_N L_N - a_F L_F) \left(\frac{1}{\Delta_{31}} - \frac{1}{\tan(\Delta_{31})} \right), \quad (3)$$

where α_+ and α_- are the slopes of the center of the ellipses as one varies θ_{13} for normal and inverted hierarchies, and a_F and a_N are the matter parameters, L_F and L_N are the baselines for the two experiments⁷. The separation between the center of the ellipses for the two

⁶ The reason for this choice of $\langle E \rangle / L$ is explained in the next paragraph.

⁷ Notice that although we are using the constant density approximation, a_F and a_N are different because the average density depends on the baseline. For T2K (NO ν A) we use an average density times electron fraction equal to $1.15 (1.40) \text{ g} \cdot \text{cm}^{-3}$.

hierarchies increases as the difference in the matter parameter times the path length for the two experiments increases. Also, since $(\Delta^{-1} - \cot \Delta)$ is a monotonically increasing function, we conclude that the smaller the energy, the larger the ratio of slopes, at least for the same $\langle E \rangle/L$.

However the width of the ellipses is crucial: even when the separation between the central axes of the two regions is substantial if the ellipses for the normal and inverted hierarchy overlap the hierarchy cannot be resolved for values of CP phase, δ , for which there is overlap. The width of the ellipses is controlled by the difference in the $\langle E \rangle/L$ of the two experiments. For fixed θ_{13} the ellipses are flat if there exists two different δ which give the same oscillation probability for both the near, P_N , and far, P_F , detectors. That is, if

$$P_N(\theta_{13}, \delta) = P_N(\theta_{13}, \delta') \quad \text{and} \quad P_F(\theta_{13}, \delta) = P_F(\theta_{13}, \delta') \quad (4)$$

has non-trivial solutions for δ and δ' . Eqs. (4) are satisfied when $\cos(\Delta + \delta) = \cos(\Delta + \delta')$ for both the near and far experiments simultaneously. Both equations can only be satisfied when $\Delta_N = \Delta_F$, that is the two experiments have the same $\langle E \rangle/L$.

Thus, we have two conditions to satisfy to optimize the determination of the neutrino mass hierarchy:

- (a) maximize the difference in the factor (aL) for both experiments and
- (b) minimize the ellipses width by performing the two experiments at the same $\langle E \rangle/L$.

The most promising way to optimize the sensitivity to the hierarchy with relatively near term data is therefore to focus on the neutrino running mode and to exploit the Phase I data of the long-baseline off-axis ν_e appearance experiments, T2K and NO ν A. T2K utilizes a steerable neutrino beam from J-PARC and Super-Kamiokande and maybe eventually Hyper-Kamiokande as the far detector. The beam will peak at 0.65 GeV with the detector off-axis by an angle of 2.5° at 295 km. For this configuration the matter effects are small but not negligible [55]. NO ν A proposes to use the Fermilab NuMI beam with a baseline of 810 km with a 30 kton low density tracking calorimeter with an efficiency of 24%. Such a detector would be located 12 km off-axis distance from the beam center at $L = 810$ km (corresponding to 0.85° off-axis angle), resulting in a mean neutrino energy of 2.0 GeV. Matter effects are quite significant for NO ν A. Therefore, the condition (a) is satisfied, since $(aL)_{\text{NO}\nu\text{A}} \simeq 3(aL)_{\text{T2K}}$. What about the condition (b)? A back-of-the-envelope calculation

indicates us that the current off-axis angles are not such that $\langle E \rangle / L$ of the two experiments will be the same. However, by placing the detector(s) at slightly different off-axis angle(s), one can arrange that the $\langle E \rangle / L$ of the two experiments to be exactly the same. This strategy would only need half of the time of data taking (because we avoid the antineutrino running), when compared to the standard one (i.e. running in neutrinos and antineutrinos at a fixed energy, $\langle E \rangle$, and baseline, L).

IV. OPTIMIZING THE NO ν A AND T2K DETECTOR LOCATIONS

In this section we present what could be achieved if NO ν A and T2K setups are carefully chosen, focusing on the physics potential of the combination of their future data. We define the Phase I of the experiments as follows. For the T2K experiment, we consider 5 years of neutrino running and SK as the far detector with a fiducial mass of 22.5 kton and 70% detection efficiencies. For the NO ν A experiment, we assume 6.5×10^{20} protons on target per year, 5 years of neutrino running and the detector described in the previous section.

We present in Figs. 3 the bi-event neutrino-neutrino plots, fixing the off-axis location of the T2K detector to 2.5° but varying the off-axis distance of the NO ν A far detector, where the number of events is computed by integrating the neutrino flux convoluted with the cross section, the oscillation probability and the efficiency over an energy window of 1 GeV centered around the mean neutrino energy (for both NO ν A and T2K experiments). If the location of the NO ν A far detector is the proposed one, i.e. 12 km off-axis, the width of the ellipses is not negligible and the determination of the hierarchy would become a really difficult task even if nature has been kind in its choice of δ , see Fig. 3 (a). As we anticipate in the previous section, to maximize the difference in the product aL , see Eq. (3), does not ensure a degeneracy-free hierarchy measurement, due to the impact of the ellipses width. Notice, however, that the width of the ellipses is minimal when the $\langle E \rangle / L$ of both experiments is almost the same: this occurs when the NO ν A far detector is located 14 km off axis, see Fig. 3 (c), for which the mean energy $\langle E \rangle \sim 1.7$ GeV. If the NO ν A far detector is placed 16 km off-axis, the $\langle E \rangle / L$ of the T2K and NO ν A experiments is no longer the same and the ellipses width has grown with respect to the 14 km off-axis setup. *The most important lesson we have learned is that if the T2K beam is off-axis by an angle of 2.5° , the ideal location for the NO ν A detector to optimize the hierarchy extraction would be*




FIG. 3: (a) Bi-event neutrino-neutrino ellipses at the $NO\nu A$ and $T2K$ experiments for normal (lower blue) and inverted (upper red) hierarchies. From bottom up, the dashed ellipses correspond to $\sin^2 2\theta_{13}$ varying from 0.01 to 0.2 with a stepsize of 0.01 and the solid ellipses illustrate the case $\sin^2 2\theta_{13} = 0.1$. The $T2K$ far detector is located off-axis by an angle of 2.5° . The $NO\nu A$ detector is placed 12 km off-axis. (b,c,d) Same as (a) but with the $NO\nu A$ detector located at 13 km, 14 km and 16 km off-axis, respectively.

14 km off-axis. Fig. 3 shows the potential of the combination of the data from the Phase I (only neutrinos) of the $T2K$ and $NO\nu A$ experiments, without relying on future second off-axis detectors (placed at a shorter distance or at the second oscillation maximum) and future upgraded proton luminosities (and/or detection technologies providing almost perfect detection efficiencies).

The disadvantage is that the configuration at 14 km off-axis implies a 30% loss in statistics

with respect to the *default* configuration at 12 km off-axis. Consequently, the best strategy would be the following: one could tune the $\langle E \rangle / L$ of the two experiments to be the same by changing the T2K off axis-angle, since the design of the T2K experiment could in principle allow for such a modification, keeping the NO ν A off-axis distance fixed to 12 km which corresponds to $\langle E \rangle = 2.0$ GeV. In order to reproduce the same $\langle E \rangle / L$ than the corresponding one for the NO ν A experiment, the energy at T2K should be 0.73 GeV, which could be easily achieved by placing the detector 2° off-axis, for which $\langle E \rangle = 0.75$ GeV. In principle, since the configuration at 2° off-axis would also imply a gain in statistics for the T2K experiment with respect to the one at 2.5° off-axis, one would expect an improvement in the sensitivity. However, we noticed that the ratio for the two solutions Eq. (3) increases as the energy decreases. Since the configuration of NO ν A detector located at 14 km off-axis and T2K located at 2.5° off-axis provides a lower $\langle E \rangle$ than the energy for the configuration obtained by placing the NO ν A detector to its default location and moving slightly T2K to 2° off-axis, the sensitivity achieved in the former is higher than in the latter.

The data fits we perform here are based on a χ^2 analysis, where the χ^2 function reads:

$$\chi^2 = \sum_{i=T2K, NO\nu A} \left(\frac{N_i^{obs} - N_i^{th}(\sin^2 2\theta_{13}, \delta)}{\sigma_i} \right)^2, \quad (5)$$

where N^{obs} is the measured number of events, $N^{th}(\sin^2 2\theta_{13}, \delta)$ is the expected number of events for a particular choice of the hierarchy and σ refers to the statistical error corresponding to N^{th} . Our analysis includes the intrinsic ν_e background. In our plots we show the 90% confidence level regions in the $(\sin^2 2\theta_{13}, \delta)$ plane resulting from the data fits, where the 90% CL is defined as $|\chi_{min}^2(\text{wrong hierarchy}) - \chi_{min}^2(\text{true hierarchy})| > 4.61$ for 2 d.o.f [56]. It is very important to notice here that our 90% CL contours translate into the 97% CL contours for 1 d.o.f.⁸ For all the sensitivity (exclusion) plots shown in this paper, the systematic error was not included but later we will make some comments on the impact of this error on our results.

We summarize the results in Figs. 4, where we present the exclusion plots in the

⁸ In order to test whether or not the 2 d.o.f approach is more appropriate, we use a Monte Carlo technique in which we simulate 10^6 experiments for each point in the $(\sin^2 2\theta_{13}, \delta)$ parameter space and we compute the probability of distinguishing the hierarchy for each point. The result of the Monte Carlo analysis agrees with the conservative approach of 2 d.o.f but it does not reproduce the 1 d.o.f confidence levels. This Monte Carlo analysis has therefore guided us to use 2 d.o.f.

($\sin^2 2\theta_{13}, \delta$) plane for a measurement of the hierarchy at the 90% CL for the several possible combinations, assuming that nature's solution is the normal hierarchy and $\Delta m_{31}^2 = 2.4 \times 10^{-3} \text{ eV}^2$ (left panel) and $\Delta m_{31}^2 = 3.0 \times 10^{-3} \text{ eV}^2$ (right panel) (in the light of the recent MINOS results, we explore here the impact of a larger Δm_{31}^2). We show as well the corresponding CHOOZ bound for $\sin^2 2\theta_{13}$. A larger value of Δm_{31}^2 implies more statistics and, consequently, a sensitivity improvement: see Fig. 4 (b), where for the sake of illustration only three representative configurations are shown.

If both T2K and NO ν A run in their *default* configurations the combination of their future Phase I data (only neutrinos) will not contribute much to our knowledge of the neutrino hierarchy, see the solid blue line in Figs. 4. If we fix the T2K off-axis location to its *default* value of 2.5° but we change the location of the NO ν A detector to 14 km the improvement is really remarkable, see the dashed red line in Figs. 4: the sensitivity to the mass hierarchy has a milder dependence on the CP-phase δ once that the $\langle E \rangle / L$ of the two experiments is chosen to be the same. The best sensitivity to the hierarchy extraction is clearly achieved when the NO ν A experiment is at 14 km off-axis and the T2K off-axis angle is the *default* one. If the T2K off-axis angle is slightly modified to 2° , see the dotted lines in Figs. 4 it would be possible to reproduce the results from the combination of the data from T2K located at 2.5° off-axis and the NO ν A detector placed at 13 km off-axis.

If the nature's choice for the neutrino mass ordering is the inverted hierarchy, the sensitivity curves depicted in Fig. 4 (a) will be shifted but in the opposite direction, see Fig. 5 for the result from the combination of data from the Phase I of both experiments. Notice that the method presented here is independent on the octant in which θ_{23} lies (if θ_{23} is not maximal). There exists though a minor impact on the sensitivity curves since $\theta_{23} > \pi/4$ ($< \pi/4$) implies more (less) statistics, see the first term in Eq. (2).

We have explored the impact of an overall 4% systematic error. The sensitivity curves, while shifted slightly for the combination of NO ν A at 14 km off-axis and T2K at 2.5° off-axis (or NO ν A at 12 km and T2K at 2.0°), get much worse if both experiments run in their default configurations. One can understand this by making use of the bi-event neutrino-neutrino plots, see Figs. 2: a larger error bar has an enormous impact if the ellipses width is not negligible, see Figs. 2 (a).

The combination of data from an upgraded phase of the T2K and/or NO ν A experiments (by increasing the proton luminosities, the years of neutrino running and/or the mass of




FIG. 4: (a) 90% CL (2 d.o.f) hierarchy resolution for different possible combinations: the default one (T2K at an off-axis angle of 2.5° and NO ν A far detector at 12 km off-axis, in solid blue), T2K at an off-axis angle of 2.5° and NO ν A far detector at 13 km off-axis (long dash-dot red curve), T2K at an off-axis angle of 2.5° and NO ν A far detector at 14 km off-axis (short dashed red curve), T2K at an off-axis angle of 2.5° and NO ν A far detector at 16 km off-axis (three dots-three dashes blue curve) and T2K at an off-axis angle of 2° and NO ν A far detector at 12 km off-axis (dotted blue curve). We have considered the statistics corresponding to the Phase I of both experiments. The vertical dashed line indicates the 95% CL CHOOZ bound for the value of Δm_{31}^2 for the panel. (b) The same as (a) but assuming that $\Delta m_{31}^2 = 3.0 \times 10^{-3} \text{ eV}^2$ and only for the three most representative combinations: the default one (in solid blue), T2K at an off-axis angle of 2° and NO ν A far detector at 12 km off-axis (dotted blue curve) and the optimal one, that is, T2K at an off-axis angle of 2.5° and NO ν A far detector at 14 km off-axis (short dashed red curve). If one reinterprets these limits for 1 d.o.f then they correspond to the 95% CL, approximately.

the far detectors) will obviously increase the statistics and will shift the sensitivity curves depicted in Fig. 4 (a), similarly to the effect of increasing Δm_{31}^2 , see Figs. 6, where we have upgraded NO ν A and T2K experiments by increasing a factor of five their expected Phase I statistics (Phase II). Fig. 7 depicts the results from an upgraded Phase II of both experiments in the inverted hierarchy nature's choice: if the neutrino mass hierarchy is inverted, the case for the Phase II of both experiments will be stronger, especially for $\Delta m_{31}^2 = 2.4 \times 10^{-3} \text{ eV}^2$.




FIG. 5: Same as Figs. 4 but assuming that the nature's choice for the neutrino mass spectrum is the inverted hierarchy.




FIG. 6: Same as Figs. 4 but increasing the statistics of T2K and NO ν A by a factor of five.

V. CONCLUSIONS

The most promising way to extract the neutrino mass hierarchy is to make use of the matter effects in neutrino oscillations. For that purpose, the fastest way would be to exploit




FIG. 7: Same as Figs. 6 but assuming that the nature's choice for the neutrino mass spectrum is the inverted hierarchy.

the neutrino data only from two near-term long baseline ν_e appearance experiments performed at the same $\langle E \rangle / L$, provided $\sin^2 2\theta_{13}$ is within their sensitivity range or within the sensitivity range of the next-generation $\bar{\nu}_e$ disappearance reactor neutrino experiments. Such a possibility could be provided by the combination of the data from the Phase I of the T2K and NO ν A experiments. We conclude that the optimal configuration for these experiments would be 14 km off-axis for the NO ν A far detector and 2.5° off-axis for the T2K experiment. The combination of their expected results could provide a 90% confidence level (using 2 d.o.f) resolution of the neutrino mass hierarchy if $\sin^2 2\theta_{13} > 0.11$ (for $\Delta m_{31}^2 = 2.4 \times 10^{-3} \text{ eV}^2$) or if $\sin^2 2\theta_{13} > 0.07$ (for $\Delta m_{31}^2 = 3.0 \times 10^{-3} \text{ eV}^2$). A modest upgraded next Phase of both NO ν A and T2K experiments (by increasing a factor of five their expected Phase I statistics) could shift the 90% CL limits quoted above to $\sin^2 2\theta_{13} > 0.03$ (for $\Delta m_{31}^2 = 2.4 \times 10^{-3} \text{ eV}^2$) and to $\sin^2 2\theta_{13} > 0.025$ (for $\Delta m_{31}^2 = 3.0 \times 10^{-3} \text{ eV}^2$). A slightly less sensitive combination is T2K at 2° off-axis angle and NO ν A at 12 km off-axis location.

Acknowledgements

We acknowledge extensive discussions with Hisakazu Minakata. The calculations presented here made extensive use of the Fermilab-General-Purpose Computing Farms [57]. Fermilab is operated by URA under DOE contract DE-AC02-76CH03000. H. N. is supported by Fundação de Amparo à Pesquisa do Estado de Rio de Janeiro (FAPERJ) and Conselho Nacional de Ciência e Tecnologia (CNPq).

-
- [1] B. T. Cleveland *et al.*, *Astrophys. J.* **496**, 505 (1998); Y. Fukuda *et al.* [Kamiokande Collaboration], *Phys. Rev. Lett.* **77**, 1683 (1996); J. N. Abdurashitov *et al.* [SAGE Collaboration], *J. Exp. Theor. Phys.* **95**, 181 (2002); W. Hampel *et al.* [GALLEX Collaboration], *Phys. Lett. B* **447**, 127 (1999); T. A. Kirsten [GNO Collaboration], *Nucl. Phys. Proc. Suppl.* **118**, 33 (2003).
- [2] S. Fukuda *et al.* [Super-Kamiokande Collaboration], *Phys. Lett. B* **539**, 179 (2002).
- [3] Q. R. Ahmad *et al.* [SNO Collaboration], *Phys. Rev. Lett.* **87**, 071301 (2001).
- [4] Q. R. Ahmad *et al.* [SNO Collaboration], *Phys. Rev. Lett.* **89**, 011301 (2002) and *ibid.* **89**, 011302 (2002).
- [5] S. N. Ahmed *et al.* [SNO Collaboration], *Phys. Rev. Lett.* **92**, 181301 (2004).
- [6] B. Aharmim *et al.* [SNO Collaboration], *Phys. Rev. C* **72**, 055502 (2005).
- [7] Y. Ashie *et al.* [Super-Kamiokande Collaboration], *Phys. Rev. D* **71**, 112005 (2005).
- [8] K. Eguchi *et al.* [KamLAND Collaboration], *Phys. Rev. Lett.* **90**, 021802 (2003).
- [9] M. H. Ahn [K2K Collaboration], hep-ex/0606032.
- [10] E. Ables *et al.* [MINOS Collaboration], FERMILAB-PROPOSAL-0875.
- [11] MINOS Collaboration, hep-ex/0607088.
- [12] A. Aguilar *et al.* [LSND Collaboration], *Phys. Rev. D* **64**, 112007 (2001).
- [13] A. A. Aguilar-Arevalo *et al.* [MiniBooNE Collaboration],
The MiniBooNE Run Plan, available at
<http://www-boone.fnal.gov/publicpages/runplan.ps.gz>
- [14] B. Pontecorvo, *Sov. Phys. JETP* **6**, 429 (1957) [*Zh. Eksp. Teor. Fiz.* **33**, 549 (1957)] and *ibid.* **7**, 172 (1958) [*ibid.* **34** (1958) 247]; Z. Maki, M. Nakagawa and S. Sakata, *Prog. Theor. Phys.* **28**, 870 (1962).

- [15] M. Apollonio *et al.* [CHOOZ Collaboration], Phys. Lett. B **466**, 415 (1999).
- [16] F. Boehm *et al.*, Phys. Rev. Lett. **84**, 3764 (2000) and Phys. Rev. D **62**, 072002 (2000).
- [17] T. Schwetz, hep-ph/0606060.
- [18] G. L. Fogli *et al.*, arXiv:hep-ph/0608060.
- [19] O. Mena and S. J. Parke, Phys. Rev. D **69**, 117301 (2004).
- [20] T. Araki *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **94**, 081801 (2005).
- [21] Y. Hayato *et al.*, Letter of Intent, available at <http://neutrino.kek.jp/jhfnu/>
- [22] D. S. Ayres *et al.* [NOvA Collaboration], hep-ex/0503053. FERMILAB-PROPOSAL-0929, March 21, 2005. Revised NO ν A Proposal available at http://www-nova.fnal.gov/NOvA_Proposal/Revised_NOvA_Proposal.html
- [23] A. Para and M. Szleper, hep-ex/0110032.
- [24] A. De Rújula, M. B. Gavela and P. Hernández, Nucl. Phys. B **547**, 21 (1999); V. Barger *et al.*, Phys. Rev. D **62**, 013004 (2000); M. Freund *et al.*, Nucl. Phys. B **578**, 27 (2000).
- [25] A. Cervera *et al.*, Nucl. Phys. B **579**, 17 (2000) [Erratum-*ibid.* B **593**, 731 (2001)].
- [26] P. Zucchelli, Phys. Lett. B **532**, 166 (2002).
- [27] M. Mezzetto, J. Phys. G **29**, 1781 (2003) and *ibid.* **29** 1771 (2003).
- [28] For a recent study of the optimal β -beam at the CERN-SPS, see: J. Burguet-Castell *et al.*, Nucl. Phys. B **725**, 306 (2005).
- [29] J. Burguet-Castell *et al.*, Nucl. Phys. B **695**, 217 (2004); P. Huber *et al.*, Phys. Rev. D **73**, 053002 (2006); A. Donini *et al.*, hep-ph/0604229.
- [30] K. Anderson *et al.*, hep-ex/0402041; F. Ardellier *et al.*, hep-ex/0405032; E. Abouzaid *et al.*, LBNL-56599; P. Huber *et al.*, JHEP **0605**, 072 (2006).
- [31] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978); V. D. Barger, K. Whisnant, S. Pakvasa and R. J. N. Phillips, Phys. Rev. D **22**, 2718 (1980); S. P. Mikheev and A. Y. Smirnov, Sov. J. Nucl. Phys. **42**, 913 (1985); S. J. Parke, Phys. Rev. Lett. **57**, 1275 (1986); H. W. Zaglauer and K. H. Schwarzer, Z. Phys. C **40**, 273 (1988); M. C. Banuls, G. Barenboim and J. Bernabeu, Phys. Lett. B **513**, 391 (2001); M. Freund *et al.*, Nucl. Phys. B **578**, 27 (2000).
- [32] E. K. Akhmedov *et al.*, JHEP **0404**, 078 (2004);
- [33] J. Arafune, M. Koike and J. Sato, Phys. Rev. D **56**, 3093 (1997) [Erratum-*ibid.* D **60**, 119905 (1999)]; H. Minakata and H. Nunokawa, Phys. Lett. B **413**, 369 (1997); A. Donini *et al.*, Nucl. Phys. B **574**, 23 (2000).

- [34] G. L. Fogli and E. Lisi, Phys. Rev. D **54**, 3667 (1996).
- [35] J. Burguet-Castell *et al.* Nucl. Phys. B **608**, 301 (2001).
- [36] H. Minakata and H. Nunokawa, JHEP **0110**, 001 (2001).
- [37] V. Barger, D. Marfatia and K. Whisnant, Phys. Rev. D **65**, 073023 (2002);
- [38] T. Kajita, H. Minakata and H. Nunokawa, Phys. Lett. B **528**, 245 (2002); H. Minakata, H. Nunokawa and S. J. Parke, Phys. Rev. D **66**, 093012 (2002); M. Freund, P. Huber and M. Lindner, Nucl. Phys. B **615**, 331 (2001); P. Huber, M. Lindner and W. Winter, Nucl. Phys. B **645**, 3 (2002); A. Donini, D. Meloni and S. Rigolin, JHEP **0406**, 011 (2004); M. Aoki, K. Hagiwara and N. Okamura, Phys. Lett. B **606**, 371 (2005); O. Yasuda, New J. Phys. **6**, 83 (2004); O. Mena and S. J. Parke, Phys. Rev. D **72**, 053003 (2005).
- [39] H. Minakata and H. Nunokawa, Phys. Lett. B **413**, 369 (1997).
- [40] A. Donini, D. Meloni and P. Migliozi, Nucl. Phys. B **646**, 321 (2002); D. Autiero *et al.*, Eur. Phys. J. C **33**, 243 (2004).
- [41] V. Barger, D. Marfatia and K. Whisnant, Phys. Rev. D **66**, 053007 (2002).
- [42] M. Ishitsuka *et al.*, Phys. Rev. D **72**, 033003 (2005); K. Hagiwara, N. Okamura and K. i. Senda, Phys. Lett. B **637**, 266 (2006).
- [43] P. Huber, M. Lindner and W. Winter, Nucl. Phys. B **654**, 3 (2003).
- [44] H. Minakata, H. Nunokawa and S. J. Parke, Phys. Rev. D **68**, 013010 (2003).
- [45] V. Barger, D. Marfatia and K. Whisnant, Phys. Lett. B **560**, 75 (2003).
- [46] J. Burguet-Castell *et al.*, Nucl. Phys. B **646**, 301 (2002); K. Whisnant, J. M. Yang and B. L. Young, Phys. Rev. D **67**, 013004 (2003); P. Huber *et al.*, Nucl. Phys. B **665**, 487 (2003); P. Huber *et al.*, Phys. Rev. D **70**, 073014 (2004); A. Donini *et al.*, Nucl. Phys. B **710**, 402 (2005); A. Donini, E. Fernández-Martínez and S. Rigolin, Phys. Lett. B **621**, 276 (2005).
- [47] O. Mena and S. J. Parke, Phys. Rev. D **70**, 093011 (2004).
- [48] P. Huber, M. Maltoni and T. Schwetz, Phys. Rev. D **71**, 053006 (2005).
- [49] O. Mena, Mod. Phys. Lett. A **20**, 1 (2005).
- [50] P. Huber *et al.*, hep-ph/0606119.
- [51] A. Blondel *et al.*, hep-ph/0606111.
- [52] A. de Gouvea, J. Jenkins and B. Kayser, Phys. Rev. D **71**, 113009 (2005); H. Nunokawa, S. Parke and R. Z. Funchal, Phys. Rev. D **72**, 013009 (2005); A. de Gouvea and W. Winter, Phys. Rev. D **73**, 033003 (2006); H. Minakata *et al.*, hep-ph/0607284.

- [53] O. Mena Requejo, S. Palomares-Ruiz and S. Pascoli, Phys. Rev. D **72**, 053002 (2005).
- [54] O. Mena, S. Palomares-Ruiz and S. Pascoli, Phys. Rev. D **73**, 073007 (2006).
- [55] M. Aoki, K. Hagiwara and N. Okamura, Phys. Lett. B **554**, 121 (2003).
- [56] W. M. Yao et al., J. Phys. G **33**, 1 (2006).
- [57] M. Albert *et al.*, “The Fermilab Computing Farms in 2001–2002,” FERMILAB-TM-22091.