

Limits on the diffuse flux of ultra-high energy neutrinos using the Pierre Auger Observatory

Sergio Pastor^{a,1}, for the Pierre Auger Collaboration^b

^a*Institut de Física Corpuscular (CSIC-Universitat de València), Ed. Institutos de Investigación, Apartado 22085, 46071 Valencia, Spain*

^b*Av. San Martín Norte 304 (5613) Malargüe, Prov. de Mendoza, Argentina*

Abstract

The surface detector array of the Pierre Auger Observatory is sensitive to ultra-high energy neutrinos in the cosmic radiation. These particles can interact close to ground, both through charged and neutral currents in the atmosphere (down-going) and, for tau neutrinos, through the Earth-skimming mechanism (up-going) where a tau lepton is produced in the Earth crust that can emerge and decay in the atmosphere. Both types of neutrino induced-events produce an inclined shower that can be identified by the presence of a broad time structure of signals in the water-Cherenkov detectors. Using data collected from the surface detector array of the Pierre Auger Observatory, we present the corresponding limits on the diffuse flux of ultra-high energy neutrinos.

Keywords: Pierre Auger Observatory, Ultra-high energy neutrinos

1. Introduction

The existence of cosmic neutrinos with energies in the EeV range and above is required by the observation of ultra-high energy cosmic rays (UHECRs). Although the nature of the very energetic cosmic radiation and its production mechanisms are still uncertain [1, 2], all models of UHECRs predict neutrino fluxes from the decay of charged pions, produced either in interactions of the cosmic rays in their sources, or in their subsequent interactions with background radiation fields. The so-called cosmogenic or GZK neutrinos [3] are produced in the interaction of UHE protons with the cosmic microwave background (CMB). This flux of cosmogenic neutrinos is to some extent uncertain since it depends on the composition of primary UHECRs and on the nature, cosmological evolution and spatial distribution of the sources (see e.g. [4]).

The observation of UHE neutrinos would open a new window to the universe, since they can give information on regions that are otherwise hidden from observation by large amounts of matter in the field of view. Moreover, UHE neutrinos travel unaffected by magnetic

fields and, hence, they essentially maintain the direction of their production places. The detection of very energetic cosmic neutrinos is the aim of many experiments which employ different techniques, from neutrino telescopes such as IceCube or Antares to experiments like ANITA, that searches for radio waves from extra-terrestrial neutrino interactions. One of the detection techniques is based on the observation of extensive air showers (EAS) in the atmosphere initiated by UHE neutrinos, which could be detectable by a large ground detector such as the Pierre Auger Observatory.

In this contribution, we describe the sensitivity of the surface detector array of the Pierre Auger Observatory to UHE neutrinos with energies around EeV and larger. We explain the identification criteria used to distinguish neutrino-induced showers from those initiated by UHECRs, such as protons or heavy nuclei. The analysis of Auger data reveals no neutrino candidates, leading to stringent limits on the diffuse flux of UHE neutrinos.

2. Detection of UHE neutrinos with the Pierre Auger Observatory

Neutrinos, even at very high energies, present a low interaction probability which means that a large amount of matter is needed to detect these elusive particles. In the case of UHE cosmic neutrinos, the Earth atmosphere is the target where the primary particles inter-

Email address: pastor@ific.uv.es (Sergio Pastor)

¹Work supported by the Spanish grants FPA2008-00319 and CSD2009-00064 (MICINN) and PROMETEO/2009/091 (Generalitat Valenciana)




Figure 1: Types of neutrino interactions that initiate particle showers in the atmosphere. In neutral current (NC) processes the scattered neutrino carries away a large fraction of the primary energy and only part is transferred to the shower. A similar case is that of charged current (CC) interactions involving UHE ν_μ 's, where the outgoing muon usually decays under the ground and does not produce an EAS. Instead, the emerging charged leptons essentially carry all the initial energy in CC processes with a ν_e or a ν_τ , with the possibility of a double-bang shower if the τ travels a long distance before decaying.




Figure 2: Simplified picture of the different types of particle showers induced by UHE neutrinos compared to proton-induced showers. The surface detector of the Pierre Auger Observatory is sensitive to both up-going (UG) or Earth-skimming tau neutrinos and down-going neutrinos (DG) of all flavours.

49 act producing an EAS that can be detected in experi- 71
 50 ments such as the Pierre Auger Observatory. These are 72
 51 the so-called down-going (DG) events, where neutrinos 73
 52 of all flavours interact at any atmospheric depth through 74
 53 charged or neutral currents, as shown in Figure 1, and 75
 54 develop an EAS. Instead, only UHE tau neutrinos can 76
 55 lead to up-going (UG) or Earth-skimming events in an 77
 56 efficient way, when they interact in the Earth crust and 78
 57 produce a tau lepton that can emerge and decay in the 79
 58 atmosphere. If the decay of such tau leptons occurs 80
 59 in flight over the detector array, they may initiate de- 81
 60 tectable air showers [5, 6]. Tau neutrinos are expected 82
 61 to be suppressed in the production processes, because 83
 62 they do not appear in the charged pion decay chain. 84
 63 However, the effect of neutrino flavour oscillations over 85
 64 cosmological distances modifies the initial composition 86
 65 and leads to approximately equal fluxes for all flavours. 87
 66 DG and UG neutrino-induced showers are depicted in 88
 67 Figure 2.

68 The main background for the detection of EAS indu- 90
 69 ced by UHE neutrinos is the particle showers initi- 91
 70 ated by UHECRs: protons or heavy nuclei and possi- 92

bly photons. UHECRs interact high in the atmosphere, 90
 producing particle showers that contain muons and an 91
 electromagnetic component of electron, positrons and 92
 photons. This latter component reaches a maximum at 93
 an atmospheric depth of order 800 g cm^{-2} , extinguish- 94
 ing gradually within the next 1000 g cm^{-2} . Thus after 95
 roughly a couple of vertical atmospheric depths only 96
 high energy muons survive. In the first stages of devel- 97
 opment, while the electromagnetic component devel- 98
 ops, the time spread of the particles in the shower 99
 front is large ($\sim \mu\text{s}$). When the shower becomes old, 100
 most of the particles in the shower front, the high energy 101
 muons, arrive in a short time window ($\sim 100 \text{ ns}$). As a 102
 consequence very inclined showers induced by UHE- 103
 CRs in the upper atmosphere reach the ground as a thin 104
 and flat front of muons accompanied by an electromag- 105
 netic halo, which is produced by bremsstrahlung, pair 106
 production, and muon decays, and has a time structure 107
 very similar to that of the muons. On the other hand, 108
 if a shower is induced by a particle that interacts deep 109
 in the atmosphere (a deep neutrino interaction in air, or 110
 a tau decay), its electromagnetic component could hit 111

the ground and give a distinct broad signal in time. The panels in Figure 3 represent these various possibilities.

The Pierre Auger Observatory [7] has been designed to measure UHECRs with unprecedented precision. It employs the two available techniques to detect EAS, namely, arrays of surface particle detectors and telescopes that detect fluorescence radiation. The surface detector array (SD) of the Southern Auger Observatory, recently completed in the Mendoza province (Argentina), consists of 1600 water Cherenkov tanks arranged in a hexagonal grid of 1.5 km that covers an effective area of 3000 km². Each cylindrical tank of 10 m² surface contains purified water, 1.2 m deep, and is instrumented with three 9" photomultiplier tubes (PMT) sampled by 40 MHz Flash Analog Digital Converters (FADCs). Each tank is regularly monitored and calibrated in units of vertical equivalent muon (VEM) corresponding to the signal produced by a muon traversing the tank vertically [8].

The signal in each station of the Auger SD is digitized using FADCs with a 25 ns time resolution, which allows unambiguous distinction between the narrow signals induced by muons and the broad signals induced by the electromagnetic component. Thus the time structure and shape of the FADC traces, characterized by several observables [9], can help us to discriminate stations hit by an EAS in the early stages of development or by an old EAS. A set of conditions has been designed and optimized to select showers induced by UHE neutrinos, either UG or DG events, rejecting those induced by UHECRs. These conditions constitute the neutrino identification criteria for SD events, described in the next section. The fluorescence detectors can also be used for neutrino searches but the nominal 10% duty cycle of this technique reduces the sensitivity. Here only the data collected with the SD of the Pierre Auger Observatory is used to search for UHE neutrinos.

3. Neutrino identification criteria

A large set of simulations of UHE neutrinos forced to interact deep in the atmosphere were produced in order to characterize the signal that their induced EAS would produce at the SD array. The first interaction of the primary neutrino in the air, either NC or CC, was simulated using HERWIG [10], while the AIRES code [11] was used for the EAS development. For ν_τ -induced showers, the TAUOLA package [12] was used to simulate the τ decay and obtain the secondary particles and their energies.

Shower simulations were performed including the geographic conditions of the Auger site (e.g. geomag-




Figure 3: Upper panel: an inclined EAS induced by a proton interacting high in the atmosphere. The electromagnetic component is absorbed and only the muons reach the array of ground detectors. Middle panel: a primary UHE neutrino can initiate a deep inclined shower whose early region has a significant electromagnetic component at the detector level. Lower panel: a "young" shower can be also produced by an up-going tau lepton produced by the interaction of an Earth-skimming neutrino.

netic field) for different zenith angles. For UG showers, primary energies ranged from 10^{17} to 3×10^{20} eV at zenith angles between 90.1° and 95.9° and at an altitude of the decay point above the Pierre Auger Observatory up to 2500 m. In this case we also simulate the propagation of up-going tau neutrinos through the Earth crust. In the case of DG neutrinos, simulations were performed at zenith angles $\theta = 75^\circ, 80^\circ, 85^\circ, 87^\circ, 88^\circ$ and 89° and random azimuth angles between 0° and 360° and different hadronic models. The primary neutrinos were forced to interact at different slant depths measured from the ground up to a maximum value depending on θ . Finally the response of the SD array is simulated in detail using the Offline simulation package [13]. The two sets of Monte Carlo (MC) neutrino simulations were used to estimate the expected neutrino signal and the detection efficiency for both UG and DG events.

The main criterion to identify young and very inclined showers consists of looking for broad time signals in the SD stations. Two different sets of identification conditions were designed to search for UG and DG neutrinos.




Figure 4: Distribution of the values of the area over peak for the first triggering station (AoP_1 , left) and the product of AoP of the first four triggering stations (right), as a function of the zenith angle for real data and MC simulated neutrinos.




Figure 5: Distribution of the Fisher discriminant in linear (left) and logarithmic (right) scale for real Auger data in the training period (1 Jan 2004 - 31 Oct 2007) and MC simulated down-going neutrinos for events with multiplicity $7 \leq N \leq 11$.

3.1. Up-going neutrinos

Young showers are expected to trigger detector stations with broad signals releasing a so-called Time Over Threshold (ToT) trigger [14]. Counting ToTs stations can help identifying young showers, and a cut in the value of the area of the signal over its peak (AoP, where the peak corresponds to the maximum measured current of recorded trace at a single water-Cherenkov detector) is applied to reject accidental muons hitting a station that could mimic a ToT local trigger. After trace cleaning, very inclined showers are identified with the elongation of their footprint, defined by the ratio of length (L) over width (W) of the shower pattern on ground, requiring $L/W > 5$, and the mean apparent velocity $\langle V \rangle$, expected to be compatible with the speed of light for quasi-horizontal showers, in the range $(0.29, 0.31) \text{ m ns}^{-1}$ with an r.m.s. scatter below 0.08 m ns^{-1} . Finally compact configurations of selected ToTs complete the expected picture of young ν_τ -induced shower footprints.

For more details, we refer the reader to refs. [15, 16].

3.2. Down-going neutrinos

DG neutrino events are also young and inclined showers, but the wider range of zenith angles requires identification criteria different from those applied to UG neutrinos, as described in [17]. For this purpose data collected with the Auger SD between 1 Jan 2004 and 31 Oct 2007 (about 1.2 years of the full SD array) was used as "training" data. Showers that trigger the SD array but arrived during periods with instabilities in data acquisition were excluded. After that the FADC traces are cleaned to remove segments that are due to accidental muons not belonging to the shower but arriving close in time with the shower front. Moreover, if 2 or more segments with areas comparable to each other appear in a trace the station is classified as ambiguous and it is not used. Then a selection of the stations actually belonging to the event is done based on space-time compatibility

among them. Events with less than 4 tanks passing the level 2 trigger algorithm [7] are rejected. This sample is then searched for inclined events requiring that the triggered tanks have elongated patterns on the ground, with a cut $L/W > 3$. The average speed $\langle V \rangle$ measured between pairs of triggered stations is required to be compatible with that expected in a simple planar model of the shower front in an inclined event with $\theta \geq 75^\circ$, allowing for some spread due to fluctuations ($\langle V \rangle \leq 0.313 \text{ m ns}^{-1}$). Only events with reconstructed zenith angle $\theta \geq 75^\circ$ are selected. Exactly the same set of conditions is applied to the simulated neutrinos.

The sample of inclined events is searched for young showers using observables characterizing the time duration of the FADC traces in the early region of the event. To optimize their discrimination power we applied the Fisher discriminant method [18] to the training data, dominated by nucleonic showers, and to the Monte Carlo (MC) simulations that are exclusively composed of neutrino-induced showers. Given two populations of events – nucleonic inclined showers and ν -induced showers in our case – characterized by a set of observables, the Fisher method produces a linear combination of the various observables (the Fisher discriminant f) so that the separation between the means of f in the two samples is maximized, while the quadratic sum of the r.m.s. of f in each of them is minimized. Since events with a large number of tanks or multiplicity N are different from events with small multiplicity the sample of training data is divided into 3 sub-samples corresponding to events with number of tanks $4 \leq N \leq 6$, $7 \leq N \leq 11$ and $N \geq 12$, and a Fisher discriminant is obtained using each of the sub-samples as training data. We use 10 discriminant variables of the Fisher estimator: the AoP and its square of the first 4 tanks in each event, their product, and a global early-late asymmetry. In Figure 4 the distributions of two of these discriminant variables are shown as a function of the zenith angle for neutrino simulated showers and real inclined events. One can see a clear separation between the two samples.

In Figure 5 we present the distribution of the Fisher discriminant for the training data and DG neutrino simulations with multiplicity $7 \leq N \leq 11$. Again both samples are well separated. The expected number of background events can be computed by extrapolating the exponential tail of the distribution of the data. This allows us to find a cut-off value f_{cut} for each of the sub-samples, so that we expect less than one background event every 20 years above its value. Events with $f > f_{\text{cut}}$ are considered to be neutrino candidates. These cuts reject all real events in the training data samples while keeping a

significant fraction of the MC neutrino simulations [17].

4. Exposure and limits on UHE neutrinos

Auger data have been analyzed to look for candidate events that fulfilled the selection criteria for Earth-skimming UHE tau neutrinos, as described in [15, 16]. This analysis has been updated with data equivalent to 2 years of the full Auger SD in [9, 17], where we also describe how we have applied the selection procedure and values of f_{cut} for the identification of DG neutrino events to real data from 1 Nov 2007 to 28 Feb 2009 (~0.8 years of the full SD array), i.e. after the training period mentioned above.

Over the period analyzed, no candidate events were found for either UG or DG neutrinos. Based on this, the Pierre Auger Observatory data can be used to place the corresponding limits on the diffuse flux of UHE neutrinos. For this purpose the total exposure of the Auger SD must be evaluated, which involves folding the SD array aperture with the interaction probability and the identification efficiency ϵ , and integrating in time taking into account changes in the array configuration due to the installation of new stations and instabilities in data taking.

For both UG and DG neutrinos, the identification efficiency depends on the position of the shower in the surface covered by the array and the time through the instantaneous configuration of the array. The efficiency for UG neutrinos is also a function of the τ energy and the altitude above ground of the central part of the shower h_c (defined at 10 km after the decay point [19]). Instead, for DG neutrinos ϵ depends on the primary energy and the depth along the atmosphere at which the neutrino interacts, as well as on the neutrino flavour and type of interaction (CC or NC), since the different combinations of both induce different type of showers. The efficiencies for each case were obtained through MC simulations of the development of the shower in the atmosphere and the simulation of the surface detector array.

The Auger exposure to UHE neutrinos was calculated using purely MC techniques and also integrating the neutrino identification efficiencies ϵ over the whole parameter space, for the details see [9, 15, 16, 17]. Several sources of systematic uncertainties have been taken into account and their effect on the exposure evaluated. Here we do not include the full list, but we note that the main systematic uncertainty for UG neutrinos is the calculation of τ energy losses [16], while for DG the dominating source is the neutrino cross section.

Once the exposure has been calculated, a limit on the flux of UHE neutrinos can be obtained assuming a




Figure 6: Integrated upper limits (90% C.L.) from the Pierre Auger Observatory on the diffuse flux of UHE neutrinos from the analysis of the equivalent of 2 (0.8) years of the full SD array for UG (DG) neutrinos, together with limits from other experiments. For comparison, a computation of the flux of cosmogenic neutrinos from Ref. [26] is included.




Figure 7: Differential upper limits (90% C.L.) on the diffuse flux of UHE neutrinos from the same Auger data as in the previous figure. The sensitivity of the Pierre Auger Observatory peaks at EeV neutrino energies.

known shape. For a $f(E) = k \cdot E^{-2}$ differential neutrino flux we have obtained a 90% C.L. limit on the all-flavour neutrino flux using DG showers [17]

$$k < 3.2 \times 10^{-7} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} . \quad (1)$$

The corresponding limit on Earth-skimming UG neutrinos is

$$k < 4.7 \times 10^{-8} \text{ GeV cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} , \quad (2)$$

which updates the limits published in [15, 16]. It is worth to mention that the topography around the Southern Site of the Pierre Auger Observatory enhances the flux of secondary tau leptons and could improve the above limit up to a factor of $\sim 20\%$.

In Figure 6 the two Auger limits on the integrated UHE neutrino flux are shown, compared with the bounds from other experiments: AMANDA-IceCube [20, 21], RICE [22], HiRes [23, 24] and ANITA-II [25]. Finally, we include in Figure 7 the Auger limits in differential format in order to emphasize the range in energies at which the sensitivity of the Pierre Auger Observatory to UHE neutrinos peaks. These differential limits were calculated as $2.44/E_\nu \varepsilon(E_\nu)$, where $\varepsilon(E_\nu)$ is the exposure. In both figures the gray line corresponds to one example of the expected cosmogenic neutrino flux, computed in [26] and consistent with HiRes and Fermi-LAT measurements.

5. Conclusions

The surface detector array of the Pierre Auger Observatory is sensitive to the EAS initiated by UHE neutrinos in the atmosphere, either down-going or Earth-skimming neutrinos. We have shown how using MC simulations and training data the identification criteria for UHE neutrinos can be found. The data collected by the Auger SD has been used to present upper limits to the diffuse flux of UHE neutrinos, providing at present the most sensitive bound on neutrinos at EeV energies, which is the most relevant energy to explore the predicted fluxes of cosmogenic neutrinos. The Pierre Auger Observatory will keep taking data for about 20 years over which the bound will improve significantly if no neutrino candidate is found.

References

- [1] F. Halzen and D. Hooper, Rep. Prog. Phys. 65 (2002) 1025.
- [2] P. Bhattacharjee and G. Sigl, Phys. Rep. 327 (2000) 109.
- [3] V.S. Berezinsky et al. Phys. Lett. B 28 (1969) 423.
- [4] L.A. Anchordoqui et al., Phys. Rev. D 76 (2007) 123008.
- [5] D. Fargion, Astrophys. J. 570 (2002) 909.

- [6] A. Letessier-Selvon, AIP Conf. Proc. 566 (2001) 157.
- [7] J. Abraham et al. [Pierre Auger Coll.], Nucl. Instr. and Meth. A 523 (2004) 50.
- [8] X. Bertou et al. [Pierre Auger Coll.], Nucl. Instr. and Meth. A 568 (2006) 839.
- [9] D. Góra [Pierre Auger Coll.], Proceedings of the 31st Int. Cosmic Ray Conference 2009, Łódź, Poland.
- [10] G. Corcella et al., JHEP 01 (2001) 010.
- [11] S. Sciutto, <http://www.fisica.unlo.edu.ar/auger/aires/>
- [12] S. Jadach et al., Comput. Phys. Commun. 76 (1993) 361.
- [13] S. Argiro et al., Nucl. Instrum. Meth. A 580 (2007) 1485.
- [14] P. Billoir and O. Blanch Bigas, Nucl. Phys. Proc. Suppl. 168 (2007) 225.
- [15] J. Abraham et al. [Pierre Auger Coll.], Phys. Rev. Lett. 100 (2008) 211101.
- [16] J. Abraham et al. [Pierre Auger Coll.], Phys. Rev. D 79 (2009) 102001.
- [17] J. Tiffenberg [Pierre Auger Coll.], Proceedings of the 31st Int. Cosmic Ray Conference 2009, Łódź, Poland.
- [18] R. Fisher, Annals Eugen. 7 (1936) 179.
- [19] X. Bertou et al., Astropart. Phys. 17 (2002) 183.
- [20] M. Ackermann et al. [IceCube Coll.], Astrophys. J. 675 (2008) 1014.
- [21] R. Abbasi et al. [IceCube Coll.], preprint arXiv:1009.1442
- [22] I. Kravchenko et al. [Rice Coll.], Phys. Rev. D 73 (2006) 082002.
- [23] K. Martens [HiRes Coll.], preprint arXiv:0707.4417
- [24] R.U. Abbasi et al. [HiRes Coll.], Astrophys. J. 684 (2008) 790.
- [25] P.W. Gorham et al. [ANITA Coll.], Phys. Rev. D 82 (2010) 022004.
- [26] M. Ahlers et al., Astropart. Phys. 34 (2010) 106.