

Cross sections for the reactions $e^+e^- \rightarrow K_S^0 K_L^0$, $K_S^0 K_L^0 \pi^+ \pi^-$, $K_S^0 K_S^0 \pi^+ \pi^-$, and $K_S^0 K_S^0 K^+ K^-$ from events with initial-state radiation

J. P. Lees, V. Poireau, and V. Tisserand

*Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP),
Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France*

E. Grauges

Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

A. Palano^{ab}

INFN Sezione di Bari^a; Dipartimento di Fisica, Università di Bari^b, I-70126 Bari, Italy

G. Eigen and B. Stugu

University of Bergen, Institute of Physics, N-5007 Bergen, Norway

D. N. Brown, L. T. Kerth, Yu. G. Kolomensky, M. J. Lee, and G. Lynch

Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

H. Koch and T. Schroeder

Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

C. Hearty, T. S. Mattison, J. A. McKenna, and R. Y. So

University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

A. Khan

Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

V. E. Blinov^{ac}, A. R. Buzykaev^a, V. P. Druzhinin^{ab}, V. B. Golubev^{ab}, E. A. Kravchenko^{ab},
A. P. Onuchin^{ac}, S. I. Serednyakov^{ab}, Yu. I. Skovpen^{ab}, E. P. Solodov^{ab}, and K. Yu. Todyshev^{ab}
*Budker Institute of Nuclear Physics SB RAS, Novosibirsk 630090^a,
Novosibirsk State University, Novosibirsk 630090^b,
Novosibirsk State Technical University, Novosibirsk 630092^c, Russia*

A. J. Lankford and M. Mandelkern

University of California at Irvine, Irvine, California 92697, USA

B. Dey, J. W. Gary, and O. Long

University of California at Riverside, Riverside, California 92521, USA

C. Campagnari, M. Franco Sevilla, T. M. Hong, D. Kovalskiy, J. D. Richman, and C. A. West

University of California at Santa Barbara, Santa Barbara, California 93106, USA

A. M. Eisner, W. S. Lockman, W. Panduro Vazquez, B. A. Schumm, and A. Seiden

University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

D. S. Chao, C. H. Cheng, B. Echenard, K. T. Flood, D. G. Hitlin, T. S. Miyashita, P. Ongmongkolkul, and F. C. Porter
California Institute of Technology, Pasadena, California 91125, USA

R. Andreassen, Z. Huard, B. T. Meadows, B. G. Pushpawela, M. D. Sokoloff, and L. Sun

University of Cincinnati, Cincinnati, Ohio 45221, USA

P. C. Bloom, W. T. Ford, A. Gaz, J. G. Smith, and S. R. Wagner

University of Colorado, Boulder, Colorado 80309, USA

R. Ayad* and W. H. Toki
Colorado State University, Fort Collins, Colorado 80523, USA

B. Spaan
Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany

D. Bernard and M. Verderi
Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS/IN2P3, F-91128 Palaiseau, France

S. Playfer
University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

D. Bettoni^a, C. Bozzi^a, R. Calabrese^{ab}, G. Cibinetto^{ab}, E. Fioravanti^{ab},
 I. Garzia^{ab}, E. Luppi^{ab}, L. Piemontese^a, and V. Santoro^a
INFN Sezione di Ferrara^a; Dipartimento di Fisica e Scienze della Terra, Università di Ferrara^b, I-44122 Ferrara, Italy

A. Calcaterra, R. de Sangro, G. Finocchiaro, S. Martellotti,
 P. Patteri, I. M. Peruzzi,[†] M. Piccolo, M. Rama, and A. Zallo
INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy

R. Contri^{ab}, M. Lo Vetere^{ab}, M. R. Monge^{ab}, S. Passaggio^a, C. Patrignani^{ab}, and E. Robutti^a
INFN Sezione di Genova^a; Dipartimento di Fisica, Università di Genova^b, I-16146 Genova, Italy

B. Bhuyan and V. Prasad
Indian Institute of Technology Guwahati, Guwahati, Assam, 781 039, India

M. Morii
Harvard University, Cambridge, Massachusetts 02138, USA

A. Adametz and U. Uwer
Universität Heidelberg, Physikalisches Institut, D-69120 Heidelberg, Germany

H. M. Lacker
Humboldt-Universität zu Berlin, Institut für Physik, D-12489 Berlin, Germany

P. D. Dauncey
Imperial College London, London, SW7 2AZ, United Kingdom

U. Mallik
University of Iowa, Iowa City, Iowa 52242, USA

C. Chen, J. Cochran, and S. Prell
Iowa State University, Ames, Iowa 50011-3160, USA

H. Ahmed
Physics Department, Jazan University, Jazan 22822, Kingdom of Saudi Arabia

A. V. Gritsan
Johns Hopkins University, Baltimore, Maryland 21218, USA

N. Arnaud, M. Davier, D. Derkach, G. Grosdidier, F. Le Diberder,
 A. M. Lutz, B. Malaescu,[‡] P. Roudeau, A. Stocchi, and G. Wormser
*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11,
 Centre Scientifique d'Orsay, F-91898 Orsay Cedex, France*

D. J. Lange and D. M. Wright
Lawrence Livermore National Laboratory, Livermore, California 94550, USA

J. P. Coleman, J. R. Fry, E. Gabathuler, D. E. Hutchcroft, D. J. Payne, and C. Touramanis
University of Liverpool, Liverpool L69 7ZE, United Kingdom

A. J. Bevan, F. Di Lodovico, and R. Sacco
Queen Mary, University of London, London, E1 4NS, United Kingdom

G. Cowan
University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom

J. Bougher, D. N. Brown, and C. L. Davis
University of Louisville, Louisville, Kentucky 40292, USA

A. G. Denig, M. Fritsch, W. Gradl, K. Griessinger, A. Hafner, and K. R. Schubert
Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany

R. J. Barlow[§] and G. D. Lafferty
University of Manchester, Manchester M13 9PL, United Kingdom

R. Cenci, B. Hamilton, A. Jawahery, and D. A. Roberts
University of Maryland, College Park, Maryland 20742, USA

R. Cowan and G. Sciolla
Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA

R. Cheaib, P. M. Patel,[¶] and S. H. Robertson
McGill University, Montréal, Québec, Canada H3A 2T8

N. Neri^a and F. Palombo^{ab}
INFN Sezione di Milano^a; Dipartimento di Fisica, Università di Milano^b, I-20133 Milano, Italy

L. Cremaldi, R. Godang,^{**} P. Sonnek, and D. J. Summers
University of Mississippi, University, Mississippi 38677, USA

M. Simard and P. Taras
Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7

G. De Nardo^{ab}, G. Onorato^{ab}, and C. Sciacca^{ab}
*INFN Sezione di Napoli^a; Dipartimento di Scienze Fisiche,
 Università di Napoli Federico II^b, I-80126 Napoli, Italy*

M. Martinelli and G. Raven
NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands

C. P. Jessop and J. M. LoSecco
University of Notre Dame, Notre Dame, Indiana 46556, USA

K. Honscheid and R. Kass
Ohio State University, Columbus, Ohio 43210, USA

E. Feltresi^{ab}, M. Margoni^{ab}, M. Morandin^a, M. Posocco^a, M. Rotondo^a, G. Simi^{ab}, F. Simonetto^{ab}, and R. Stroili^{ab}
INFN Sezione di Padova^a; Dipartimento di Fisica, Università di Padova^b, I-35131 Padova, Italy

S. Akar, E. Ben-Haim, M. Bomben, G. R. Bonneaud, H. Briand,
 G. Calderini, J. Chauveau, Ph. Leruste, G. Marchiori, and J. Ocariz
*Laboratoire de Physique Nucléaire et de Hautes Energies,
 IN2P3/CNRS, Université Pierre et Marie Curie-Paris6,
 Université Denis Diderot-Paris7, F-75252 Paris, France*

M. Biasini^{ab}, E. Manoni^a, S. Pacetti^{ab}, and A. Rossi^a
INFN Sezione di Perugia^a; Dipartimento di Fisica, Università di Perugia^b, I-06123 Perugia, Italy

C. Angelini^{ab}, G. Batignani^{ab}, S. Bettarini^{ab}, M. Carpinelli^{ab,††}, G. Casarosa^{ab}, A. Cervelli^{ab}, M. Chrzaszcz^{ab},
 F. Forti^{ab}, M. A. Giorgi^{ab}, A. Lusiani^{ac}, B. Oberhof^{ab}, E. Paoloni^{ab}, A. Perez^a, G. Rizzo^{ab}, and J. J. Walsh^a
INFN Sezione di Pisa^a; Dipartimento di Fisica, Università di Pisa^b; Scuola Normale Superiore di Pisa^c, I-56127 Pisa, Italy

D. Lopes Pegna, J. Olsen, and A. J. S. Smith
Princeton University, Princeton, New Jersey 08544, USA

R. Faccini^{ab}, F. Ferrarotto^a, F. Ferroni^{ab}, M. Gaspero^{ab}, L. Li Gioi^a, and G. Piredda^a
*INFN Sezione di Roma^a; Dipartimento di Fisica,
 Università di Roma La Sapienza^b, I-00185 Roma, Italy*

C. Büniger, S. Dittrich, O. Grünberg, T. Hartmann, M. Hess, T. Leddig, C. Voß, and R. Waldi
Universität Rostock, D-18051 Rostock, Germany

T. Adye, E. O. Olaiya, and F. F. Wilson
Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom

S. Emery and G. Vasseur
CEA, Irfu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France

F. Anulli,^{‡‡} D. Aston, D. J. Bard, C. Cartaro, M. R. Convery, J. Dorfan, G. P. Dubois-Felsmann,
 W. Dunwoodie, M. Ebert, R. C. Field, B. G. Fulsom, M. T. Graham, C. Hast, W. R. Innes, P. Kim,
 D. W. G. S. Leith, P. Lewis, D. Lindemann, S. Luitz, V. Luth, H. L. Lynch, D. B. MacFarlane,
 D. R. Muller, H. Neal, M. Perl, T. Pulliam, B. N. Ratcliff, A. Roodman, A. A. Salnikov,
 R. H. Schindler, A. Snyder, D. Su, M. K. Sullivan, J. Va'vra, W. J. Wisniewski, and H. W. Wulsin
SLAC National Accelerator Laboratory, Stanford, California 94309 USA

M. V. Purohit, R. M. White,^{§§} and J. R. Wilson
University of South Carolina, Columbia, South Carolina 29208, USA

A. Randle-Conde and S. J. Sekula
Southern Methodist University, Dallas, Texas 75275, USA

M. Bellis, P. R. Burchat, and E. M. T. Puccio
Stanford University, Stanford, California 94305-4060, USA

M. S. Alam and J. A. Ernst
State University of New York, Albany, New York 12222, USA

R. Gorodeisky, N. Guttman, D. R. Peimer, and A. Soffer
Tel Aviv University, School of Physics and Astronomy, Tel Aviv, 69978, Israel

S. M. Spanier
University of Tennessee, Knoxville, Tennessee 37996, USA

J. L. Ritchie, A. M. Ruland, R. F. Schwitters, and B. C. Wray
University of Texas at Austin, Austin, Texas 78712, USA

J. M. Izen and X. C. Lou
University of Texas at Dallas, Richardson, Texas 75083, USA

F. Bianchi^{ab}, F. De Mori^{ab}, A. Filippi^a, and D. Gamba^{ab}
INFN Sezione di Torino^a; Dipartimento di Fisica, Università di Torino^b, I-10125 Torino, Italy

L. Lanceri^{ab} and L. Vitale^{ab}

INFN Sezione di Trieste^a; Dipartimento di Fisica, Università di Trieste^b, I-34127 Trieste, Italy

F. Martinez-Vidal, A. Oyanguren, and P. Villanueva-Perez
IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain

J. Albert, Sw. Banerjee, A. Beaulieu, F. U. Bernlochner, H. H. F. Choi, G. J. King, R. Kowalewski,
M. J. Lewczuk, T. Lueck, I. M. Nugent, J. M. Roney, R. J. Sobie, and N. Tasneem
University of Victoria, Victoria, British Columbia, Canada V8W 3P6

T. J. Gershon, P. F. Harrison, and T. E. Latham
Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

H. R. Band, S. Dasu, Y. Pan, R. Prepost, and S. L. Wu
University of Wisconsin, Madison, Wisconsin 53706, USA

We study the processes $e^+e^- \rightarrow K_s^0 K_L^0 \gamma$, $K_s^0 K_L^0 \pi^+ \pi^- \gamma$, $K_s^0 K_s^0 \pi^+ \pi^- \gamma$, and $K_s^0 K_s^0 K^+ K^- \gamma$, where the photon is radiated from the initial state, providing cross section measurements for the hadronic states over a continuum of center-of-mass energies. The results are based on 469 fb⁻¹ of data collected with the *BABAR* detector at SLAC. We observe the $\phi(1020)$ resonance in the $K_s^0 K_L^0$ final state and measure the product of its electronic width and branching fraction with about 3% uncertainty. We present a measurement of the $e^+e^- \rightarrow K_s^0 K_L^0$ cross section in the energy range from 1.06 to 2.2 GeV and observe the production of a resonance at 1.67 GeV. We present the first measurements of the $e^+e^- \rightarrow K_s^0 K_L^0 \pi^+ \pi^-$, $K_s^0 K_s^0 \pi^+ \pi^-$, and $K_s^0 K_s^0 K^+ K^-$ cross sections, and study the intermediate resonance structures. We obtain the first observations of J/ψ decay to the $K_s^0 K_L^0 \pi^+ \pi^-$, $K_s^0 K_s^0 \pi^+ \pi^-$, and $K_s^0 K_s^0 K^+ K^-$ final states.

PACS numbers: 13.66.Bc, 14.40.-n, 13.25.Jx

I. INTRODUCTION

The idea to use electron-positron annihilation events with initial-state radiation (ISR) to study processes with energies below the nominal e^+e^- center-of-mass ($E_{c.m.}$) energy was outlined in Ref. [1]. The possibility of exploiting ISR to measure low-energy cross sections at high-luminosity ϕ and B factories is discussed in Refs. [2–4], and motivates the study described in this paper. This is of particular interest because of a three-standard-deviation discrepancy between the current measured value of the muon anomalous magnetic moment ($g-2$) and that predicted by the Standard Model [5], in which hadronic loop contributions are obtained from experimental e^+e^- annihilation cross sections at low $E_{c.m.}$ energies. The study of ISR events at B factories provides

independent results over a continuum of energy values for hadronic cross sections in this energy region and also contributes to the investigation of low-mass resonance spectroscopy.

Studies of the ISR processes $e^+e^- \rightarrow \mu^+ \mu^- \gamma$ [6, 7] and $e^+e^- \rightarrow X_h \gamma$, using data of the *BABAR* experiment at SLAC, where X_h represents any of several exclusive multihadron final states, have been reported previously. The studied final states include: charged hadron pairs $\pi^+ \pi^-$ [7], $K^+ K^-$ [8], and $p \bar{p}$ [9]; four or six charged mesons [10–12]; charged mesons plus one or two π^0 mesons [11–14]; and a K_s^0 plus charged mesons [15]. Together, these demonstrate good detector efficiency for events of this kind, and well understood tracking, particle identification, and π^0 and K_s^0 reconstruction.

This paper reports measurements of the $K_s^0 K_L^0$, $K_s^0 K_L^0 \pi^+ \pi^-$, $K_s^0 K_s^0 \pi^+ \pi^-$, and $K_s^0 K_s^0 K^+ K^-$ final states, produced in conjunction with a hard photon, that is assumed to result from ISR. Candidate K_s^0 decays are reconstructed in the $\pi^+ \pi^-$ decay mode. This is the first ISR measurement from *BABAR* that includes K_L^0 mesons, which we detect via their nuclear interactions in the electromagnetic calorimeter. We use the $e^+e^- \rightarrow \gamma \phi \rightarrow \gamma K_s^0 K_L^0$ reaction to measure the K_L^0 detection efficiency directly from the data. The $e^+e^- \rightarrow K_s^0 K_L^0$ cross section is measured from threshold to 2.2 GeV. For the other final states, we measure cross sections from threshold to 4 GeV, study the internal structure of the events, and perform the first measurements of their J/ψ branching fractions. Together with our previous measurements [8, 11], these results provide a much more com-

*Now at the University of Tabuk, Tabuk 71491, Saudi Arabia

†Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

‡Now at Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Paris, France

§Now at the University of Huddersfield, Huddersfield HD1 3DH, UK

¶Deceased

**Now at University of South Alabama, Mobile, Alabama 36688, USA

††Also with Università di Sassari, Sassari, Italy

‡‡Also with INFN Sezione di Roma, Roma, Italy

§§Now at Universidad Técnica Federico Santa María, Valparaiso, Chile 2390123

plete understanding of the $K\bar{K}$, $K\bar{K}\pi\pi$, and $K\bar{K}K\bar{K}$ final states in e^+e^- annihilations.

II. THE BABAR DETECTOR AND DATA SET

The data used in this analysis were collected with the BABAR detector at the PEP-II asymmetric-energy e^+e^- storage ring. The total integrated luminosity used is 468.6 fb^{-1} [16], which includes data collected at the $\Upsilon(4S)$ resonance (424.7 fb^{-1}) and at a c.m. energy 40 MeV below this resonance (43.9 fb^{-1}).

The BABAR detector is described in detail elsewhere [17]. Charged particles are reconstructed using the BABAR tracking system, which comprises the silicon vertex tracker (SVT) and the drift chamber (DCH) inside the 1.5 T solenoid. Separation of pions and kaons is accomplished by means of the detector of internally reflected Cherenkov light and energy-loss measurements in the SVT and DCH. The hard ISR photon, photons from π^0 decays, and K_L^0 are detected in the electromagnetic calorimeter (EMC). Muon identification, provided by the instrumented flux return, is used to select the $\mu^+\mu^-\gamma$ final state.

In order to study the detector acceptance and efficiency, we have developed a special package of simulation programs for radiative processes based on the approach suggested by Kühn and Czyż [18]. Multiple collinear soft-photon emission from the initial e^+e^- state is implemented with the structure-function technique [19, 20], while additional photon radiation from the final-state particles is simulated using the PHOTOS package [21]. The precision of the radiative simulation does not contribute more than 1% uncertainty to the efficiency calculation.

The four-meson final states are generated according to a phase-space distribution. We simulate the $K_S^0 K_L^0 \gamma$ channel using a model that includes the $\phi(1020)$ and two additional resonances, fitted to all available $e^+e^- \rightarrow K_S^0 K_L^0$ cross section measurements [22–26], which cover the range from threshold up to about 2.5 GeV. Samples of roughly five times the number of expected events are generated for each final state and processed through the detector response simulation [27]. These events are then reconstructed using the same software chain as the data. Variations in detector and background conditions are taken into account.

We also simulate a number of background processes. Based on our experience with final states including kaons, we consider the ISR processes $K_S^0 K_L^0(\phi)\eta$, $K_S^0 K^\pm \pi^\mp$, $K_S^0 K^\pm \pi^\mp \pi^0$, $K_S^0 K_L^0 \pi^+ \pi^- \pi^0$, and $K_S^0 K_L^0 \pi^0$, with normalizations based on our previous measurements and isospin relations. In addition, we generate a large sample of the as-yet unobserved final state $K_S^0 K_L^0 \pi^0 \pi^0 \gamma$, which is a potential background. We also simulate several non-ISR backgrounds, including $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) events using the *Jetset* 7.4 [28] generator, and $e^+e^- \rightarrow \tau^+\tau^-$ events using the *KORALB* [29] generator.




FIG. 1: The $\pi^+\pi^-$ invariant mass distribution for the selected K_S^0 candidates for the data (points) and simulation (histogram). The vertical lines indicate the signal region.

III. THE ISR PHOTON AND K_S^0 SELECTION

Photons are reconstructed as clusters of energy deposits in contiguous crystals of the EMC. We consider the cluster in the event with the highest energy in the e^+e^- c.m. frame, and require ISR event candidates to contain a cluster with $E_{\text{c.m.}}^\gamma > 3 \text{ GeV}$, which we denote as the ISR photon. The ISR photon detection efficiency has been studied using $\mu\mu\gamma$ events [7], and we apply a polar-angle-dependent correction of typically $-1.5 \pm 0.5\%$ to the simulated efficiency.

In these events, we reconstruct K_S^0 candidates decaying to two charged pions from pairs of oppositely charged tracks not identified as electrons. They must have a well reconstructed vertex between 0.2 and 40.0 cm in radial distance from the beam axis, and their total momentum must be consistent with originating from the interaction region. The $m(\pi^+\pi^-)$ invariant mass distribution for these K_S^0 candidates is shown in Fig. 1 for both data (points) and a simulation (histogram) containing only genuine K_S^0 . The background level is relatively low and we select candidates in the $482 < m(\pi^+\pi^-) < 512 \text{ MeV}/c^2$ mass range (vertical lines on Fig. 1), and use the sidebands $472\text{--}482$ and $512\text{--}522 \text{ MeV}/c^2$ to estimate the contribution from non- K_S^0 backgrounds.

A few thousand events (about 1% of the total number of events) have more than one selected K_S^0 candidate, and we use these to study the $K_S^0 K_S^0 \pi^+ \pi^-$ and $K_S^0 K_S^0 K^+ K^-$ final states. Considering only the “best” K_S^0 candidate, with $m(\pi^+\pi^-)$ closest to the nominal [30] K_S^0 mass, we also include these events in the $K_S^0 K_L^0$ and $K_S^0 K_L^0 \pi^+ \pi^-$ measurements. The K_S^0 detection efficiency has been studied very carefully at BABAR, with data-MC

differences in the efficiency determined as a function of the K_s^0 direction and momentum. We apply a correction event by event, which introduces an overall correction $+1.1 \pm 1.0\%$ to the number of K_s^0 .

We also require the event to contain exactly zero or two tracks that are consistent with originating from the interaction region, excluding those in the selected K_s^0 candidate(s). Any number of additional tracks and EMC clusters is allowed.

IV. K_L^0 DETECTION AND EFFICIENCY

The decay length of the K_L^0 meson is large, and the probability to detect a K_L^0 decay in the DCH is low. Instead, we look for a cluster in the EMC resulting from the interaction of a K_L^0 with a nucleus in the EMC material. Such clusters are indistinguishable from photon-induced clusters, and give poor resolution on the K_L^0 energy.

In this section, we describe the use of a clean sample of $e^+e^- \rightarrow \phi\gamma$, $\phi \rightarrow K_s^0 K_L^0$ events to optimize our selection of K_L^0 clusters and measure their detection efficiency and angular resolution. In sections VI and VII, we describe the use of the selected K_L^0 candidate clusters to study the ϕ resonance and measure the $e^+e^- \rightarrow K_s^0 K_L^0$ cross section above the ϕ region, respectively.

A. The $e^+e^- \rightarrow \phi\gamma \rightarrow K_s^0 K_L^0 \gamma$ process

Using the four-momenta of the best selected K_s^0 , the ISR photon, and the initial electron and positron, we can calculate the recoil-mass-squared,

$$m_{\text{rec}}^2 = (E_0 - E_\gamma - E_{K_s^0})^2 - (\vec{p}_0 - \vec{p}_\gamma - \vec{p}_{K_s^0})^2, \quad (1)$$

where $E_0 = E^+ + E^-$ and $\vec{p}_0 = \vec{p}^+ + \vec{p}^-$ are the energy and total momentum vector of the initial e^+e^- system, E_γ and \vec{p}_γ (with $E_\gamma \equiv |\vec{p}_\gamma|$) are the energy and momentum vector of the photon, and $E_{K_s^0}$ and $\vec{p}_{K_s^0}$ are the energy and momentum vector of the K_s^0 candidate. The presence of the reaction $e^+e^- \rightarrow K_s^0 K_L^0 \gamma$ would be evident as a peak in the m_{rec} distribution at the mass of the K_L^0 .

Due to the large uncertainty of the measured ISR photon energy, the calculated value of m_{rec} also has a large uncertainty. However, if we assume the reaction $e^+e^- \rightarrow \gamma\phi(1020) \rightarrow \gamma K_s^0 K_L^0$, we can calculate the constrained ISR photon energy E_γ^c according to:

$$E_\gamma^c = \frac{E_0^2 - p_0^2 - m_\phi^2}{2(E_0 - \vec{p}_0 \cdot \vec{n}_\gamma)}, \quad (2)$$

where \vec{n}_γ is a unit vector along the ISR photon direction and m_ϕ is the ϕ meson mass [30]. Using E_γ^c instead of the measured E_γ in Eq. 1, we obtain a much better resolution on the recoil mass m_{rec}^c for genuine events of that type. The m_{rec}^c distribution for our data is shown in

Fig. 2 as the points. A simulated distribution for genuine $e^+e^- \rightarrow \gamma\phi \rightarrow \gamma K_s^0 K_L^0$ events is shown as the histogram. Selecting events with $m_{\text{rec}}^c > 0.4$ GeV/ c^2 (corresponding to $m(K_s^0 K_L^0) < 1.1$ GeV/ c^2), with the additional requirement that there be no other track within a 0.2 cm radius of the interaction point, we obtain a very clean sample of $K_s^0 K_L^0 \gamma$ events, without any need to detect the K_L^0 meson.

The non- K_s^0 background, estimated from the sidebands of the $m(\pi^+\pi^-)$ distribution in Fig. 1, contributes 0.8% of the events in Fig. 2. This background arises from $e^+e^- \rightarrow \gamma\gamma$ events in which one photon converts to a misidentified electron-positron pair. We estimate backgrounds from other ISR final states containing a real K_s^0 using the simulation. Normalized contributions to the m_{rec}^c distribution for $K_s^0 K_L^0 2\pi^0$, $K_s^0 K_L^0 \pi^0 (K^{*0} \bar{K})$, and $\phi(K_s^0 K_L^0) \eta$ are shown in Fig. 3, cumulatively, as shaded, hatched, and open histograms. The simulated backgrounds from $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$), and $e^+e^- \rightarrow \tau\tau$ events are found to be negligible.




FIG. 2: The distribution of constrained recoil mass m_{rec}^c , obtained according to Eqs. 1 and 2, for selected γK_s^0 candidates. The points represent the data, and the histogram an MC simulation of $e^+e^- \rightarrow \gamma\phi \rightarrow \gamma K_s^0 K_L^0$ events, normalized to the two most populated bins.

Fitting the simulated non- K_s^0 and ISR backgrounds with smooth functions and summing them together with the signal simulation, we obtain excellent agreement with the observed spectrum. The total background is $6.9 \pm 0.5\%$ of the selected events.

The position of the K_L^0 peak in Fig. 3 is very sensitive to both the reconstructed K_s^0 candidate mass and the assumed ϕ -meson mass (see Eqs. 1,2). There is a small 0.21 ± 0.02 MeV/ c^2 data-MC difference in the K_s^0 peak position in Fig. 1. As a cross check, we correct the data for this difference and vary m_ϕ in Eq. 2 for the data so that the experimental m_{rec}^c peak position matches




FIG. 3: The experimental m_{rec}^c distribution (points) compared with our estimated background contributions from (cumulatively): $K_s^0 K_L^0 2\pi^0 \gamma$ (shaded area), $K_s^0 K_L^0 \pi^0 \gamma$ (hatched), and $\phi(K_s^0 K_L^0) \eta \gamma$ (open histogram). The simulated signal distribution is shown as the dashed histogram and the sum of all simulated events as the solid histogram.

that of the simulation. This results in an estimate of $m_\phi = 1019.480 \pm 0.040 \pm 0.036 \text{ MeV}/c^2$, where the systematic uncertainty includes the effects of the nominal K^0 mass ($0.024 \text{ MeV}/c^2$ [30]), the K_s^0 momentum measurement in the DCH ($0.020 \text{ MeV}/c^2$), and the DCH-EMC mis-alignment ($0.018 \text{ MeV}/c^2$). This is consistent with the value, tabulated by the Particle Data Group (PDG) $m_\phi = 1019.455 \pm 0.020 \text{ MeV}/c^2$ [30].

Subtracting the non- K_s^0 and ISR-produced backgrounds, we obtain 81012 ± 285 (447434 for the MC simulation) $K_s^0 K_L^0 \gamma$ events in the ϕ mass region without requiring K_L^0 detection.

These events must satisfy our trigger and software filters, which were designed for various classes of events. We study efficiencies in data and simulation using prescaled events not subject to these filters, and obtain a correction of $(+3.9 \pm 2.3)\%$. Furthermore, the pions from K_s^0 decays in this particular reaction have a relatively large probability to overlap in the DCH, and the reconstruction efficiency for overlapping tracks is not well simulated. We introduce a $+1.5 \pm 0.6\%$ correction for this effect.

B. The K_L^0 detection efficiency

We select events with $m_{\text{rec}}^c > 0.47 \text{ MeV}/c^2$ (vertical line in Fig. 2), reducing the background level from 6.9% to 2.8%. Using the K_s^0 and ISR photon angles and momenta, we calculate the hypothetical K_L^0 direction for

each event, and look for an EMC cluster in that direction.




FIG. 4: The EMC cluster energy versus the opening angle between the measured cluster direction and the predicted K_L^0 direction for all non-ISR clusters in the data.




FIG. 5: The difference in azimuthal angle between the EMC cluster direction and the predicted K_L^0 direction for selected clusters in the data (points) and MC-simulated $K_s^0 K_L^0(\phi) \gamma$ events (histogram).

Figure 4 shows a two-dimensional plot of the EMC cluster energy versus the opening angle $\Delta\psi$ between the predicted K_L^0 direction and measured cluster direction for all clusters in the data except those assigned to the ISR photon. A clean signal is observed at high cluster energies, but the background from low-energy clusters is large. We consider clusters with energy greater than 0.2 GeV, and select the one closest to the predicted K_L^0 direction if it is within 0.5 radians. This yields K_L^0 detection probabilities of about 48% in the data and 51% in the MC simulation.

We then study the resolution in polar (θ) and azimuthal (ϕ) angle of the selected K_L^0 clusters as a function of their position in the detector and the predicted K_L^0 energy. The resolutions in the two angles are consistent, with no significant dependence on position or energy. The overall $\Delta\phi$ distributions are shown for data and simulation in Fig. 5. Good agreement is seen, with root-mean-square deviations of 0.035 radians. We use this value in the kinematic fits.

V. THE KINEMATIC FIT PROCEDURE

Each candidate event selected in Sec. III is subjected to a set of constrained kinematic fits in which the four-momenta and covariance matrices of the initial e^+e^- , the ISR photon, the best K_S^0 candidate, and the two tracks from the interaction region, if present, are taken into account. The three-momentum vectors for each particle including the photon obtained from these fits are determined with better accuracy, and are used in further calculations.

First, we consider each neutral cluster with $E > 0.2$ GeV (excluding the ISR photon) as a K_L^0 candidate, and perform a three-constraint (3C) kinematic fit under the $K_S^0 K_L^0 \gamma$ or $K_S^0 K_L^0 \pi^+ \pi^- \gamma$ hypothesis. The angular resolutions for K_L^0 clusters discussed in the previous section are used, and the K_L^0 momentum is determined in the fit. We retain the K_L^0 candidate cluster giving the best χ^2 value in each event.

We then perform a kinematic fit under the $K_S^0 K^\pm \pi^\mp \pi^0 \gamma$ hypothesis, where the cluster is assumed to be one photon from a π^0 decay, rather than a K_L^0 . Such events can enter the sample if a charged kaon is misidentified as a pion and only one photon from the π^0 decay is considered. Similarly, we perform fits under the hypotheses of the other backgrounds discussed in Sec. II, giving us additional χ^2 variables with which to suppress these processes.

We perform additional fits to the events with more than one K_S^0 candidate under the $K_S^0 K_S^0 \pi^+ \pi^- \gamma$ and $K_S^0 K_S^0 K^+ K^- \gamma$ hypotheses. For each pair of K_S^0 candidates, a four-constraint (4C) kinematic fit is performed using the four-momenta and covariance matrices of all initial- and final-state particles. The combination with the best χ^2 for each hypothesis is retained.

VI. THE $K_S^0 K_L^0$ FINAL STATE ($m(K_S^0 K_L^0) < 1.08$ GeV/ c^2)

A. Additional selection criteria and background subtraction

To study this mass region, we consider events selected as described in Sec. IV A, with $m_{\text{rec}}^c > 0.4$ GeV/ c^2 (see Fig. 2). We select a K_L^0 cluster where possible, using

the 3C fits described in Sec. V, and obtain the χ^2 distribution for the best $K_S^0 K_L^0 \gamma$ candidate shown in Fig. 6 as the points. The unshaded histogram is for the corresponding MC-simulated pure $K_S^0 K_L^0 \gamma$ events, normalized to the data in the region $\chi^2 < 10$ where we expect very low background.




FIG. 6: The three-constraint χ^2 distribution for K_S^0 -cluster- γ events in the data (points) fitted under the $K_S^0 K_L^0 \gamma$ hypothesis. The open histogram represents the same distribution for the MC-simulated signal events, normalized to the data in the region $\chi^2 < 10$, and the shaded histogram represents the estimated background in the data.

The experimental and simulated distributions are broader than a typical 3C χ^2 distribution due to multiple soft-photon emission from the initial state, which is not taken into account in the fit, but is present in both the data and simulation. The observed difference at higher χ^2 values is due to background in the data and possibly a data-MC difference in the angular uncertainty of the K_L^0 cluster.

For further analysis we require $\chi^2(K_S^0 K_L^0) < 15$ (vertical line in Fig. 6), and for these events we calculate the K_L^0 candidate mass according to Eqs. 1,2 and perform the background subtraction described in Sec. IV A. We obtain $27\,925 \pm 176$ events for the data (871 background events are subtracted) and 164 179 events for the MC-simulation, representing samples with the K_L^0 detected. Dividing by the corresponding numbers of events before K_L^0 cluster selection, we obtain K_L^0 detection efficiencies, including the effects of the kinematic fit and χ^2 selection, of 0.3447 ± 0.0017 for the data and 0.3724 ± 0.0008 for the simulation. The double ratio 0.9394 ± 0.0052 is applied as a correction factor to account for this data-MC difference. This ratio is independent of momentum and polar angle of the K_L^0 .

We use the four-vectors returned by the kinematic fit to calculate the $K_S^0 K_L^0$ invariant mass, the distribution of




FIG. 7: The $K_S^0 K_L^0$ invariant mass distribution in the data (points) and signal-MC simulation (histogram) for candidate events in the signal region of Fig. 6. The shaded histogram represents the estimated background, and the line is a smooth parametrization thereof.

which is shown in Fig. 7. The $\phi(1020)$ resonance is clearly visible, with a width of about 10 MeV, much larger than the nominal width of the resonance [30] due to the resolution of this final state. The background, estimated as described above, is shown as the shaded histogram. We fit it with a smooth, empirical function, shown as the line, and use the fit result in each bin for background subtraction.

B. Fit for the $\phi(1020)$ parameters

To obtain the parameters of the $\phi(1020)$, we fit the background-subtracted distribution in Fig. 7 with a cross section $\sigma(s)$ convolved with a resolution matrix $\text{Res}(j, i)$. In each mass bin i :

$$N_{K_S^0 K_L^0}(i) = \sum_{j=1}^{100} N_0(j) \cdot \text{Res}(j, i),$$

$$N_0(j) = \int_{m_j}^{m_{j+1}} \sigma(s) \epsilon(s) \epsilon_{K_S^0 K_L^0}^{\text{corr}} L(s) ds, \quad (3)$$

where $s = m(K_S^0 K_L^0)^2$, $\epsilon(s)$ is the simulated detection efficiency, $\epsilon_{K_S^0 K_L^0}^{\text{corr}} = 0.939 \cdot 0.985 \cdot 0.961$ is the data-MC efficiency correction factor for the χ^2 cut, track overlap, and event filter, $L(s)$ is the ISR luminosity, calculated at leading-order [4], and $N_0(j)$ is the acceptance-corrected number of events expected for bin j .

The 100×100 resolution matrix is obtained from simulation by binning the reconstructed versus simulated $K_S^0 K_L^0$ invariant mass for signal events in 1×1 MeV/ c^2




FIG. 8: The simulated distribution of differences between the reconstructed and generated $K_S^0 K_L^0$ invariant mass for the 1 MeV bin of reconstructed mass at the ϕ peak.

intervals. The distribution of differences between the reconstructed and simulated masses near 1.020 GeV/ c^2 , corresponding to a row of this matrix, is shown in Fig. 8. The $K_S^0 K_L^0$ threshold and a radiative tail are visible. We normalize each row to unit area, and introduce an additional variable Gaussian smearing σ_{add} , to account for any data-MC difference in the resolution.

We describe the cross section near the ϕ resonance using formulae discussed in detail in Refs. [22, 31]:

$$\sigma(s) = \frac{1}{s^{5/2}} \frac{q_{K_S^0 K_L^0}^3(s)}{q_{K_S^0 K_L^0}^3(m_\phi^2)} \left| \frac{\Gamma_\phi m_\phi^3 \sqrt{m_\phi \sigma_{\phi \rightarrow K_S^0 K_L^0}} / C}{D_\phi(s)} - \frac{\sqrt{\Gamma_\phi \Gamma_\rho m_\rho^3 m_\phi^2 6\pi \mathcal{B}(\rho \rightarrow e^+ e^-) \mathcal{B}(\phi \rightarrow K_S^0 K_L^0)}}{D_\rho(s)} + \frac{\sqrt{\Gamma_\phi \Gamma_\omega m_\omega^3 m_\phi^2 6\pi \mathcal{B}(\omega \rightarrow e^+ e^-) \mathcal{B}(\phi \rightarrow K_S^0 K_L^0)}}{D_\omega(s)} \right|^2 + A_{K_S^0 K_L^0}, \quad (4)$$

where $q_{K_S^0 K_L^0}(s) = \sqrt{s - 4m_{K^0}^2}$ is a threshold term, $\sigma_{\phi \rightarrow K_S^0 K_L^0}$ is the peak cross section value, $D_V(s) = s - m_V^2 + i\sqrt{s}\Gamma_V(s)$ is the propagator for a vector resonance V , $C = 0.389 \times 10^{12}$ nb MeV²/ c^4 [30], and

$$\Gamma_V(s) = \Gamma_V \cdot \sum_{V \rightarrow f} \mathcal{B}(V \rightarrow f) \frac{P_{V \rightarrow f}(s)}{P_{V \rightarrow f}(m_V^2)} \quad (5)$$

describes the energy-dependent width, and for the ϕ we use the set of final states $f = K^+ K^-$, $K_S^0 K_L^0$, $\pi^+ \pi^- \pi^0$,

and $\eta\gamma$, with corresponding branching fractions $\mathcal{B}(V \rightarrow f)$ and phase space factors $P_{V \rightarrow f}(s)$. We include the influence of the $\rho(770)$ and $\omega(782)$ resonances in the in the energy-dependent width according to the “ideal” quark model, which assumes their decay rates to $K_S^0 K_L^0$ are a factor of two lower than that of the ϕ . We use the relation

$$\sigma(V \rightarrow f) = \frac{12\pi\mathcal{B}(V \rightarrow e^+e^-)\mathcal{B}(V \rightarrow f)}{m_V^2} C \quad (6)$$

in Eq. 4 for the corresponding cross sections.

We introduce a complex constant $A_{K_S^0 K_L^0}$ to describe the contributions of higher radial excitations of the ρ , ω and ϕ mesons to the cross section, as well as any deviations from the “ideal” quark structure relations for the $\rho(770)$ and $\omega(782)$. It can be written in terms of two free parameters, a non-resonant cross section σ_{bkg} , and a phase Ψ ,

$$A_{K_S^0 K_L^0} = m_\phi^2 \sqrt{\sigma_{\text{bkg}} m_\phi / C} \cdot e^{-i\Psi}. \quad (7)$$

The fitted value of Ψ is consistent with zero, and we fix it to zero in the final fit, but propagate its fitted uncertainty as a systematic uncertainty to account for model dependence.




FIG. 9: The simulated detection efficiency $\epsilon(s)$ versus the generated $K_S^0 K_L^0$ invariant mass, calculated by dividing the number of events in the signal region of Fig. 6 by the number generated in each bin.

The detection efficiency, shown as a function of mass in Fig. 9, is obtained by dividing the number of selected MC-simulated events in each $0.001 \text{ GeV}/c^2$ mass interval by the number generated in the same interval. The mass dependence is well described by a linear fit, which we use in all calculations. This efficiency includes the geometrical acceptance of the detector for the final-state photon and the charged pions from the K_S^0 decay, the inefficiency




FIG. 10: The $K_S^0 K_L^0$ invariant mass distribution in the $\phi(1020)$ region. Only statistical uncertainties are shown. The curve represents the result of the fit described in the text.

of the detector subsystems, and event losses due to additional soft-photon emission from the initial state. It is not sensitive to the detector mass resolution.

The result of the fit is projected on the background-subtracted invariant mass distribution in Fig. 10. We obtain the resonance parameters:

$$\begin{aligned} \sigma_\phi &= 1409 \pm 33 \pm 42 \pm 15 \text{ nb}, \\ m_\phi &= 1019.462 \pm 0.042 \pm 0.050 \pm 0.025 \text{ MeV}/c^2, \\ \Gamma_\phi &= 4.205 \pm 0.103 \pm 0.050 \pm 0.045 \text{ MeV}, \\ \sigma_{\text{bkg}} &= 0.022 \pm 0.012 \text{ nb}, \end{aligned}$$

where the first uncertainties are statistical, the second systematic, and the third due to model dependence, evaluated by varying σ_{bkg} by its uncertainty.

We introduce an additional Gaussian smearing to describe an uncertainty in the detector resolution, and obtain $\sigma_{\text{add}} = 0.6 \pm 0.2 \text{ MeV}/c^2$, which improves the χ^2 of the fit in the $1.0\text{--}1.05 \text{ GeV}/c^2$ region from 59 to 53, for 51 degrees of freedom. We estimate systematic uncertainties of $0.05 \text{ MeV}/c^2$ in mass and 0.05 MeV in width from the uncertainty of the σ_{add} value. The other systematic uncertainties are summarized in Table I, along with the corrections applied to the measurements. A total correction of $+14.1 \pm 2.9\%$ is applied to the number of events. The largest contribution to the uncertainty is from the software filter, due to the limited number of available prescaled events.

Our parameter values are consistent with the most precise cross section measurement, $\sigma_\phi = 1376 \pm 24 \text{ nb}$ [22], and with the PDG values $m_\phi = 1019.455 \pm 0.020 \text{ MeV}/c^2$ and $\Gamma_\phi = 4.26 \pm 0.04 \text{ MeV}$ [30]. Since each row of the resolution matrix is normalized to unit area, the smearing procedure does not affect the total number of events,

TABLE I: Summary of corrections and systematic uncertainties for the measurement of the $e^+e^- \rightarrow K_S^0 K_L^0$ process in the ϕ resonance region.

Source	Correction	Uncertainty
Background filter efficiency	+3.9%	2.3%
Photon detection efficiency	+1.5%	0.5%
K_L^0 detection efficiency	+6.1%	0.6%
K_S^0 detection efficiency	+1.1%	1.0%
Track overlap	+1.5%	0.6%
ISR luminosity	-	0.5%
Backgrounds	-	0.5%
Radiative corrections	-	1.0%
Total (sum in quadrature)	+14.1%	2.9%

which is proportional to the product $\Gamma_\phi \sigma_0$ of the total width and peak cross section of the ϕ . Using this product as a free parameter in the fit, we obtain the product of the electronic width of the ϕ and its branching fraction to $K_S^0 K_L^0$,

$$\Gamma_{ee} \mathcal{B}_{K_S^0 K_L^0} = 0.4200 \pm 0.0033 \pm 0.0122 \pm 0.0019 \text{ keV},$$

where the first uncertainty is statistical, the second systematic, and the third due to model dependence. Using $\mathcal{B}_{K_S^0 K_L^0} = 0.342 \pm 0.004$ or $\Gamma_\phi = 4.26 \pm 0.04$ MeV from Ref. [30], we obtain $\Gamma_{ee} = 1.228 \pm 0.037 \pm 0.014$ keV or $\mathcal{B}_{ee} \mathcal{B}_{K_S^0 K_L^0} = 0.986 \pm 0.030 \pm 0.009$, respectively, where the first uncertainty is our total experimental uncertainty and the second is from the PDG tables. These values are consistent with the most recent measurement of $\Gamma_{ee} = 1.235 \pm 0.022$ keV [32], and with the PDG values of $\Gamma_{ee} = 1.27 \pm 0.04$ keV and $\mathcal{B}_{ee} \mathcal{B}_{K_S^0 K_L^0} = 1.006 \pm 0.016$ [30], and have comparable precision.

VII. THE $K_S^0 K_L^0$ FINAL STATE ($m(K_S^0 K_L^0) > 1.06$ GeV/ c^2)

In this section we consider events with $m(K_S^0 K_L^0) > 1.06$ GeV/ c^2 . Since the $e^+e^- \rightarrow K_S^0 K_L^0$ cross section drops much more rapidly with increasing mass than the background, we apply additional selection criteria compared to the criteria of Sec. VI A. In all cases, we consider K_S^0 candidates with $0.482 < m(\pi^+ \pi^-) < 0.512$ MeV/ c^2 (see Fig. 1) and use sideband data to subtract non- K_S^0 background from all studied quantities.

A. Additional selection criteria

We consider all EMC clusters except those assigned to the ISR photon and the K_L^0 as photon candidates, and combine each pair into a π^0 candidate. Figure 11 shows a scatter plot of the higher of the energies $E_{\gamma \text{max}}$ of the two photons assigned to the pair versus the corresponding diphoton mass $m_{\gamma\gamma}$. A large signal from events containing a π^0 is observed. To reduce this background,




FIG. 11: Two-dimensional plot of the higher cluster energy in a photon-candidate pair versus the corresponding diphoton mass $m_{\gamma\gamma}$ for all pairs of EMC clusters, containing neither the ISR photon nor the K_L^0 candidate.

we require $E_{\gamma \text{max}} < 0.5$ GeV (horizontal line in Fig. 11). Since signal event may contain several background clusters, this reduces the signal efficiency. We measure this loss using events in the ϕ region, where no π^0 signal is observed, but background clusters are present in both data and simulation. We find losses of 10% in the data and 7% in the simulation, and apply the 3% difference as a correction.

The 3C χ^2 distribution for the remaining candidate events with $m(K_S^0 K_L^0) > 1.06$ GeV/ c^2 is shown as the points in Fig. 12. The open histogram shows the corresponding simulated distribution for genuine $K_S^0 K_L^0 \gamma$ events, normalized to the data in the region $\chi^2 < 3$. The shaded, cross-hatched and hatched areas represent the simulated contributions from the ISR channels $\phi\eta$, $K_S^0 K_L^0 \pi^0$, and $K_S^0 K_L^0 \pi^0 \pi^0$, respectively. These channels contribute significant background, and almost entirely account for the difference between the data and signal-MC χ^2 distributions. We find no significant contribution from simulated non-ISR backgrounds.

We select events with $\chi^2(K_S^0 K_L^0) < 10$, and use events from the control region $10 < \chi^2(K_S^0 K_L^0) < 20$ (vertical lines in Fig. 12) to estimate the background in the signal region. The signal region contains 6264 data and 13 292 MC-simulated events, while the control region contains 2968 and 2670, respectively.

B. Background subtraction

To obtain any distribution of the $K_S^0 K_L^0$ signal events $N_0^d(m)$, we take the experimental events in the signal




FIG. 12: The 3C χ^2 distributions for $K_S^0 K_L^0 \gamma$ candidate events in the data (points) and signal simulation (open histogram), fitted under the $K_S^0 K_L^0$ hypothesis. The shaded, cross-hatched, and hatched areas represent the simulated contributions from the ISR $\phi\eta$, $K_S^0 K_L^0 \pi^0$, and $K_S^0 K_L^0 \pi^0 \pi^0$ channels, respectively.

region of Fig. 12, $N_s^d(m)$, and subtract the background events, taken from the control region $N_c^d(m)$, corrected for the presence of signal events, estimated from MC-simulation $N_c^{MC}(m)$:

$$N_0^d(m) = N_s^d(m) - b \cdot (N_c^d(m) - a \cdot N_c^{MC}(m)), \quad (8)$$

where $b = 1.15$ is the simulated ratio of background events in the signal and control regions, and $a = N_0^d(m)/N_s^{MC}(m)$ is a factor equalizing the number of signal and simulated events.

This procedure relies on good agreement between data and simulation in both the χ^2 and mass distributions. As noted above, the MC simulation uses a “world average” cross section, well measured below 1.4 GeV, but based on only the DM1 measurement [24], which has large statistical uncertainties, in the 1.4-2.4 GeV $E_{c.m.}$ region. We adopt an iterative procedure, in which we reweight the simulated mass distribution to match our measurement and repeat the subtraction until there is no change in the results.

Figure 13a shows the $K_S^0 K_L^0$ mass distribution for data events in the χ^2 control region of Fig. 12 as points, with the shaded histogram showing the distribution for signal MC at the final iteration. The signal contribution is not large, and the difference between the data and weighted MC-simulated distributions is scaled by $b = 1.15$ to estimate the background in the χ^2 signal region. The squares in Fig. 13b represent this background estimate in each $K_S^0 K_L^0$ invariant mass bin. We also estimate the background directly from the MC simulation of the ISR $\phi\eta$, $K_S^0 K_L^0 \pi^0$, and $K_S^0 K_L^0 2\pi^0$ processes, shown as the histogram in Fig. 13b. The two estimates agree relatively

well, but the MC simulation does not incorporate the correct mass distributions for these processes, and other unknown processes might contribute. The mass distribution after background subtraction is shown in Fig. 13c. In the 1.4-2.4 GeV/ c^2 mass region we select about 1000 events, compared with only 58 events found by the DM1 [24] experiment.

C. Simulated detection efficiency

The selection procedures applied to the data are also applied to the MC-simulated event sample. The resulting $K_S^0 K_L^0$ invariant mass distribution is shown in Fig. 14(a) for the signal and control (shaded histogram) regions. The mass dependence of the detection efficiency is obtained by dividing the number of reconstructed MC events in each mass interval by the number generated in that interval. The results are shown in Fig. 14(b). The 40 MeV/ c^2 mass intervals used are wider than the detector resolution of 10 MeV/ c^2 , but a small effect of the resolution on the efficiency is visible, due to the very steep decrease in the cross section with increasing mass.

D. The $e^+e^- \rightarrow K_S^0 K_L^0$ cross section for c.m. energies above 1.06 GeV

The cross section for e^+e^- annihilation into $K_S^0 K_L^0$ can be calculated from

$$\sigma(K_S^0 K_L^0)(E_{c.m.}) = \frac{dN_{K_S^0 K_L^0 \gamma}(E_{c.m.})}{d\mathcal{L}(E_{c.m.}) \cdot \epsilon_{K_S^0 K_L^0}^{\text{corr}} \cdot \epsilon_{K_S^0 K_L^0}^{\text{MC}}(E_{c.m.}) \cdot R}, \quad (9)$$

where: $E_{c.m.} \equiv m(K_S^0 K_L^0)$; $dN_{K_S^0 K_L^0 \gamma}$ is the number of selected $K_S^0 K_L^0$ events after background subtraction in the interval $dE_{c.m.}$; $\epsilon_{K_S^0 K_L^0}^{\text{MC}}(E_{c.m.})$ is the corresponding detection efficiency, estimated from the MC simulation with corrections to the K_S^0 and ISR photon detection efficiencies; and the factor $\epsilon_{K_S^0 K_L^0}^{\text{corr}} = 0.939 \cdot 0.961 \cdot 0.97$ takes into account the data-MC differences in K_L^0 detection and background filter efficiencies, and the energy requirement on additional photon clusters. The radiative correction R is within one percent of unity, with an estimated precision of about 1%. The differential luminosity $d\mathcal{L}(E_{c.m.})$ associated with the interval $dE_{c.m.}$ centered at an effective collision energy of $E_{c.m.}$ is calculated using the leading order formula (see for example Ref. [13]), and the systematic uncertainty associated with the luminosity determination is estimated to be 0.5%.

The cross section for the reaction $e^+e^- \rightarrow K_S^0 K_L^0$ after all corrections is shown as a function of energy in Fig. 15 and listed in Table II. Only statistical uncertainties are shown. The systematic uncertainty is dominated by the background subtraction procedure and is strongly correlated across the entire energy range. It is about 10% at 1.6 GeV, and increases with decreasing cross section to




FIG. 13: The $K_S^0 K_L^0$ invariant mass distribution for data (circles) in the χ^2 control (a) and signal (b) regions (see Fig. 12). The histogram in (a) represents MC-simulated signal events in the control region, and that in (b) represents the simulated background from $\phi\eta$, $K_S^0 K_L^0 \pi^0$, and $K_S^0 K_L^0 2\pi^0$ events; The squares show the total estimated background, obtained as the difference between the data and simulated distributions in (a), normalized to the data in the signal region. (c) The $K_S^0 K_L^0$ invariant mass distribution above 1.06 GeV/ c^2 for the data after background subtraction.

TABLE II: Summary of the $e^+e^- \rightarrow K_S K_L$ cross section measurement. Uncertainties are statistical only.

$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)
1.08	6.86 ± 0.43	1.36	0.40 ± 0.13	1.64	0.83 ± 0.13	1.92	0.07 ± 0.05
1.12	2.86 ± 0.30	1.40	0.22 ± 0.11	1.68	0.51 ± 0.11	1.96	0.09 ± 0.05
1.16	1.78 ± 0.24	1.44	0.32 ± 0.12	1.72	0.26 ± 0.11	2.00	0.02 ± 0.03
1.20	1.48 ± 0.23	1.48	0.36 ± 0.11	1.76	0.11 ± 0.07	2.04	0.02 ± 0.03
1.24	0.87 ± 0.18	1.52	0.66 ± 0.13	1.80	0.03 ± 0.05	2.08	0.00 ± 0.03
1.28	0.54 ± 0.14	1.56	0.67 ± 0.13	1.84	0.04 ± 0.04	2.12	0.01 ± 0.02
1.32	0.54 ± 0.15	1.60	0.84 ± 0.12	1.88	0.02 ± 0.04	2.16	0.02 ± 0.03

~ 30 (50)% for values below 0.5 (0.3) nb. Also shown are all other available data, which are consistent with our results. Below 1.4 GeV, our measurement has precision comparable to the measurements by the CMD2 [23] and SND [25] experiments at VEPP-2M, and is much more precise than the result from the OLYA experiment [26]. In the 1.4-2.4 GeV region, our result is much more precise than the only other available measurement, from the DM1 [24] experiment.

The measured cross section exhibits a distinctive structure around 1.6 GeV, indicating the presence of a vector resonance, perhaps the $\phi(1680)$. Denoting it ϕ' , we fit the cross section using Eq. 4 with the additional amplitude

$$-A_{\phi'} = \frac{\Gamma_{\phi'} m_{\phi'}^3 \sqrt{m_{\phi'} \sigma_{\phi' \rightarrow K_S^0 K_L^0} / C} q_{K_S^0 K_L^0}^{3/2}(m_{\phi'}^2)}{D_{\phi'}(s) q_{K_S^0 K_L^0}^{3/2}(m_{\phi'}^2)}. \quad (10)$$

The energy-dependent width (see Eq. 5) assumes the branching fractions and phase space factors of the major $\phi(1680)$ decay modes, $f = K^* K$, $\phi\eta$, $\phi\pi\pi$, and $K_S^0 K_L^0$ taken from Refs. [11, 15]. We fix the $\phi(1020)$ parameters to the values obtained in Sec. VIB, and float the parameters of the ϕ' . Since the other vector states in this energy range, such as $\omega(1420, 1650)$ and $\rho(1450, 1700)$,

are relatively wide and overlap considerably, we again describe the sum of their contributions using the non-resonant cross section σ_{bkg} and phase Ψ of Eq. 7. First, we fix both to zero, and the fit yields a relatively good description of the data, with $\chi^2 = 30$ for (29-4) degrees of freedom. The result of the fit (solid curve) is compared with the data in Figs. 16 and 17.

Next, we allow σ_{bkg} and Ψ to float in the fit, and obtain $\Psi = 0.2 \pm 0.6$ radians. Since this is consistent with zero, we fix it to zero and repeat the fit. The result is shown as the dashed curves in Figs. 16 and 17. We obtain an improved description of the cross section, with $\chi^2 = 21$ for (29-6) degrees of freedom and the fitted parameter values:

$$\begin{aligned} \sigma_{\phi'} &= 0.46 \pm 0.10 \pm 0.05 \text{ nb}, \\ m_{\phi'} &= 1674 \pm 12 \pm 6 \text{ MeV}/c^2, \\ \Gamma_{\phi'} &= 165 \pm 38 \pm 70 \text{ MeV}, \\ \sigma_{\text{bkg}} &= 0.36 \pm 0.16 \text{ nb}, \end{aligned}$$

where the first uncertainties are statistical and the second systematic, dominated by the difference between fixed and floated Ψ . The relative phase between the non-resonant background and the ϕ resonance is consistent with that between the ϕ' and ϕ resonances, but the un-




FIG. 14: a) The $K_S^0 K_L^0$ mass distributions from the MC simulation for the signal (unshaded) and control (shaded) regions of Fig. 12. b) The mass dependence of the simulated net reconstruction and selection efficiency for the $K_S^0 K_L^0$ final state in the $m(K_S^0 K_L^0) > 1.06 \text{ GeV}/c^2$ region.




FIG. 16: The $e^+e^- \rightarrow K_S^0 K_L^0$ cross section (points) compared with the results of the fits described in the text with the non-resonant amplitude fixed to zero (solid lines) and floating (dashed lines).




FIG. 15: The $e^+e^- \rightarrow K_S^0 K_L^0$ cross section compared with all available data in this $E_{c.m.}$ region.

certainty is very large.

Our parameter values for this resonance are consistent with those of the PDG for the $\phi(1680)$, and with the results of similar fits performed in Refs. [11, 15] for the K^*K , $\phi\eta$, and $\phi\pi\pi$ decay modes of the $\phi(1680)$. However, as shown in Fig. 17, the cross section for $e^+e^- \rightarrow K^+K^-$ is quite different from that for $K_S^0 K_L^0$, indicating substantial interference between the iso-scalar and iso-vector amplitudes in this energy range. The fitting function used above is not able to reproduce the




FIG. 17: Comparison of the $e^+e^- \rightarrow K_S^0 K_L^0$ cross section (points) with that for $e^+e^- \rightarrow K^+K^-$ [8] (crosses).

K^+K^- data [8], and therefore the results should be taken with caution. A simultaneous fit to the cross sections for $e^+e^- \rightarrow \pi^+\pi^-$ (pure iso-vector), $\pi^+\pi^-\pi^0$ (pure iso-scalar), K^+K^- , $K_S^0 K_L^0$, and perhaps other multihadron final states is needed to extract the iso-scalar and iso-vector components, together with reliable resonance parameter values.

The product $\Gamma_{\phi'}\sigma_{\phi'}$ is proportional to the total number of events and does not depend on the experimental

resolution. Using this product as a free parameter in the fit and Eq. 6 we obtain for the $\phi(1680)$ candidate

$$\Gamma_{ee}\mathcal{B}_{K_S^0 K_L^0} = (14.3 \pm 2.4 \pm 1.5 \pm 6.0) \text{ eV} , \quad (11)$$

where the first uncertainty is statistical, the second systematic, and the third due to model dependence. There is no independent measurement of the $\phi(1680) \rightarrow K_S^0 K_L^0$ branching fraction that could be used to calculate Γ_{ee} . However, we have also measured $\Gamma_{ee}\mathcal{B}_{K^* K} = 369 \pm 53 \text{ eV}$, $\Gamma_{ee}\mathcal{B}_{\phi\eta} = 138 \pm 43 \text{ eV}$ [15], and $\Gamma_{ee}\mathcal{B}_{\phi\pi\pi} = 42 \pm 5 \text{ eV}$ [11]. We assume these are the four dominant decay modes, estimate their rates, and use them in our $\Gamma(s)$ calculation.

VIII. THE $K_S^0 K_L^0 \pi^+ \pi^-$ FINAL STATE

We now consider the events with exactly two tracks not assigned to the K_S^0 candidate, but consistent with originating from the same event vertex. This final state has four charged particles, and therefore large backgrounds from ISR and non-ISR multihadron events. We make additional requirements on the two tracks and the rest of the event in order to suppress these backgrounds.

A. Additional selection criteria

The two additional tracks must not be identified as K^\pm , and are required to extrapolate to within $\pm 3 \text{ cm}$ of the collision point in the direction along the beam axis and 0.25 cm in the perpendicular direction. The event must contain no other tracks that extrapolate to within 1 cm of the axis, which is also the lower limit on the radial position of the $K_S^0 \rightarrow \pi^+ \pi^-$ vertex. Considering all pairs of EMC clusters except those assigned to the ISR photon and K_L^0 candidates, we observe a large signal from π^0 , similar to that shown in Fig. 11. As in that case, we require $E_{\gamma, \text{max}} < 0.5 \text{ GeV}$, reducing backgrounds from several sources with a loss of 3% in signal efficiency, as shown in Sec. VII A.

ISR $K_S^0 K^\pm \pi^\mp \pi^0$ events with the charged kaon misidentified as a pion and a cluster from a π^0 photon taken as the K_L^0 candidate are indistinguishable from signal events. To reduce this background, we pair the K_L^0 cluster with all other EMC clusters. For every such pair with $M(\gamma\gamma)$ within $0.03 \text{ GeV}/c^2$ of the π^0 mass, we perform a kinematic fit to the $K_S^0 K^\pm \pi^\mp \pi^0 \gamma$ hypothesis and require $\chi^2(K_S^0 K \pi \pi^0) > 100$.

The 3C χ^2 distribution for the remaining candidate events under the $K_S^0 K_L^0 \pi^+ \pi^- \gamma$ hypothesis is shown as the points in Fig. 18a, with the corresponding MC-simulated pure $K_S^0 K_L^0 \pi^+ \pi^- \gamma$ events shown as the open histogram. The simulated distribution is normalized to the data in the region $\chi^2 < 1$, where the contribution of higher-order ISR is small and the background contamination is lowest, but still amounts to about 15% of the signal. The shaded, cross-hatched, and hatched areas represent the simulated

contributions from the ISR $\phi\eta$, ISR $K_S^0 K^\pm \pi^\mp \pi^0$, and non-ISR $q\bar{q}$ channels, respectively. These backgrounds account for only half of the observed data-MC difference in the distribution at large χ^2 values.

We define a signal region $\chi^2(K_S^0 K_L^0 \pi^+ \pi^-) < 25$ and a control region $25 < \chi^2(K_S^0 K_L^0 \pi^+ \pi^-) < 50$ (vertical lines in Fig. 18), from which we estimate backgrounds in the signal region. The signal region contains 10 788 data and 6825 MC events, while the control region contains 5756 and 633 events, respectively.

B. Background subtraction

The background to the $K_S^0 K_L^0 \pi^+ \pi^-$ mass distribution is subtracted in two stages. The χ^2 distributions for the $K_S^0 K^\pm \pi^\mp \pi^0$ and non-ISR $q\bar{q}$ events peak at low values, since their kinematics are similar to those of signal events. We therefore subtract their MC-simulated contribution from both the signal and control regions of Fig. 18(a). There are large uncertainties in their normalizations, but this has little effect on the total uncertainty. The mass distribution for the data in the signal region before background subtraction is shown in Fig. 18(b) as the points, with the simulated $K_S^0 K^\pm \pi^\mp \pi^0$ and $q\bar{q}$ events shown as the shaded and cross-hatched histograms, respectively.

We estimate the remaining background using the mass distributions for the remaining events in the signal and control regions, according to Eq. 8 of Sec. VII B. The contribution is shown as the hatched area in Fig. 18(b). We fit the sum of all backgrounds with a polynomial function to reduce the statistical fluctuations (curve in Fig. 18(b)) and use this fit for the background subtraction. The resulting mass distribution for $e^+e^- \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$ events is shown in Fig. 18(c). We observe 3320 events in the mass range from threshold to $4.0 \text{ GeV}/c^2$. In addition to a main peak around $2 \text{ GeV}/c^2$, a J/ψ signal and a possible structure just below $3 \text{ GeV}/c^2$ are visible.

We estimate the systematic uncertainty due to the background subtraction to be about 10% for $m(K_S^0 K_L^0 \pi^+ \pi^-) < 2.5 \text{ GeV}/c^2$ (i.e., a 30% uncertainty on a 30% total background), increasing to about 30% in the 2.5-3.0 GeV/c^2 region and reaching 100% above $3.4 \text{ GeV}/c^2$, where background dominates.

C. Simulated detection efficiency

The selection procedures applied to the data are also applied to the MC-simulated event sample. The resulting $K_S^0 K_L^0 \pi^+ \pi^-$ invariant mass distribution is shown in Fig. 19(a) for the signal and control (shaded histogram) regions. The detection efficiency as a function of mass is obtained by dividing the number of reconstructed MC events in each $0.05 \text{ GeV}/c^2$ mass interval by the number generated in that interval, and is shown in Fig. 19(b). The $50 \text{ MeV}/c^2$ mass interval used is wider than the detector resolution of about $25 \text{ MeV}/c^2$. Since the cross




FIG. 18: (a) The three-constraint χ^2 distributions for data (points) and MC simulated $K_S^0 K_L^0 \pi^+ \pi^- \gamma$ events (open histogram). The shaded, cross-hatched, and hatched histograms represent the simulated contributions from ISR $\phi\eta$, ISR $K_S^0 K \pi \pi^0$, and non-ISR $q\bar{q}$ events, respectively. (b) The $K_S^0 K_L^0 \pi^+ \pi^-$ invariant mass distribution for data events in the signal region of (a) (points). The shaded and cross-hatched histograms represent the simulated contributions from ISR $\phi\eta + K_S^0 K \pi \pi^0$ and non-ISR $q\bar{q}$ events, respectively, and the hatched area represents that estimated from the control region. The curve shows the empirical fit used for background subtraction. (c) The $K_S^0 K_L^0 \pi^+ \pi^-$ invariant mass distribution after background subtraction.




FIG. 19: (a) The $K_S^0 K_L^0 \pi^+ \pi^-$ invariant mass distribution for MC-simulated signal events in the signal (open histogram) and control region (shaded) of Fig. 18. (b) The net reconstruction efficiency from the simulation.

section has no sharp structures (except for the J/ψ signal, which is discussed below), we apply no corrections for the resolution. We apply all the corrections discussed above for data-MC differences in the tracking, photon, and K_L^0 detection efficiencies.




FIG. 20: The $e^+e^- \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$ cross section. The error bars are statistical only.

D. The $e^+e^- \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$ cross section

The cross section for the reaction $e^+e^- \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$ is calculated using Eq. 9 with the corrections described above, plus an additional 3% correction for the requirement on the maximum energy of extra EMC clusters. The cross section is shown as a function of energy in Fig. 20, and listed in Table III. There are no previous measurements for this final state. The cross section shows a threshold rise at 1.5 GeV, a maximum value of about 1 nb near 2 GeV, and a slow decrease toward

TABLE III: Summary of the $e^+e^- \rightarrow K_S K_L \pi^+ \pi^-$ cross section measurement. Uncertainties are statistical only.

$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)
1.425	0.03 ± 0.02	2.075	0.99 ± 0.09	2.725	0.11 ± 0.05	3.375	0.07 ± 0.04
1.475	0.12 ± 0.04	2.125	0.81 ± 0.08	2.775	0.24 ± 0.05	3.425	0.01 ± 0.04
1.525	0.15 ± 0.04	2.175	0.69 ± 0.08	2.825	0.21 ± 0.05	3.475	-0.01 ± 0.03
1.575	0.17 ± 0.05	2.225	0.56 ± 0.08	2.875	0.15 ± 0.05	3.525	0.03 ± 0.03
1.625	0.20 ± 0.05	2.275	0.51 ± 0.07	2.925	0.22 ± 0.05	3.575	0.03 ± 0.03
1.675	0.27 ± 0.06	2.325	0.40 ± 0.07	2.975	0.17 ± 0.05	3.625	0.00 ± 0.03
1.725	0.39 ± 0.06	2.375	0.43 ± 0.07	3.025	0.18 ± 0.05	3.675	-0.01 ± 0.03
1.775	0.58 ± 0.07	2.425	0.31 ± 0.06	3.075	0.60 ± 0.06	3.725	-0.06 ± 0.03
1.825	0.60 ± 0.08	2.475	0.27 ± 0.06	3.125	0.44 ± 0.05	3.775	0.02 ± 0.03
1.875	0.83 ± 0.08	2.525	0.22 ± 0.06	3.175	0.16 ± 0.05	3.825	-0.03 ± 0.03
1.925	1.09 ± 0.09	2.575	0.17 ± 0.06	3.225	0.05 ± 0.04	3.875	0.00 ± 0.03
1.975	1.03 ± 0.09	2.625	0.23 ± 0.06	3.275	0.08 ± 0.04	3.925	0.04 ± 0.03
2.025	0.94 ± 0.09	2.675	0.18 ± 0.05	3.325	0.05 ± 0.04	3.975	0.01 ± 0.03

higher energies, perturbed by the J/ψ signal.

Only statistical uncertainties are shown. The total systematic uncertainty is dominated by the background subtraction procedure. It amounts to about 10% at 2 GeV, where the cross section peaks, and increases with decreasing cross section to $\sim 30\%$ near 1.5 and 3 GeV, and to 100% well above 3 GeV.




FIG. 21: The $K_L^0 \pi^+$ invariant mass versus the $K_S^0 \pi^-$ invariant mass (two entries per event).

E. The $K^*(892)^\pm$ and $K_2^*(1430)^\pm$ contributions

Figure 21 shows a scatter plot of the $K_L^0 \pi^+$ invariant mass versus the $K_S^0 \pi^-$ invariant mass, with two entries per event. Clear bands corresponding to the $K^*(892)^\pm$ resonances are visible. Indications of $K_2^*(1430)^\pm$ production are also seen in the projections shown in Fig. 22.




FIG. 22: The (a) $K_S^0 \pi^\pm$ and (b) $K_L^0 \pi^\pm$ mass projections of Fig. 21. The curves represent the results of the fits described in the text, with the hatched areas representing the non-resonant components.

We fit these projections with a sum of two Breit-Wigner functions and a function describing the non-resonant contribution, yielding $3335 \pm 115 K^*(892)^\pm \rightarrow K_S^0 \pi^\pm$ decays, $3200 \pm 151 K^*(892)^\pm \rightarrow K_L^0 \pi^\pm$ decays, and a total of $286 \pm 99 K_2^*(1430)^\pm$ decays. The total number of $K^*(892)^\pm$ decays is larger than the number of $K_S^0 K_L^0 \pi^+ \pi^-$ events, indicating correlated production of $K^*(892)^+ K^*(892)^-$ pairs. In each $0.04 \text{ GeV}/c^2$ bin of $K_L^0 \pi^\mp$ mass we fit the $K_S^0 \pi^\pm$ mass distribution with the same function, and the resulting numbers of $K^*(892)^\pm$ decays are shown in Fig. 23.

A strong signal of $2098 \pm 61 \pm 200 K^*(892)^\pm$ is observed, where the second uncertainty is due to variations of the fitting procedure. This corresponds to the production of $K^*(892)^+ K^*(892)^-$ pairs in about 63% of all observed $K_S^0 K_L^0 \pi^+ \pi^-$ events. We also find $105 \pm 23 \pm 50$ events at the $K_2^*(1430)^\pm$ mass, corresponding to $K^*(892)^\pm K_2^*(1430)^\mp$ correlated production. We have observed such correlated production previously in the $K^+ K^- \pi^0 \pi^0$ channel [11]; these results are compared and discussed below.




FIG. 23: The number of $K^*(892)^\pm$ events obtained from fits to the $K_S^0\pi^\pm$ invariant mass distribution in each $0.04 \text{ GeV}/c^2$ interval of $K_L^0\pi^\mp$ mass. The curve represents the result of the fit described in the text, with the hatched areas representing the non-resonant component.




FIG. 24: (a) The $K_S^0 K_L^0$ invariant mass distribution for the selected $K_S^0 K_L^0 \pi^+ \pi^-$ events. The solid and dashed lines represent the result of the fit described in the text and its non- ϕ component, respectively. (b) The $\pi^+ \pi^-$ invariant mass distribution for events in the ϕ peak (see text).

F. The $\phi(1020)\pi^+\pi^-$ contribution

Figure 24(a) shows the $K_S^0 K_L^0$ invariant mass distribution for the selected $K_S^0 K_L^0 \pi^+ \pi^-$ events. A clear $\phi(1020)$ signal is visible. Fitting with a Gaussian plus polynomial function yields $424 \pm 30 \phi \rightarrow K_S^0 K_L^0$ decays, corresponding to about 13% of the events.

We calculate the $\pi^+ \pi^-$ invariant mass for events in the ϕ region, $1.01 < m(K_S^0 K_L^0) < 1.04 \text{ GeV}/c^2$, and subtract the non-resonant contribution using events in the sideband $1.04 < m(K_S^0 K_L^0) < 1.07 \text{ GeV}/c^2$. We show the resulting $m(\pi^+ \pi^-)$ distribution in Fig. 24(b). It is consistent with those observed in the $\phi\pi^+\pi^-$ and $\phi\pi^0\pi^0$ final states [11], where $f_0(980)$ signals were clearly seen. Fit-

ting the $m(K_S^0 K_L^0)$ distribution in bins of the $K_S^0 K_L^0 \pi^+ \pi^-$ mass, we obtain a $\phi\pi^+\pi^-$ invariant mass spectrum consistent with those observed in the $K^+ K^- \pi^+ \pi^-$ and $K^+ K^- \pi^0 \pi^0$ final states [11]. However, the statistical uncertainties are quite large, and so we do not present the distribution or calculate a cross section for this intermediate state.

IX. THE $K_S^0 K_S^0 \pi^+ \pi^-$ FINAL STATE

A. Final Selection and Backgrounds

This final state contains six charged pions and no neutral particles other than the ISR photon. We consider the events from Sec. III with at least two K_S^0 candidates, and the combination of two K_S^0 candidates and two charged tracks in each event giving the best χ^2 for a 4C fit under the $K_S^0 K_S^0 \pi^+ \pi^-$ hypothesis (see Sec. V). To reduce the background from multihadronic $q\bar{q}$ events, we reject events in which both of the charged tracks not in a K_S^0 candidate are identified as kaons.

The $\chi^2(K_S^0 K_S^0 \pi^+ \pi^-)$ distribution for the selected events in the data is shown in Fig. 25 (points), along with that for selected simulated ISR $K_S^0 K_S^0 \pi^+ \pi^-$ events (open histogram), which is normalized to the data in the region $\chi^2(K_S^0 K_S^0 \pi^+ \pi^-) < 10$ where the backgrounds and radiative corrections do not exceed 5%. Both distributions are broader than those for a typical 4C χ^2 distribution due to higher-order ISR, and the data include contributions from background processes.




FIG. 25: The four-constraint χ^2 distributions for $K_S^0 K_S^0 \pi^+ \pi^- \gamma$ candidate events selected in the data (points) and signal-MC simulation (open histogram) fitted under the $K_S^0 K_S^0 \pi^+ \pi^-$ hypothesis. The cross-hatched histogram represents the simulated background contribution from non-ISR $q\bar{q}$ events.

The cross-hatched histogram in Fig. 25 represents the background from non-ISR $e^+e^- \rightarrow q\bar{q}$ events. These predominantly contain a hard π^0 , giving a false ISR photon, and have kinematics similar to the signal, giving a peak at low values of $\chi^2(K_s^0 K_s^0 \pi^+ \pi^-)$. We evaluate this background in a number of $E_{c.m.}$ ranges using the selected data and $q\bar{q}$ events simulated with JETSET. Combining each ISR photon candidate with all other EMC clusters in the same event, we compare the π^0 signals in the resulting data and simulated $\gamma\gamma$ invariant mass distributions. The simulation gives an $E_{c.m.}$ dependence consistent with the data, so we normalize its prediction using the overall data-over-MC ratio of π^0 signals, and subtract that from the data.

All remaining background sources are either negligible or yield a $\chi^2(K_s^0 K_s^0 \pi^+ \pi^-)$ distribution that is nearly uniform over the range shown in Fig. 25. We define signal and control regions, $\chi^2(K_s^0 K_s^0 \pi^+ \pi^-) < 25$ and $25 < \chi^2(K_s^0 K_s^0 \pi^+ \pi^-) < 50$, respectively (see Fig. 25), and use them to estimate and subtract the sum of the remaining backgrounds as described in Sec. VII B. The signal region of Fig. 25 contains 1704 data and 8309 MC-simulated events; the control region contains 219 data and 580 simulated events.

We recalculate the masses of the two K_s^0 candidates using the results of the kinematic fit. Figure 26 shows a scatter plot of the invariant mass of one K_s^0 candidate versus that of the other for events in the signal region. Any background from events not containing two K_s^0 mesons is very low.




FIG. 26: Scatter plot of the $\pi^+ \pi^-$ invariant mass of one K_s^0 candidate versus that of the other K_s^0 candidate calculated using the results of the constrained fit.

The $m(K_s^0 K_s^0 \pi^+ \pi^-)$ distribution for the events in the signal region of Fig. 25 is shown in Fig. 27 as the points. The contributions from non-ISR events and the background estimated from the control region are shown as




FIG. 27: The $K_s^0 K_s^0 \pi^+ \pi^-$ invariant mass distribution (points) for events in the signal region of Fig. 25. The cross-hatched and hatched histograms represent the backgrounds from non-ISR $q\bar{q}$ events and others estimated from the χ^2 control region of Fig. 25, respectively. The curve represents the smooth empirical fit to the total background used for subtraction.

cross-hatched and hatched histograms, respectively. We fit the sum of all backgrounds with a second order polynomial to reduce fluctuations, and use the result (curve in Fig. 27) for the background subtraction. This gives 1479 signal events with masses between threshold and 4.0 GeV/c^2 . We estimate the systematic uncertainty due to background subtraction to be about 5% of the signal for $m(K_s^0 K_s^0 \pi^+ \pi^-) < 2.5 \text{ GeV}/c^2$, increasing to about 20% in the 2.5-3.0 MeV/c^2 region and 50-70% above 3.0 GeV/c^2 , where background dominates.

B. Simulated detection efficiency

The MC-simulated $K_s^0 K_s^0 \pi^+ \pi^-$ invariant-mass distribution is shown in Fig. 28(a) for events in the signal and control (shaded histogram) regions. The mass dependence of the detection efficiency is shown in Fig. 28(b). The mass interval used, 50 MeV/c^2 per bin, is wider than the 10 MeV/c^2 detector resolution, and the cross section has no sharp structure (except the J/ψ signal, discussed below), so we apply no corrections for the resolution. We apply all the corrections discussed above for data-MC differences in track, K_s^0 , and photon detection efficiency.

C. Cross section for $e^+e^- \rightarrow K_s^0 K_s^0 \pi^+ \pi^-$

We calculate the $e^+e^- \rightarrow K_s^0 K_s^0 \pi^+ \pi^-$ cross section as a function of the effective c.m. energy using Eq. 9 shown in




FIG. 28: (a) The $K_S^0 K_S^0 \pi^+ \pi^-$ invariant mass distribution for the MC-simulated signal events in the signal and control (shaded) regions of Fig. 25. (b) The net reconstruction and selection efficiency from the simulation.

Sec. VIID. The fully corrected cross section is shown in Fig. 29 and listed in Table IV, with statistical uncertainties only. There are no other measurements for this final state. The cross section shows a slow rise from threshold at 1.5 GeV, a maximum value of about 0.5 nb near 2 GeV, and a slow decrease with increasing energy, punctuated by a clear J/ψ signal. The systematic uncertainty is dominated by the uncertainty of the backgrounds, and totals 5% relative at the peak of the cross section, increasing to 20% at 3 GeV, and 50-70% at higher energies.




FIG. 29: The $e^+ e^- \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$ cross section.




FIG. 30: The $K_S^0 \pi^-$ invariant mass versus the $K_S^0 \pi^+$ invariant mass (two entries per event).




FIG. 31: The (a) $m(K_S^0 \pi^+)$ and (b) $m(K_S^0 \pi^-)$ projections of Fig. 30. The lines and hatched areas represent the results of the fits described in the text and their non- K^* components, respectively.


D. The $K^*(892)^\pm$ and $K_2^*(1430)^\pm$ contributions

Figure 30 shows a scatter plot of the $K_S^0 \pi^-$ invariant mass versus the $K_S^0 \pi^+$ invariant mass, with two entries per event. Clear bands associated with the $K^*(892)^\pm$ are visible here, as are peaks in the projections shown in Fig. 31. The projections also show indications of $K_2^*(1430)^\pm$ production.

Fitting the projections with a sum of two Breit-Wigner functions and a threshold function yields 827 ± 29 $K^*(892)^+ \rightarrow K_S^0 \pi^+$ and 856 ± 50 $K^*(892)^- \rightarrow K_S^0 \pi^-$ decays, as well as 116 ± 40 $K_2^*(1430)^+$ and 70 ± 34 $K_2^*(1430)^-$ decays. The total number of $K^*(892)^\pm$ decays is larger than the number of $K_S^0 K_S^0 \pi^+ \pi^-$ events, indicating correlated production of $K^*(892)^+ K^*(892)^-$ pairs. We fit the $K_S^0 \pi^+$ invariant mass distribution in $0.04 \text{ GeV}/c^2$ bins of the $K_S^0 \pi^-$ mass, and show the number of $K^*(892)^+$ decays in each bin in Fig. 32. A clear $K^*(892)^+$ signal is observed; a fit yields $742 \pm 30 \pm 100$


TABLE IV: Summary of the $e^+e^- \rightarrow K_S K_S \pi^+ \pi^-$ cross section measurement. Uncertainties are statistical only.

$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)
1.63	0.02 ± 0.01	2.22	0.22 ± 0.03	2.83	0.04 ± 0.01	3.42	0.01 ± 0.01
1.67	0.03 ± 0.01	2.28	0.25 ± 0.03	2.88	0.04 ± 0.01	3.47	0.00 ± 0.01
1.73	0.05 ± 0.02	2.33	0.13 ± 0.02	2.92	0.05 ± 0.01	3.53	0.04 ± 0.01
1.77	0.13 ± 0.03	2.38	0.12 ± 0.02	2.97	0.07 ± 0.02	3.58	0.02 ± 0.01
1.83	0.22 ± 0.03	2.42	0.10 ± 0.02	3.03	0.05 ± 0.01	3.63	0.02 ± 0.01
1.88	0.24 ± 0.03	2.47	0.12 ± 0.02	3.08	0.28 ± 0.03	3.67	0.04 ± 0.01
1.92	0.27 ± 0.04	2.53	0.11 ± 0.02	3.13	0.19 ± 0.02	3.72	0.01 ± 0.01
1.98	0.38 ± 0.04	2.58	0.12 ± 0.02	3.17	0.04 ± 0.01	3.78	0.02 ± 0.01
2.03	0.47 ± 0.05	2.63	0.07 ± 0.02	3.22	0.07 ± 0.01	3.83	0.02 ± 0.01
2.08	0.37 ± 0.04	2.67	0.08 ± 0.02	3.28	0.05 ± 0.01	3.88	0.00 ± 0.01
2.13	0.38 ± 0.04	2.72	0.08 ± 0.02	3.33	0.03 ± 0.01	3.92	0.01 ± 0.01
2.17	0.25 ± 0.03	2.78	0.05 ± 0.01	3.38	0.01 ± 0.01	3.97	0.02 ± 0.01

FIG. 32: The fitted number of $K^*(892)^+$ events in each $0.04 \text{ GeV}/c^2$ interval of the $K_S^0 \pi^-$ mass. The curve represents the result of the fit described in the text, with the hatched area representing the non-resonant component.

pair production events, $e^+e^- \rightarrow K^*(892)^+ K^*(892)^- \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$, where the second uncertainty is due to variation of the starting values of the fit parameters. This accounts for 50% of the selected events and 88% of the $K^*(892)^\pm$ production. We find no significant signal at the $K_2^*(1430)^+$ mass, and hence no evidence for $e^+e^- \rightarrow K^*(892)^\pm K_2^*(1430)^\mp$ events.

The number of correlated $K^*(892)^+ K^*(892)^-$ production events in this channel (742 ± 104 events with 4.5% efficiency) can be compared with the corresponding numbers in the $K_S^0 K_L^0 \pi^+ \pi^-$ channel (2098 ± 209 events with 5% efficiency), presented above, and in the $K^+ K^- \pi^0 \pi^0$ final state (1750 ± 60 events with 8% efficiency), from our previous measurement [11] using the same integrated luminosity. Normalizing these to the same 5% efficiency, we obtain the ratios $(824 \pm 116):(2098 \pm 209):(1094 \pm 38)$.

FIG. 33: The (a) $\pi^+ \pi^-$ and (b) $K_S^0 K_S^0$ invariant mass distributions for the selected $K_S^0 K_S^0 \pi^+ \pi^-$ events with $K^*(892)^+ K^*(892)^-$ events excluded (see text).

These are consistent with the 1:2:1 ratios expected assuming equal production of K_S^0 and K_L^0 in $K^*(892)^\pm$ decays.

The size of the data sample is not large enough to apply this procedure to every $m(K_S^0 K_S^0 \pi^+ \pi^-)$ bin and extract the $e^+e^- \rightarrow K^*(892)^+ K^*(892)^-$ cross section. However, considering events with both $m(K_S^0 \pi^+)$ and $m(K_S^0 \pi^-)$ within $\pm 0.15 \text{ GeV}/c^2$ of the nominal $K^*(892)^\pm$ mass [30], we conclude that the $K^*(892)^+ K^*(892)^-$ contribution almost completely dominates for $m(K_S^0 K_S^0 \pi^+ \pi^-)$ below 2.5 GeV . For the events outside this box, we show the $\pi^+ \pi^-$ and $K_S^0 K_S^0$ invariant mass distributions in Fig. 33. The $\rho(770)$ resonance is prominent in the $\pi^+ \pi^-$ spectrum, whereas the $K_S^0 K_S^0$ spectrum shows no significant structure. The three resonant channels $K^*(892)^+ K^*(892)^-$, $K^*(892)^\pm K_S^0 \pi^\mp$ (see Fig. 30), and $\rho(770) K_S^0 K_S^0$ dominate the $K_S^0 K_S^0 \pi^+ \pi^-$ cross section within our measured range, and there is a small contribution from $K^*(1430)^\pm K_S^0 \pi^\mp$.

X. THE $K_s^0 K_s^0 K^+ K^-$ FINAL STATE

A. Final selection and background

We consider the events from Sec. III with at least two K_s^0 candidates, and the combination of two K_s^0 candidates and two charged tracks in each event giving the best χ^2 for a 4C fit under the $K_s^0 K_s^0 K^+ K^-$ hypothesis (see Sec. V). To reduce the background from multi-pionic events, we require that both of the charged tracks not in the K_s^0 candidates be identified as kaons.




FIG. 34: The four-constraint χ^2 distributions for $K_s^0 K_s^0 K^+ K^- \gamma$ candidate events in the data (points) and signal MC simulation (open histogram) fitted under the $K_s^0 K_s^0 K^+ K^-$ hypothesis. The cross-hatched histogram represents the simulated contribution from non-ISR $q\bar{q}$ events.

The $\chi^2(K_s^0 K_s^0 K^+ K^-)$ distribution for the selected events is shown in Fig. 34 (points) along with that for simulated ISR $K_s^0 K_s^0 K^+ K^-$ events (open histogram), where the latter distribution is normalized to the data in the region $\chi^2(K_s^0 K_s^0 \pi^+ \pi^-) < 8$. There is very little background: simulated ISR events in other channels do not satisfy the selection; there is no significant π^0 peak in the data; and the signal MC describes the data well, even at high χ^2 values. The simulation predicts only a few $e^+e^- \rightarrow q\bar{q} \rightarrow K_s^0 K_s^0 K^+ K^- \pi^0$ events, which are shown as the hatched histogram in Fig. 34.

We select events with $\chi^2(K_s^0 K_s^0 K^+ K^-) < 40$, obtaining 129 events in the data with masses between threshold and 4.5 GeV/c^2 , and 2544 events in the signal MC simulation. The $K_s^0 K_s^0 K^+ K^-$ invariant mass distribution is shown as the open histogram in Fig. 35. We do not subtract any background, nor do we assign any systematic uncertainty to account for possible background contribu-




FIG. 35: The $K_s^0 K_s^0 K^+ K^-$ invariant mass distribution for data events in the signal region, $\chi^2(K_s^0 K_s^0 K^+ K^-) < 40$ (open histogram). The subset of events with $m(K^+ K^-) < 1.04 \text{ GeV}/c^2$, predominantly $K_s^0 K_s^0 \phi(1020)$ events, is shown as the shaded histogram.




FIG. 36: (a) The $K_s^0 K_s^0 K^+ K^-$ invariant mass distribution for the MC-simulated signal events in the signal and control (shaded) regions of Fig. 34. (b) The net reconstruction and selection efficiency from the simulation.

B. Simulated detection efficiency

The MC-simulated $K_s^0 K_s^0 K^+ K^-$ invariant-mass distribution is shown in Fig. 36(a) for events in the signal and control (shaded histogram) regions. The mass dependence of the detection efficiency is shown in

TABLE V: Summary of the $e^+e^- \rightarrow K_S^0 K_S^0 K^+ K^-$ cross section measurement. Uncertainties are statistical only.

$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)	$E_{c.m.}$ (GeV)	σ (nb)
2.05	0.003 ± 0.003	2.75	0.010 ± 0.004	3.45	0.013 ± 0.004	4.15	0.003 ± 0.002
2.15	0.000 ± 0.003	2.85	0.010 ± 0.004	3.55	0.006 ± 0.003	4.25	0.000 ± 0.003
2.25	0.008 ± 0.004	2.95	0.011 ± 0.005	3.65	0.004 ± 0.002	4.35	0.001 ± 0.001
2.35	0.010 ± 0.005	3.05	0.012 ± 0.005	3.75	0.005 ± 0.003	4.45	0.003 ± 0.002
2.45	0.007 ± 0.004	3.15	0.005 ± 0.003	3.85	0.009 ± 0.003		
2.55	0.007 ± 0.004	3.25	0.010 ± 0.004	3.95	0.002 ± 0.002		
2.65	0.017 ± 0.006	3.35	0.009 ± 0.004	4.05	0.004 ± 0.002		

Fig. 36(b). The mass interval used, $50 \text{ MeV}/c^2$ per bin, is wider than the $10 \text{ MeV}/c^2$ detector resolution, and the cross section has no sharp structure (except the J/ψ signal, discussed below), so we apply no corrections for the resolution. We apply all the corrections discussed above for data-MC differences in track, K_S^0 , and photon detection efficiency.




FIG. 37: The $e^+e^- \rightarrow K_S^0 K_S^0 K^+ K^-$ cross section. Events with invariant mass within $0.05 \text{ GeV}/c^2$ of the J/ψ mass are excluded.

C. Cross section for $e^+e^- \rightarrow K_S^0 K_S^0 K^+ K^-$

We remove the events within $\pm 0.05 \text{ GeV}/c^2$ of the J/ψ signal (which is discussed below), and calculate the $e^+e^- \rightarrow K_S^0 K_S^0 K^+ K^-$ cross section using Eq. 9. The fully corrected cross section is shown in as a function of energy in Fig. 37 and listed in Table V. There are no previous measurements of this final state. The systematic uncertainties are smaller than the statistical terms and do not exceed 5%.

D. Internal structure of the $K_S^0 K_S^0 K^+ K^-$ system

Figure 38(a) shows a scatter plot of the $K^+ K^-$ invariant mass versus the $K_S^0 K_S^0$ invariant mass for all selected events. A strong $\phi(1020)$ band is evident. Requiring $m(K^+ K^-) < 1.04 \text{ GeV}/c^2$, we obtain the contribution from $\phi K_S^0 K_S^0$ events shown in Fig 35 as the shaded histogram. This mode dominates at all masses.

There is also structure for $m(K_S^0 K_S^0)$ near $1.5 \text{ GeV}/c^2$, which is more visible as a peak in the $m(K_S^0 K_S^0)$ projection of Fig. 38(b). We fit this mass region with a Breit-Wigner plus a second-order polynomial function. An expanded view is shown in Fig. 38(c), along with the result of the fit. We obtain 29 ± 7 events with Breit-Wigner mass and width

$$m = 1.526 \pm 0.007 \text{ GeV}/c^2$$

$$\Gamma = 0.037 \pm 0.012 \text{ GeV}.$$

These parameters may be compared with the averages [30] for the $f_2'(1525)$ resonance, $m(f_2') = 1.525 \pm 0.005 \text{ GeV}/c^2$ and $\Gamma(f_2') = 0.073_{-0.005}^{+0.006} \text{ GeV}$; the mass is consistent but the width is about 3 standard deviations lower.

XI. THE CHARMONIUM REGION

Figures 39(a), (b), and (c) show expanded views of the mass distributions in Figs. 18(c), 27, and 35, respectively, in the J/ψ mass region. Fitting with Gaussian plus second-order polynomial functions yields 248 ± 27 $J/\psi \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$ decays, 133 ± 13 $J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$ decays, and 28.5 ± 5.5 $J/\psi \rightarrow K_S^0 K_S^0 K^+ K^-$ decays. Using the respective simulated efficiencies with all the corrections described above, and the differential luminosity, we calculate the products of the J/ψ electronic width and branching fractions to these modes, and list them in Table VI. Using the PDG value of $\Gamma_{ee}(J/\psi) = 5.55 \text{ keV}$ [30], we obtain the corresponding branching fractions, also presented in Table VI. These are the first observations of these J/ψ decay modes and measurements of their branching fractions. They can be compared with $\mathcal{B}(J/\psi \rightarrow K^+ K^- \pi^+ \pi^-) = (6.8 \pm 0.3) \times 10^{-3}$ [30], which is dominated by the *BABAR* measurement.




FIG. 38: (a) The K^+K^- versus $K_S^0K_S^0$ invariant mass for all selected $K_S^0K_S^0K^+K^-$ events in the data. (b) The $m(K_S^0K_S^0)$ projection of (a). (c) An expanded view of (b) in which the line represents the result of the fit described in the text.




FIG. 39: Expanded views of the invariant mass distributions near the J/ψ mass for the (a) $K_S^0K_S^0\pi^+\pi^-$, (b) $K_S^0K_S^0\pi^+\pi^-$, and (c) $K_S^0K_S^0K^+K^-$ final states. The lines represent the results of the fits described in the text.

A. Internal structure of the $J/\psi \rightarrow K_S^0K_S^0\pi^+\pi^-$ and $K_S^0K_S^0K^+K^-$ decays

The J/ψ signal in the $K_S^0K_L^0\pi^+\pi^-$ mode has a large non-resonant background (see Fig. 39(a)), and we are unable to quantify the contributions from the $K^*(892)K_S^0\pi$ and $\phi\pi^+\pi^-$ intermediate states with reasonable accuracy. The $J/\psi \rightarrow \phi\pi^+\pi^-$ decay rate is relatively well measured [30], dominated by *BABAR*.

The $K_S^0K_S^0\pi^+\pi^-$ channel has much lower background (see Fig. 39(b)), and we use the 157 events with invariant mass within 30 MeV/ c^2 of the nominal J/ψ mass to study intermediate states. We use events in the 30 MeV/ c^2 intervals on each side of the signal region to estimate a non- J/ψ contribution of 24 events, and to subtract the corresponding contributions from the histograms that follow. The resulting $m(K_S^0K_S^0)$ distribution (four entries per event) is shown in Fig. 40(a). Fitting with two Breit-Wigner (BW) functions plus a polynomial, we obtain 53 ± 14 events containing $K^*(892)K_S^0\pi$ and 35 ± 15 con-

taining $K_2^*(1430)K_S^0\pi$. To estimate decays to correlated $K^*(892)^+K^*(892)^-$ or $K_2^*(1430)^\mp K^*(892)^\pm$ pairs, we consider events from the $K^*(892)^+$ and $K^*(892)^-$ bands (see Fig. 30) defined by $|m(K_S^0\pi) - 0.892| < 0.15$ GeV/ c^2 ; a pairing in the overlap region gives only one entry, and there can be as many as two entries per event. Fitting the invariant mass distribution of the other $K_S^0\pi$ pair, shown in Fig. 40(b), with two BW functions plus a polynomial, we obtain 0.7 ± 5.0 and 8 ± 8 events for the $K^*(892)^+K^*(892)^-$ and $K_2^*(1430)^\mp K^*(892)^\pm$ combinations, respectively. Both are consistent with zero, i.e., no correlated production.

For each of these intermediate states we calculate the product of its J/ψ branching fraction, $\Gamma_{ee}^{J/\psi}$, and the relevant branching fractions for the intermediate resonances, and list the values in Table VI. Using $\Gamma_{ee}^{J/\psi} = 5.55$ eV, known branching fractions [30], and the assumptions that K^* mesons decay equally to charged and neutral kaons, and equally to K_S^0 and K_L^0 (e.g., $\mathcal{B}_{K_2^*(1430) \rightarrow K_S^0\pi} = 0.125$), we calculate the corresponding branching fractions, also

TABLE VI: Summary of the J/ψ parameters obtained in this analysis.

Measured Quantity	Measured Value (eV)	J/ψ Branching Fraction (10^{-3})	
		This work	PDG2012
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow K_S^0 K_L^0 \pi^+ \pi^-}$	$20.8 \pm 2.3 \pm 2.1$	$3.7 \pm 0.6 \pm 0.4$	no entry
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-}$	$9.3 \pm 0.9 \pm 0.5$	$1.68 \pm 0.16 \pm 0.08$	no entry
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow K_S^0 K_S^0 K^+ K^-}$	$2.3 \pm 0.4 \pm 0.1$	$0.42 \pm 0.08 \pm 0.02$	no entry
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow K^*(892) K_S^0 \pi} \cdot \mathcal{B}_{K^*(892) \rightarrow K_S^0 \pi}$	$3.7 \pm 1.2 \pm 0.3$	$2.6 \pm 0.9 \pm 0.2$	no entry
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow K_2^*(1430) K_S^0 \pi} \cdot \mathcal{B}_{K_2^*(1430) \rightarrow K_S^0 \pi}$	$2.5 \pm 1.2 \pm 0.2$	$3.6 \pm 1.7 \pm 0.3$	no entry
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow K^*(892)^+ K^*(892)^-} \cdot \mathcal{B}_{K^*(892) \rightarrow K_S^0 \pi}^2$	$0.05 \pm 0.03 \pm 0.02$	< 1.7 90% C.L.	$1.0_{-0.4}^{+0.2}$
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow K_2^*(1430) K^*(892)} \cdot \mathcal{B}_{K_2^*(1430) \rightarrow K_S^0 \pi} \cdot \mathcal{B}_{K^*(892) \rightarrow K_S^0 \pi}$	$0.58 \pm 0.50 \pm 0.02$	< 7.8 90% C.L.	no entry
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow K_S^0 K_S^0 \phi(1020)} \cdot \mathcal{B}_{\phi \rightarrow K^+ K^-}$	$1.6 \pm 0.4 \pm 0.1$	$0.58 \pm 0.14 \pm 0.03$	no entry
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow f_2'(1525) \phi(1020)} \cdot \mathcal{B}_{\phi \rightarrow K^+ K^-} \cdot \mathcal{B}_{f_2'(1525) \rightarrow K_S^0 K_S^0}$	$0.88 \pm 0.34 \pm 0.04$	$0.45 \pm 0.17 \pm 0.02$	0.8 ± 0.4 (S=2.7)
$\Gamma_{ee}^{J/\psi} \cdot \mathcal{B}_{J/\psi \rightarrow f_2'(1525) K^+ K^-} \cdot \mathcal{B}_{f_2'(1525) \rightarrow K_S^0 K_S^0}$	$1.28 \pm 0.42 \pm 0.05$	$0.32 \pm 0.11 \pm 0.02$	no entry




FIG. 40: (a) The $K_S^0 K_S^0 \pi^+ \pi^-$ invariant mass distribution (four entries per event) for the $K_S^0 K_S^0 \pi^+ \pi^-$ events under the J/ψ peak with the non- J/ψ contribution subtracted (see text). (b) The distribution of the other $m(K_S^0 \pi^+)$ for those events with one $m(K_S^0 \pi^+)$ value within $0.15 \text{ GeV}/c^2$ of the $K^*(892)^\pm$ mass (up to two entries per event, pairings in the overlap region taken once). The lines represent the results of the fits described in the text, with the shaded areas representing the combinatorial components.




FIG. 41: (a) The $m(\pi^+ \pi^-)$ distribution for the $J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$ events. (b) The $m(K_S^0 K_S^0)$ distribution for those events in the $\rho(770)$ region, $0.6 < m(\pi^+ \pi^-) < 1.0 \text{ GeV}/c^2$.

listed in Table VI. The only entry in the PDG tables for any of these channels is $\mathcal{B}_{J/\psi \rightarrow K^*(892)^+ K^*(892)^-} = (1.0_{-0.4}^{+0.2}) \times 10^{-3}$.

Figure 41(a) shows the $\pi^+ \pi^-$ invariant mass distribution for the considered events. A clear signal from

the $\rho(770)$ resonance is seen, corresponding to $J/\psi \rightarrow \rho K_S^0 K_S^0$ decays. The $K_S^0 K_S^0$ invariant mass distribution for those events with $0.6 < m(\pi^+ \pi^-) < 1.0 \text{ GeV}/c^2$, shown in Fig. 41(b), features a narrow spike containing 9.4 ± 4.6 events near $1.53 \text{ GeV}/c^2$. We observe this same signal when no requirement is placed on the $\pi^+ \pi^-$ invariant mass. Attributing this entirely to $J/\psi \rightarrow \rho(770) f_2'(1525)$ decays, we calculate the measured product and branching fraction, using $\mathcal{B}(f_2'(1525) \rightarrow K \bar{K}) = 0.71$ [30], and list them in Table VI. This channel also has no listing in the PDG tables [30]. Due to uncertainties in the mass distributions for events without a ρ or f_2' meson, however, we do not attempt to quantify the more inclusive $\rho K_S^0 K_S^0$ or $\pi^+ \pi^- f_2'$ contributions.




FIG. 42: (a) The $K^+ K^-$ versus $K_S^0 K_S^0$ invariant mass for the $K_S^0 K_S^0 K^+ K^-$ events under the J/ψ peak (see text). (b) The $K_S^0 K_S^0$ invariant mass distribution for the events in (a) with $m(K^+ K^-) < 1.04 \text{ GeV}/c^2$. The line represents the result of the fit described in the text.

Figure 42(a) shows the $K^+ K^-$ versus $K_S^0 K_S^0$ invariant mass for the 30 $K_S^0 K_S^0 K^+ K^-$ events with total invariant mass within $30 \text{ MeV}/c^2$ of the nominal J/ψ mass, 29 \pm 6 of which are J/ψ events. Horizontal and vertical bands are visible, corresponding to the $\phi(1020)$ and $f_2'(1525)$ resonances, respectively. We select 20 $J/\psi \rightarrow \phi(1020) K_S^0 K_S^0$ candidate decays by requiring $m(K^+ K^-) < 1.04 \text{ GeV}/c^2$, and plot their $m(K_S^0 K_S^0)$ distribution in Fig. 42(b). Fitting with a

Breit-Wigner plus a constant function, we obtain 11 ± 4 $J/\psi \rightarrow f_2'(1525)\phi(1020)$ decays; including the five events with $m(K_s^0 K_s^0)$ near 1525 MeV/ c^2 but higher $m(K^+ K^-)$ values (see Fig. 42(b)) gives 16 ± 5 $J/\psi \rightarrow f_2'(1525)K^+ K^-$ decays.

Using these numbers we calculate the products of $\Gamma_{ee}^{J/\psi}$ and the relevant branching fractions, and list them in Table VI. Using the PDG values of $\Gamma_{ee}^{J/\psi}$, $\mathcal{B}(\phi \rightarrow K^+ K^-) = 0.49$, and $\mathcal{B}(f_2'(1525) \rightarrow K\bar{K}) = 0.71$ [30], we obtain the corresponding branching fractions, also shown in Table VI. Only one value can be compared with an existing PDG listing [30], namely $\mathcal{B}(J/\psi \rightarrow f_2'(1525)\phi(1020)) = (8 \pm 4) \times 10^{-4}$, which has a scale factor of 2.7. Our result can be compared to the MarkII value $(4.8 \pm 1.8) \times 10^{-4}$, and to the DM2 measurement $(12.3 \pm 0.26 \pm 2.0) \times 10^{-4}$ [30].

XII. SUMMARY

We have presented a study of the processes $e^+e^- \rightarrow K_s^0 K_L^0$ and $e^+e^- \rightarrow K_s^0 K_L^0 \pi^+ \pi^-$ at low center-of-mass energies using using events with initial-state radiation (ISR) collected with the *BABAR* detector. From the dominant $e^+e^- \rightarrow \phi\gamma \rightarrow K_s^0 K_L^0 \gamma$ process near $K_s^0 K_L^0$ threshold, we measure the probability of detecting the K_L^0 via its nuclear interaction in the electromagnetic calorimeter with about 0.6% uncertainty, as well as its angular resolution. Using the positions of candidate K_L^0 clusters in the calorimeter as input to kinematic fits, we obtain clean samples of $K_s^0 K_L^0 \gamma$ and $K_s^0 K_L^0 \pi^+ \pi^- \gamma$ events, and extract the $e^+e^- \rightarrow K_s^0 K_L^0$ and $e^+e^- \rightarrow K_s^0 K_L^0 \pi^+ \pi^-$ cross sections from threshold to 2.2 and 4 GeV, respectively.

For the $K_s^0 K_L^0$ final state, we perform fits to the $\phi(1020)$ and $\phi(1680)$ resonances, and report the resonance parameters and $\Gamma_{ee} \cdot \mathcal{B}(K_s^0 K_L^0)$ values. The results are consistent with previous measurements, and much more precise for c.m. energies above 1.2 GeV, especially for the $\phi(1680)$ mass region. The $e^+e^- \rightarrow K_s^0 K_L^0 \pi^+ \pi^-$ cross section is measured for the first time, and is dominated by the $K^*(892)^+ K^*(892)^-$ intermediate state. Additional contributions from the $K^*(892)^\pm K_2^*(1430)^\mp$ and $\phi\pi^+\pi^-$ intermediate states are observed.

We also obtain the first measurements of the $e^+e^- \rightarrow K_s^0 K_s^0 \pi^+ \pi^-$ and $e^+e^- \rightarrow K_s^0 K_s^0 K^+ K^-$ cross sections, and provide results from threshold to 4 and 4.5 GeV, respectively. For the former process, we again find the $K^*(892)^+ K^*(892)^-$ intermediate state to be dominant, and measure a contribution from $\rho(770)K_s^0 K_s^0$. However, no significant contribution from $K^*(892)^\pm K_2^*(1430)^\mp$ is observed. For the latter process, we observe contributions from the $K_s^0 K_s^0 \phi(1020)$ and $f_2'(1525)\phi(1020)$ intermediate states.

We observe the $J/\psi \rightarrow K_s^0 K_L^0 \pi^+ \pi^-$, $K_s^0 K_s^0 \pi^+ \pi^-$, and $K_s^0 K_s^0 K^+ K^-$ decays for the first time, and measure the product of the J/ψ electronic width and branching fraction to each of these modes. We

study the substructure of these decays, and obtain the first measurements of the $J/\psi \rightarrow K^*(892)^\pm K_s^0 \pi^\pm$, $K_2^*(1430)^\pm K_s^0 \pi^\pm$, $\rho(770)f_2'(1525)$, $\phi(1020)K_s^0 K_s^0$, and $f_2'(1525)K^+ K^-$ branching fractions. In addition, we measure the $J/\psi \rightarrow f_2'(1525)\phi(1020)$ branching fraction with improved precision, and observe the $\rho(770)K_s^0 K_s^0$ and $f_2'(1525)\pi^+ \pi^-$ decay modes. We do not observe $K^*(892)^+ K^*(892)^-$ or $K_2^*(1430)^\pm K^*(890)^\mp$ decays and set limits on their contributions.

Acknowledgements

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), the Commissariat à l'Énergie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Education and Science of the Russian Federation, Ministerio de Economía y Competitividad (Spain), the Science and Technology Facilities Council (United Kingdom), and the Binational Science Foundation (U.S.-Israel). Individuals have received support from the Marie-Curie IEF program (European Union) and the A. P. Sloan Foundation (USA).

-
- [1] V. N. Baier and V. S. Fadin, Phys. Lett. B **27**, 223 (1968).
- [2] A. B. Arbuzov *et al.*, J. High Energy Phys. **9812**, 009 (1998).
- [3] S. Binner, J.H. Kühn and K. Melnikov, Phys. Lett. B **459**, 279 (1999).
- [4] M. Benayoun *et al.*, Mod. Phys. Lett. A **14**, 2605 (1999).
- [5] M. Davier, S. Eidelman, A. Höcker; Z. Zhang, Eur. Phys. J. C **31**, 503 (2003); M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, Eur. Phys. J. C **71**, 1515 (2011); K. Hagiwara *et al.*, J. Phys. G **38**, 085003 (2011).
- [6] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **69**, 011103 (2004).
- [7] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. **103**, 231801 (2009); J. P. Lees *et al.* (BABAR Collaboration), Phys. Rev. D **86**, 032013 (2012).
- [8] J. P. Lees *et al.* (BABAR Collaboration), Phys. Rev. D **88**, 032013 (2013).
- [9] J.P. Lees *et al.* (BABAR Collaboration), Phys. Rev. D **88**, 072009 (2013).
- [10] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **71**, 052001 (2005).
- [11] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **76**, 012008 (2007).
- [12] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **73**, 052003 (2006).
- [13] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **70**, 072004 (2004).
- [14] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **76**, 092005 (2007).
- [15] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **77**, 092002 (2008).
- [16] J. P. Lees *et al.* (BABAR Collaboration), Nucl. Instrum. and Meth. A **726**, 203 (2013).
- [17] B. Aubert *et al.* (BABAR Collaboration), Nucl. Instrum. and Meth. A **479**, 1 (2002); B. Aubert *et al.* (BABAR Collaboration), Nucl. Instrum. and Meth. A **729**, 615 (2013).
- [18] H. Czyż and J. H. Kühn, Eur. Phys. J. C **18**, 497 (2001).
- [19] A. B. Arbuzov *et al.*, J. High Energy Phys. **9710**, 001 (1997).
- [20] M. Caffo, H. Czyż, E. Remiddi, Nuovo Cim. A **110**, 515 (1997); Phys. Lett. B **327**, 369 (1994).
- [21] E. Barberio, B. van Eijk and Z. Was, Comput. Phys. Commun. **66**, 115 (1991).
- [22] R.R. Akhmetshin *et al.* (CMD-2 Collaboration), Phys. Lett. B **466**, 385 (1999), Phys. Lett. B **508**, 217 (2001).
- [23] R.R. Akhmetshin *et al.* (CMD-2 Collaboration), Phys. Lett. B **551**, 27 (2003).
- [24] F. Mane *et al.*, Phys. Lett. B **99**, 261 (1981).
- [25] M. N. Achasov *et al.* (SND Collaboration), e-Print: hep-ex/9809013.
- [26] P. Ivanov *et al.*, JETP Lett. **36**, 112 (1982).
- [27] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [28] T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994).
- [29] S. Jadach and Z. Was, Comput. Phys. Commun. **85**, 453 (1995).
- [30] C. J. Beringer *et al.* (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).
- [31] M. N. Achasov *et al.* (SND Collaboration), e-Print: hep-ex/0009036v2 (2001).
- [32] R.R. Akhmetshin *et al.* (CMD-2 Collaboration), Phys. Lett. B **695**, 412 (2011).

