

# Search for Scalar Leptoquarks from $Z^0$ decays

DELPHI Collaboration

## Abstract

We have searched for pair-produced scalar leptoquarks each decaying to a quark and a charged lepton in a sample of 116 000 hadronic  $Z^0$  events produced at LEP. No candidate was detected and cross section and branching ratio limits are set for the above process at 95% CL. Mass limits are found to be about  $42 \text{ GeV}/c^2$  depending only slightly on the models used and a coupling times branching ratio exclusion line is drawn for a scalar leptoquark with a free coupling. We have also probed the mass region above  $45 \text{ GeV}/c^2$  for a singly-produced scalar leptoquark and set limits on the cross section and the coupling  $\lambda^2/4\pi$  up to 60 GeV.

(Submitted to Physics Letters B)

P.Abreu<sup>17</sup>, W.Adam<sup>43</sup>, F.Adami<sup>34</sup>, T.Adye<sup>32</sup>, T.Akesson<sup>20</sup>, G.D.Alekseev<sup>13</sup>, P.Allen<sup>42</sup>, S.Almhed<sup>20</sup>,  
 S.J.Alvsvaag<sup>4</sup>, U.Arnaldi<sup>7</sup>, E.Anassontzis<sup>3</sup>, P.Antilogus<sup>21</sup>, W-D.Apel<sup>14</sup>, R.J.Apsimon<sup>32</sup>, B.Åsman<sup>38</sup>, P.Astier<sup>19</sup>,  
 J-E.Augustin<sup>16</sup>, A.Augustinus<sup>7</sup>, P.Baillon<sup>7</sup>, P.Bambade<sup>16</sup>, F.Barao<sup>17</sup>, R.Barate<sup>11</sup>, G.Barbiellini<sup>40</sup>,  
 D.Y.Bardin<sup>13</sup>, A.Baroncelli<sup>35</sup>, O.Barring<sup>20</sup>, W.Bartl<sup>43</sup>, M.J.Bates<sup>30</sup>, M.Battaglia<sup>25</sup>, M.Baubillier<sup>19</sup>,  
 K-H.Becks<sup>45</sup>, C.J.Beeston<sup>30</sup>, M.Begalli<sup>10</sup>, P.Beilliere<sup>6</sup>, Yu.Belokopytov<sup>37</sup>, P.Beltran<sup>9</sup>, D.Benedic<sup>8</sup>,  
 J.M.Benloch<sup>42</sup>, M.Berggren<sup>38</sup>, D.Bertrand<sup>2</sup>, F.Bianchi<sup>39</sup>, M.S.Bilenky<sup>13</sup>, P.Billoir<sup>19</sup>, J.Bjarne<sup>20</sup>, D.Bloch<sup>8</sup>,  
 S.Blyth<sup>30</sup>, V.Bocci<sup>33</sup>, P.N.Bogolubov<sup>13</sup>, T.Bolognese<sup>34</sup>, M.Bonapart<sup>27</sup>, M.Bonesini<sup>25</sup>, W.Bonivento<sup>25</sup>,  
 P.S.L.Booth<sup>18</sup>, M.Borataw<sup>19</sup>, P.Borgeaud<sup>34</sup>, H.Borner<sup>30</sup>, C.Bosio<sup>35</sup>, B.Bostjancic<sup>7</sup>, O.Botner<sup>41</sup>, B.Bouquet<sup>16</sup>,  
 M.Bozzo<sup>10</sup>, S.Braibant<sup>7</sup>, P.Branchini<sup>35</sup>, K.D.Brand<sup>31</sup>, R.A.Brenner<sup>12</sup>, C.Bricman<sup>2</sup>, R.C.A.Brown<sup>7</sup>,  
 N.Brummer<sup>27</sup>, J-M.Brunet<sup>6</sup>, L.Bugge<sup>29</sup>, T.Buran<sup>29</sup>, H.Burmeister<sup>7</sup>, J.A.M.A.Buytaert<sup>2</sup>, M.Caccia<sup>7</sup>, M.Calvi<sup>25</sup>,  
 A.J.Camacho Rozas<sup>26</sup>, J-E.Campagne<sup>7</sup>, A.Campion<sup>18</sup>, T.Camporesi<sup>7</sup>, V.Canale<sup>33</sup>, F.Cao<sup>2</sup>, F.Carena<sup>7</sup>,  
 L.Carroll<sup>18</sup>, C.Caso<sup>10</sup>, E.Castelli<sup>40</sup>, M.V.Castillo Gimenez<sup>42</sup>, A.Cattai<sup>7</sup>, F.R.Cavallo<sup>5</sup>, L.Cerrito<sup>33</sup>, A.Chan<sup>1</sup>,  
 M.Chapkin<sup>37</sup>, P.Charpentier<sup>7</sup>, P.Checchia<sup>31</sup>, G.A.Chelkov<sup>13</sup>, L.Chevalier<sup>34</sup>, P.Chliapnikov<sup>37</sup>, V.Chorowicz<sup>19</sup>,  
 R.Cirio<sup>39</sup>, M.P.Clara<sup>39</sup>, P.Collins<sup>30</sup>, J.L.Contreras<sup>22</sup>, R.Contri<sup>10</sup>, G.Cosme<sup>16</sup>, F.Couchot<sup>16</sup>, H.B.Crawley<sup>1</sup>,  
 D.Crennell<sup>32</sup>, G.Crosetti<sup>10</sup>, M.Crozon<sup>6</sup>, J.Cuevas Maestro<sup>36</sup>, S.Czellar<sup>12</sup>, S.Dagoret<sup>16</sup>, E.Dahl-Jensen<sup>26</sup>,  
 B.Dalmagne<sup>16</sup>, M.Dam<sup>7</sup>, G.Damgaard<sup>26</sup>, G.Darbo<sup>10</sup>, E.Daubie<sup>2</sup>, P.D.Dauncey<sup>30</sup>, M.Davenport<sup>7</sup>, P.David<sup>19</sup>,  
 A.De Angelis<sup>40</sup>, M.De Beer<sup>34</sup>, H.De Boeck<sup>2</sup>, W.De Boer<sup>14</sup>, C.De Clercq<sup>2</sup>, M.D.M.De Fez Laso<sup>42</sup>, N.De Groot<sup>27</sup>,  
 C.De La Vaissiere<sup>19</sup>, B.De Lotto<sup>40</sup>, A.De Min<sup>25</sup>, C.Defoix<sup>6</sup>, D.Delikaris<sup>7</sup>, S.Delorme<sup>7</sup>, P.Delpierre<sup>6</sup>,  
 N.Demaria<sup>39</sup>, J.Derkaoui<sup>39,23</sup>, L.Di Ciaccio<sup>33</sup>, H.Dijkstra<sup>7</sup>, F.Djama<sup>8</sup>, J.Dolbeau<sup>6</sup>, M.Donszelmann<sup>27</sup>,  
 K.Doroba<sup>44</sup>, M.Dracos<sup>7</sup>, J.Drees<sup>45</sup>, M.Dris<sup>28</sup>, Y.Dufour<sup>9</sup>, W.Dulinski<sup>8</sup>, R.Dzhelyadin<sup>37</sup>, L-O.Eek<sup>41</sup>,  
 P.A.-M.Eerola<sup>12</sup>, T.Ekelof<sup>41</sup>, G.Ekspong<sup>38</sup>, A.Elliot Peisert<sup>31</sup>, J-P.Engel<sup>8</sup>, D.Fassouliotis<sup>28</sup>, M.Feindt<sup>7</sup>,  
 M.Fernandez Alonso<sup>36</sup>, A.Ferrer<sup>42</sup>, T.A.Filippas<sup>28</sup>, A.Firestone<sup>1</sup>, H.Foeth<sup>7</sup>, E.Fokitis<sup>28</sup>, P.Folegati<sup>40</sup>,  
 F.Fontanelli<sup>10</sup>, K.A.J.Forbes<sup>18</sup>, H.Forsbach<sup>45</sup>, B.Franek<sup>32</sup>, P.Frenkiel<sup>8</sup>, D.C.Fries<sup>14</sup>, A.G.Frodesen<sup>4</sup>,  
 R.Fruhworth<sup>43</sup>, F.Fulda-Quenzer<sup>16</sup>, K.Furnival<sup>18</sup>, H.Furstenau<sup>14</sup>, J.Fuster<sup>7</sup>, G.Galeazzi<sup>31</sup>, D.Gamba<sup>39</sup>,  
 C.Garcia<sup>42</sup>, J.Garcia<sup>36</sup>, U.Gasparini<sup>31</sup>, P.Gavillet<sup>7</sup>, E.N.Gazis<sup>28</sup>, J-P.Gerber<sup>8</sup>, P.Giacomelli<sup>5</sup>, K-W.Glitza<sup>45</sup>,  
 R.Gokiel<sup>7</sup>, V.M.Golovatyuk<sup>13</sup>, J.J.Gomez Y Cadenas<sup>7</sup>, A.Goobar<sup>38</sup>, G.Gopal<sup>32</sup>, M.Gorski<sup>44</sup>, V.Gracco<sup>10</sup>,  
 A.Grant<sup>7</sup>, F.Grad<sup>2</sup>, E.Graziani<sup>35</sup>, M-H.Gros<sup>16</sup>, G.Grosdidier<sup>16</sup>, E.Gross<sup>7</sup>, B.Grossetete<sup>19</sup>, J.Guy<sup>32</sup>, F.Hahn<sup>7</sup>,  
 M.Hahn<sup>14</sup>, S.Haider<sup>27</sup>, Z.Hajduk<sup>27</sup>, A.Hakansson<sup>20</sup>, A.Hallgren<sup>41</sup>, K.Hamacher<sup>45</sup>, G.Hamel De Monchenault<sup>34</sup>,  
 F.J.Harris<sup>30</sup>, B.W.Heck<sup>7</sup>, T.Henkes<sup>7</sup>, I.Herbst<sup>45</sup>, J.J.Hernandez<sup>42</sup>, P.Herquet<sup>2</sup>, H.Herr<sup>7</sup>, I.Hietanen<sup>12</sup>,  
 C.O.Higgins<sup>18</sup>, E.Higon<sup>42</sup>, H.J.Hilke<sup>7</sup>, S.D.Hodgson<sup>30</sup>, T.Hofmoki<sup>44</sup>, R.Holmes<sup>1</sup>, S-O.Holmgren<sup>38</sup>,  
 D.Holthuisen<sup>27</sup>, P.F.Honore<sup>6</sup>, J.E.Hooper<sup>28</sup>, M.Houlden<sup>18</sup>, J.Hrubec<sup>43</sup>, P.O.Hulth<sup>38</sup>, K.Hultqvist<sup>38</sup>, D.Husson<sup>8</sup>,  
 B.D.Hyams<sup>7</sup>, P.Ioannou<sup>3</sup>, D.Isenhower<sup>7</sup>, P-S.Iversen<sup>4</sup>, J.N.Jackson<sup>18</sup>, P.Jalocha<sup>15</sup>, G.Jarlskog<sup>20</sup>, P.Jarry<sup>34</sup>,  
 B.Jean-Marie<sup>16</sup>, E.K.Johansson<sup>38</sup>, D.Johnson<sup>18</sup>, M.Jonker<sup>7</sup>, L.Jonsson<sup>20</sup>, P.Juillot<sup>8</sup>, G.Kalkanis<sup>3</sup>, G.Kalmus<sup>32</sup>,  
 G.Kantardjian<sup>7</sup>, F.Kapusta<sup>19</sup>, S.Katsanevas<sup>3</sup>, E.C.Katsoufis<sup>28</sup>, R.Keranen<sup>12</sup>, J.Kesteman<sup>2</sup>, B.A.Khomenko<sup>13</sup>,  
 N.N.Khovanski<sup>13</sup>, B.King<sup>18</sup>, N.J.Kjaer<sup>26</sup>, H.Klein<sup>7</sup>, W.Klempt<sup>7</sup>, A.Klovning<sup>4</sup>, P.Kluit<sup>27</sup>, J.H.Koehne<sup>14</sup>,  
 B.Koene<sup>27</sup>, P.Kokkinias<sup>9</sup>, M.Kopf<sup>14</sup>, M.Koratzinos<sup>7</sup>, K.Korcyk<sup>15</sup>, A.V.Korytov<sup>13</sup>, B.Korsen<sup>7</sup>, V.Kostiukhin<sup>37</sup>,  
 C.Kourkoumelis<sup>3</sup>, T.Kreuzberger<sup>43</sup>, J.Krolkowski<sup>44</sup>, I.Kronkvist<sup>20</sup>, U.Kruener-Marquis<sup>45</sup>, W.Krupinski<sup>15</sup>,  
 W.Kucewicz<sup>26</sup>, K.Kurvinen<sup>12</sup>, C.Lacasta<sup>42</sup>, C.Lambropoulos<sup>9</sup>, J.W.Lamsa<sup>1</sup>, L.Lanceri<sup>40</sup>, V.Lapin<sup>37</sup>,  
 J-P.Laugier<sup>34</sup>, R.Lauhakangas<sup>12</sup>, G.Leder<sup>43</sup>, F.Ledroit<sup>6</sup>, J.Lemonne<sup>2</sup>, G.Lenzen<sup>45</sup>, V.Lepeltier<sup>16</sup>,  
 A.Letessier-Selvon<sup>19</sup>, D.Liko<sup>43</sup>, E.Lieb<sup>45</sup>, E.Lillethun<sup>4</sup>, J.Lindgren<sup>12</sup>, A.Lipniacka<sup>44</sup>, I.Lippi<sup>31</sup>, R.Llosa<sup>22</sup>,  
 B.Loerstad<sup>20</sup>, M.Lokajicek<sup>13</sup>, J.G.Loken<sup>30</sup>, M.A.Lopez Aguera<sup>36</sup>, A.Lopez-Fernandez<sup>16</sup>, M.Los<sup>27</sup>, D.Loukas<sup>9</sup>,  
 A.Lounis<sup>8</sup>, J.J.Lozano<sup>42</sup>, R.Lucock<sup>32</sup>, P.Lutz<sup>6</sup>, L.Lyons<sup>30</sup>, G.Maehlum<sup>7</sup>, J.Maillard<sup>6</sup>, A.Maltezos<sup>9</sup>, F.Mandl<sup>43</sup>,  
 J.Marco<sup>36</sup>, M.Margoni<sup>31</sup>, J-C.Marin<sup>7</sup>, A.Markou<sup>9</sup>, S.Marti<sup>42</sup>, L.Mathis<sup>6</sup>, F.Matorras<sup>36</sup>, C.Matteuzzi<sup>25</sup>,  
 G.Matthiae<sup>33</sup>, M.Mazzucato<sup>31</sup>, M.Mc Cubbin<sup>18</sup>, R.Mc Kay<sup>1</sup>, R.Mc Nulty<sup>18</sup>, E.Menichetti<sup>39</sup>, G.Meola<sup>10</sup>,  
 C.Meroni<sup>25</sup>, W.T.Meyer<sup>1</sup>, M.Michelotto<sup>31</sup>, W.A.Mitaroff<sup>13</sup>, G.V.Mitselmakher<sup>13</sup>, U.Mjoernmark<sup>20</sup>, T.Moa<sup>38</sup>,  
 R.Moeller<sup>26</sup>, K.Moenig<sup>45</sup>, M.R.Monge<sup>10</sup>, P.Morettini<sup>10</sup>, H.Mueller<sup>14</sup>, H.Muller<sup>7</sup>, W.J.Murray<sup>32</sup>, G.Myatt<sup>30</sup>,  
 F.Naraghi<sup>19</sup>, U.Nau-Korzen<sup>45</sup>, F.L.Navarria<sup>5</sup>, P.Negri<sup>25</sup>, B.S.Nielsen<sup>26</sup>, B.Nijjhar<sup>18</sup>, V.Nikolaenko<sup>37</sup>,  
 V.Obrastsov<sup>37</sup>, A.G.Olshevski<sup>13</sup>, R.Orava<sup>12</sup>, A.Ostankov<sup>37</sup>, A.Ouraou<sup>34</sup>, R.Pain<sup>19</sup>, H.Palka<sup>27</sup>,  
 T.Papadopoulou<sup>28</sup>, L.Pape<sup>7</sup>, A.Passeri<sup>35</sup>, M.Pegoraro<sup>31</sup>, V.Perevozchikov<sup>37</sup>, M.Pernicka<sup>43</sup>, A.Perrotta<sup>5</sup>,  
 F.Pierre<sup>34</sup>, M.Pimenta<sup>17</sup>, O.Pingot<sup>2</sup>, M.E.Pol<sup>7</sup>, G.Polok<sup>15</sup>, P.Poropat<sup>40</sup>, P.Privitera<sup>14</sup>, A.Pullia<sup>25</sup>, J.Pyyhtia<sup>12</sup>,  
 D.Radojicic<sup>30</sup>, S.Ragazzi<sup>25</sup>, P.N.Ratoff<sup>30</sup>, A.L.Read<sup>29</sup>, N.G.Redaeli<sup>25</sup>, M.Regler<sup>43</sup>, D.Reid<sup>18</sup>, P.B.Renton<sup>30</sup>,  
 L.K.Resvanis<sup>3</sup>, F.Richard<sup>16</sup>, M.Richardson<sup>18</sup>, J.Ridky<sup>13</sup>, G.Rinaudo<sup>39</sup>, I.Roditi<sup>7</sup>, A.Romero<sup>39</sup>, I.Roncagliolo<sup>10</sup>,  
 P.Ronchese<sup>31</sup>, C.Ronnqvist<sup>12</sup>, E.I.Rosenberg<sup>1</sup>, U.Rossi<sup>5</sup>, E.Rosso<sup>7</sup>, P.Roudeau<sup>16</sup>, T.Rovelli<sup>5</sup>, W.Ruckstuhl<sup>27</sup>,  
 V.Ruhlmann<sup>34</sup>, A.Ruiz<sup>36</sup>, K.Rybicki<sup>15</sup>, H.Saarikko<sup>12</sup>, Y.Sacquin<sup>34</sup>, G.Sajot<sup>11</sup>, J.Salt<sup>42</sup>, E.Sanchez<sup>42</sup>,  
 J.Sanchez<sup>22</sup>, M.Sannino<sup>10</sup>, M.Schaeffer<sup>8</sup>, S.Schael<sup>14</sup>, H.Schneider<sup>14</sup>, M.A.E.Schyns<sup>7</sup>, F.Scuri<sup>40</sup>, A.M.Segar<sup>30</sup>,  
 R.Sekulin<sup>32</sup>, M.Sessa<sup>40</sup>, G.Sette<sup>10</sup>, R.Seufert<sup>14</sup>, P.Siegrist<sup>34</sup>, S.Simonetti<sup>10</sup>, F.Simonetto<sup>31</sup>, A.N.Sissakian<sup>13</sup>,  
 T.B.Skaali<sup>29</sup>, G.Skjjevling<sup>29</sup>, G.Smadja<sup>34,21</sup>, G.R.Smith<sup>32</sup>, N.Smirnov<sup>37</sup>, R.Sosnowski<sup>44</sup>, T.S.Spassoff<sup>11</sup>,  
 E.Spiriti<sup>35</sup>, S.Squarcia<sup>10</sup>, H.Staack<sup>45</sup>, C.Stanescu<sup>35</sup>, G.Stavropoulos<sup>9</sup>, F.Stichelbaut<sup>2</sup>, A.Stocchi<sup>16</sup>, J.Strauss<sup>43</sup>,

R.Strub<sup>8</sup>, C.J.Stubenrauch<sup>7</sup>, M.Szczekowski<sup>44</sup>, M.Szeptycka<sup>44</sup>, P.Szymanski<sup>44</sup>, T.Tabarelli<sup>25</sup>, S.Tavernier<sup>2</sup>, G.Theodosiou<sup>9</sup>, A.Tilquin<sup>24</sup>, J.Timmermans<sup>27</sup>, V.G.Timofeev<sup>13</sup>, L.G.Tkatchev<sup>13</sup>, T.Todorov<sup>13</sup>, D.Z.Toet<sup>27</sup>, A.Tomaradze<sup>37</sup>, L.Tortora<sup>35</sup>, M.T.Trainor<sup>30</sup>, D.Treille<sup>7</sup>, U.Trevisan<sup>10</sup>, W.Trischuk<sup>7</sup>, G.Tristram<sup>6</sup>, C.Troncon<sup>25</sup>, A.Tsirou<sup>7</sup>, E.N.Tsyganov<sup>13</sup>, M.Turala<sup>15</sup>, R.Turchetta<sup>8</sup>, M-L.Turluer<sup>34</sup>, T.Tuuva<sup>12</sup>, I.A.Tyapkin<sup>13</sup>, M.Tyndel<sup>32</sup>, S.Tzamarias<sup>7</sup>, B.Ueberschaer<sup>45</sup>, S.Ueberschaer<sup>45</sup>, O.Ullaland<sup>7</sup>, V.A.Uvarov<sup>37</sup>, G.Valenti<sup>5</sup>, E.Vallazza<sup>39</sup>, J.A.Valls Ferrer<sup>42</sup>, G.W.Van Apeldoorn<sup>27</sup>, P.Van Dam<sup>27</sup>, W.K.Van Doninck<sup>2</sup>, N.Van Eijndhoven<sup>7</sup>, C.Vander Velde<sup>2</sup>, J.Varela<sup>17</sup>, P.Vaz<sup>7</sup>, G.Vegni<sup>25</sup>, L.Ventura<sup>31</sup>, W.Venus<sup>32</sup>, F.Verbeure<sup>2</sup>, L.S.Vertogradov<sup>13</sup>, L.Vibert<sup>19</sup>, D.Vilanova<sup>34</sup>, N.Vishnevsky<sup>37</sup>, E.V.Vlasov<sup>37</sup>, S.Vlassopoulos<sup>28</sup>, A.S.Vodopyanov<sup>13</sup>, M.Vollmer<sup>45</sup>, S.Volponi<sup>5</sup>, G.Voulgaris<sup>3</sup>, M.Voutilainen<sup>12</sup>, V.Vrba<sup>35</sup>, H.Wahlen<sup>45</sup>, C.Walck<sup>38</sup>, F.Waldner<sup>40</sup>, M.Wayne<sup>1</sup>, P.Weilhammer<sup>7</sup>, J.Werner<sup>45</sup>, A.M.Wetherell<sup>7</sup>, J.H.Wickens<sup>2</sup>, J.Wikne<sup>29</sup>, G.R.Wilkinson<sup>30</sup>, W.S.C.Williams<sup>30</sup>, M.Winter<sup>8</sup>, D.Wormald<sup>29</sup>, G.Wormser<sup>16</sup>, K.Woschnagg<sup>41</sup>, N.Yamdagni<sup>38</sup>, P.Yepes<sup>7</sup>, A.Zaitsev<sup>37</sup>, A.Zalewska<sup>15</sup>, P.Zalewski<sup>44</sup>, D.Zavrtanik<sup>7</sup>, E.Zevgolatakos<sup>9</sup>, G.Zhang<sup>45</sup>, N.I.Zimin<sup>13</sup>, M.Zito<sup>34</sup>, R.Zitoun<sup>19</sup>, R.Zukanovich Funchal<sup>6</sup>, G.Zumerle<sup>31</sup>, J.Zuniga<sup>42</sup>

<sup>1</sup> Ames Laboratory and Department of Physics, Iowa State University, Ames IA 50011, USA

<sup>2</sup> Physics Department, Univ. Instelling Antwerpen, Universiteitsplein 1, B-2610 Wilrijk, Belgium and IHE, ULB-VUB, Pleinlaan 2, B-1050 Brussels, Belgium

and Service de Phys. des Part. Elém., Faculté des Sciences, Université de l'Etat Mons, Av. Maistriau 19, B-7000 Mons, Belgium

<sup>3</sup> Physics Laboratory, University of Athens, Solonos Str. 104, GR-10680 Athens, Greece

<sup>4</sup> Department of Physics, University of Bergen, Allégaten 55, N-5007 Bergen, Norway

<sup>5</sup> Dipartimento di Fisica, Università di Bologna and INFN, Via Irnerio 46, I-40126 Bologna, Italy

<sup>6</sup> Collège de France, Lab. de Physique Corpusculaire, 11 pl. M. Berthelot, F-75231 Paris Cedex 05, France

<sup>7</sup> CERN, CH-1211 Geneva 23, Switzerland

<sup>8</sup> Division des Hautes Energies, CRN - Groupe DELPHI and LEPsi, B.P.20 CRO, F-67037 Strasbourg Cedex, France

<sup>9</sup> Institute of Nuclear Physics, N.R.C. Demokritos, P.O. Box 60628, GR-15310 Athens, Greece

<sup>10</sup> Dipartimento di Fisica, Università di Genova and INFN, Via Dodecaneso 33, I-16146 Genova, Italy

<sup>11</sup> Institut des Sciences Nucléaires, Université de Grenoble 1, F-38026 Grenoble, France

<sup>12</sup> Research Institute for High Energy Physics, University of Helsinki, Siltavuorenpenger 20 C, SF-00170 Helsinki 17, Finland

<sup>13</sup> Joint Institute for Nuclear Research, Dubna, Head Post Office, P.O. Box 79, 101 000 Moscow, USSR.

<sup>14</sup> Institut für Experimentelle Kernphysik, Universität Karlsruhe, Postfach 6980, D-7500 Karlsruhe 1, FRG

<sup>15</sup> High Energy Physics Laboratory, Institute of Nuclear Physics, Ul. Kawiora 26 a, PL-30055 Krakow 30, Poland

<sup>16</sup> Université de Paris-Sud, Lab. de l'Accélérateur Linéaire, Bat 200, F-91405 Orsay, France

<sup>17</sup> LIP, Av. Elias Garcia 14 - 1e, P-1000 Lisbon Codex, Portugal

<sup>18</sup> Department of Physics, University of Liverpool, P.O. Box 147, GB - Liverpool L69 3BX, UK

<sup>19</sup> LPNHE, Universités Paris VI et VII, Tour 33 (RdC), 4 place Jussieu, F-75230 Paris Cedex 05, France

<sup>20</sup> Department of Physics, University of Lund, Sölvegatan 14, S-22363 Lund, Sweden

<sup>21</sup> Université Claude Bernard de Lyon, 43 Bd du 11 Novembre 1918, F-69622 Villeurbanne Cedex, France

<sup>22</sup> Departamento de Fisica Atomica Molecular y Nuclear, Universidad Complutense, Avda. Complutense s/n, E-28040 Madrid, Spain

<sup>23</sup> Permanent address: Département de Physique, Faculté des Sciences d'Oujda, Maroc

<sup>24</sup> Faculté des Sciences de Luminy, Univ. d'Aix - Marseille II Case 907 - 70, route Léon Lachamp, F-13288 Marseille Cedex 09, France

<sup>25</sup> Dipartimento di Fisica, Università di Milano and INFN, Via Celoria 16, I-20133 Milan, Italy

<sup>26</sup> Niels Bohr Institute, Blegdamsvej 17, DK-2100 Copenhagen 0, Denmark

<sup>27</sup> NIKHEF-H, Postbus 41882, NL-1009 DB Amsterdam, The Netherlands

<sup>28</sup> National Technical University, Physics Department, Zografou Campus, GR-15773 Athens, Greece

<sup>29</sup> Physics Department, University of Oslo, Blindern, N-1000 Oslo 3, Norway

<sup>30</sup> Nuclear Physics Laboratory, University of Oxford, Keble Road, GB - Oxford OX1 3RH, UK

<sup>31</sup> Dipartimento di Fisica, Università di Padova and INFN, Via Marzolo 8, I-35131 Padua, Italy

<sup>32</sup> Rutherford Appleton Laboratory, Chilton, GB - Didcot OX11 0QX, UK

<sup>33</sup> Dipartimento di Fisica, Università di Roma II and INFN, Tor Vergata, I-00173 Rome, Italy

<sup>34</sup> CEN-Saclay, DPhPE, F-91191 Gif-sur-Yvette Cedex, France

<sup>35</sup> Istituto Superiore di Sanità, Ist. Naz. di Fisica Nucl. (INFN), Viale Regina Elena 299, I-00161 Rome, Italy

<sup>36</sup> Facultad de Ciencias, Universidad de Santander, av. de los Castros, E - 39005 Santander, Spain

<sup>37</sup> Inst. for High Energy Physics, Serpukov P.O. Box 35, Protvino, (Moscow Region), USSR.

<sup>38</sup> Institute of Physics, University of Stockholm, Vanadisvägen 9, S-113 46 Stockholm, Sweden

<sup>39</sup> Dipartimento di Fisica Sperimentale, Università di Torino and INFN, Via P. Giuria 1, I-10125 Turin, Italy

<sup>40</sup> Dipartimento di Fisica, Università di Trieste and INFN, Via A. Valerio 2, I-34127 Trieste, Italy

and Istituto di Fisica, Università di Udine, I-33100 Udine, Italy

<sup>41</sup> Department of Radiation Sciences, University of Uppsala, P.O. Box 535, S-751 21 Uppsala, Sweden

<sup>42</sup> Inst. de Fisica Corpuscular IFIC, Centro Mixto Univ. de Valencia-CSIC, and Departamento de Fisica Atomica Molecular y Nuclear, Univ. de Valencia, Avda. Dr. Moliner 50, E-46100 Burjassot (Valencia), Spain

<sup>43</sup> Institut für Hochenergiephysik, Österreich Akad. d. Wissensch., Nikolsdorfergasse 18, A-1050 Vienna, Austria

<sup>44</sup> Inst. Nuclear Studies and, University of Warsaw, Ul. Hoza 69, PL-00681 Warsaw, Poland

<sup>45</sup> Fachbereich Physik, University of Wuppertal, Postfach 100 127, D-5600 Wuppertal 1, FRG

# 1 Introduction

Scalar coloured leptoquarks appear naturally as a common feature of various theories extending the standard model. Besides being present in the strong coupling version of the standard model, they also emerge in grand unifying models, in composite models, in technicolor or horizontal symmetry schemes and more recently, in  $E_6$  superstring inspired models [1][2].

Since the leptoquark couples to a quark and a lepton, leptoquark pair production at  $Z^0$  could provide a characteristic signal with an unlike-sign lepton pair which is accompanied by distant jet activity from the hadronization of the two quarks and could involve no missing energy.

As a guide in this search,  $E_6$  compactification [2] results are used to consider mass degenerate scalar elementary leptoquarks  $D_0$   $D_0^c$  in the first two families. Each one is assumed to be a weak-isospin singlet, color triplet, of charge  $-1/3$  and  $+1/3$  respectively, with a branching ratio  $2/3$  to the up quark and the charged lepton of its family. The other decay mode would be to the down quark and neutrino of the family with branching ratio  $1/3$ .  $D_0$ ,  $D_0^c$  and  $D_{\frac{1}{2}}$  constitute the new supersymmetric multiplet predicted by  $E_6$ .  $D_0$  and  $D_0^c$  are the two supersymmetric partners of the corresponding fermion field  $D_{\frac{1}{2}}$  in the same way as the left and right-handed selectrons correspond to the electron field.

At the  $Z^0$  peak, the pair production cross-section [3] is almost independent of the unknown Yukawa coupling  $\lambda$  of the leptoquark to the lepton and the quark, depending mostly on the  $Z^0 D_0 \bar{D}_0$  gauge coupling  $c$ , i.e.

$$\frac{d\sigma}{d\cos\theta} = 3\pi \frac{\alpha^2}{4s} \beta^3 \sin^2\theta \frac{s^2}{(s-M^2)^2 + M^2\Gamma^2} c^2 (v^2 + a^2)$$

Here  $\theta$  is the production angle of the  $D_0$  with respect to the  $e^-$  beam,  $\beta = \sqrt{1 - 4m^2/s}$  is the center-of-mass velocity of the leptoquark,

$$v = \frac{1}{\sin\theta_w \cos\theta_w} \left( \frac{1}{4} - \sin^2\theta_w \right), \quad a = \frac{1}{4 \sin\theta_w \cos\theta_w}$$

are the standard model electron couplings and

$$c = \frac{1}{\sin\theta_w \cos\theta_w} (I_3^W - Q \sin^2\theta_w)$$

is fixed by gauge symmetry for the leptoquark of charge  $Q$  and weak isospin  $I_3^W$ . The above expressions are valid for the leptoquarks of most other models. Phenomenologically, the leptoquark coupling  $g^2 = c^2$  to the neutral current and the branching ratio into quarks and charged leptons could be considered as free parameters.

In the following, besides searching for pairs of scalar leptoquarks, one also looked for the process

$$\begin{aligned} e^+e^- &\rightarrow D_0 \ell^+ \bar{q} & \text{or} & \quad \bar{D}_0 \ell^- q & \quad (+D_0 \leftrightarrow D_0^c) \\ &\hookrightarrow \ell^- q & & \quad \hookrightarrow \ell^+ \bar{q} \end{aligned}$$

where only one of the leptoquarks  $D_0$ ,  $\bar{D}_0$ , ( $D_0^c$ ,  $\bar{D}_0^c$ ), is produced on-shell and thus probe [4] part of the mass region  $\frac{1}{2}m_{Z^0} < m_D < m_{Z^0}$ . The cross-section for this process depends on the  $D_0 \ell q$  Yukawa coupling  $\lambda$  and is proportional to  $\alpha_\lambda = \lambda^2/4\pi$ , assumed equal to  $\alpha$  (fine structure constant) [3][4].

The width of the scalar leptoquark is of the order  $\alpha_\lambda m$  which corresponds to a life-time of  $\sim 10^{-23}$  sec for a mass of 40 GeV/ $c^2$ .

Various experimental constraints have been imposed on scalar composite leptoquarks with  $Q = 2/3$ . The JADE Collaboration [5] has excluded masses in the range 5-20.8 GeV/ $c^2$  for a leptoquark decaying into second family fermions. The UA1 Collaboration [6] has also placed a limit of  $m > 33$  GeV/ $c^2$  for second family composite leptoquarks with  $Q=2/3$  and  $BR(\mu + \text{quark}) \geq 0.1$ . KEK experiments from the AMY and VENUS Collaborations [7] searching for similar  $Q = 2/3$  composite leptoquarks and considering only the  $\gamma$  propagator, imply a limit  $m > 21$  GeV/ $c^2$  for  $Q = -1/3$  and  $BR = 2/3$  in the second family at KEK energies.

## 2 The Detector

The DELPHI detector is described in detail elsewhere [8]; here we just give a brief summary of the tracking detectors and calorimeters used in this analysis. Charged particle tracks are measured in the 1.2 T magnetic field with three cylindrical detectors: the Inner Detector (ID) covering polar angles between  $29^\circ$  and  $151^\circ$ , the Time Projection Chamber (TPC) covering angles between  $21^\circ$  and  $159^\circ$  and the Outer Detector (OD) covering angles between  $42^\circ$  and  $138^\circ$ .

The calorimeters are the High Density Projection Chamber (HPC), which measures the electromagnetic energy with high granularity covering angles between  $40^\circ$  and  $140^\circ$ , the Electromagnetic Calorimeter in the endcaps (EMF), consisting of  $2 \times 4500$  lead glass blocks and covering angles between  $10^\circ$  and  $35.5^\circ$  and between  $144.5^\circ$  and  $170^\circ$ , and the Hadron Calorimeter (HCAL), measuring the hadronic energy and covering all angles between  $10^\circ$  and  $170^\circ$ .

## 3 Data Sample

The exotic events were sought in a hadronic sample that initially was selected from the 1990 data so that:

- i) the number of charged particle tracks was at least five.
- ii) all the particles, charged and neutral, had to form at least two clusters as defined by the LUND cluster algorithm LUCLUS [9] used with default parameters ( $d_{join}=2.5$  GeV). This condition is very efficient in eliminating beam-gas events.

Charged particles were considered if they had momentum exceeding 100 MeV/ $c$  and if their extrapolated distance to the vertex in the transverse plane and along the beam axis was less than 10 cm each.

To estimate the number of produced  $Z^0$  hadronic events to which the above selected sample corresponds, the preceding cuts as well as a  $|\cos \theta_{thrust}| < 0.8$  cut were applied to the 1990 data sample taken on and around the  $Z^0$  mass peak. This left a sample of 88195 hadronic decays of  $Z^0$ . In a simulated sample of 40K hadronic reconstructed events the above thrust cut led to a 22.9% loss. This taken together with a 1.54% loss due to (i) + (ii) and with a hadronic trigger efficiency of 99.5%, led to a total number of  $116\,175 \pm 445$   $Z^0$  hadronic events produced.

## 4 Search for leptoquark pairs

The topology that was looked for was that of two jets and two isolated particles. Such events were sought within four jets reconstructed from charged and neutral particles using the LUCCLUS algorithm with default parameters. The event charged particle multiplicity was required to be at least six and the charged multiplicity of each jet at least one. In addition,

- i) Two of the reconstructed jets should be single charged particles of opposite sign. These two particles were then considered as isolated if they satisfied (iii), as well.
- ii) In the samples of Monte Carlo simulated  $D_0 \bar{D}_0$  pairs it was found that one (case iia) or both leptons (case iib) from the  $D_0 \bar{D}_0$  decays were often interpreted as one or two jets, respectively. The clustering algorithm assigned to them several very low energy fragments produced at wide angles from the fragmentation of the quarks. Therefore if condition (i) was not satisfied, the jets were further examined: if a jet had at most total multiplicity 3 and the charged leader in the jet carried at least 90% of the jet energy and satisfied the kinematic conditions of (iii), then this "jet" was considered as an isolated particle (case iia). If two such isolated particles could be recovered, the event would be classified in case iib. Figure 1 shows the distribution for the total multiplicity for the hadronic jets or isolated particles defined as above in the 180 selected MC events out of a sample of 428 for  $m = 42 \text{ GeV}/c^2$ . The two peaks seen are due to the two isolated particles and the two jets.
- iii) Both isolated particles from case (i) or (ii) were required to satisfy the angular condition with respect to the beams  $|\cos\theta_i| < 0.85$  and have momentum greater than  $3 \text{ GeV}/c$ . The hadronic jets were also required to have energy greater than  $3 \text{ GeV}$ .

The efficiencies with these selection criteria were studied using samples of events at various masses  $m$  generated with a Monte Carlo program using a  $D_0 \bar{D}_0 \rightarrow c\mu^- \bar{c}\mu^+$  generator which was based on LUND program[9] Jetset 6.3 for hadronization. Subsequently, the detailed detector simulation was used and the events were processed through the same processing chain as the real data. It was found that the events with two isolated particles of class i satisfying the above selection criteria, also had both particles satisfying the condition  $\rho > 1.8 \text{ GeV}^{\frac{1}{2}}$  on the isolation parameter

$$\rho = \sqrt{2E_{\text{isol}}(1 - \cos\theta_{\text{isol,jet}})}$$

where  $\theta_{\text{isol,jet}}$  is the minimum angle between an isolated particle and the hadronic jets and  $E_{\text{isol}}$  is the energy of the isolated particle in GeV. This condition  $\rho > 1.8 \text{ GeV}^{\frac{1}{2}}$  was then applied on the events in the Monte Carlo samples (iia) and (iib).

The same selection procedure was applied on the hadronic data sample and on a Monte Carlo sample of 97 000 hadronic  $Z^0$  events. Compatible numbers of candidates were obtained from the two samples, when the selection conditions were relaxed from the ones mentioned above. For example, when the leader in the candidate for isolated particle had energy greater than 50 % of the jet energy and this jet had multiplicity up to five, then the numbers obtained from the two samples were  $185 \pm 14$  and  $165 \pm 13$  (normalized), respectively.

The main background for this signal comes from the semileptonic decays of the  $b\bar{b}$  and  $c\bar{c}$  events giving topologies of 2 jets + 2 charged leptons. Using a sample of a mixture of 2000 such Monte Carlo events where both heavy quarks decay semileptonically, a simulation program allowed to estimate that, in this hadronic sample, approximately 2

heavy flavour events having 2 jets and 2 isolated leptons with isolation parameter  $\rho > 1.8 \text{ GeV}^{\frac{1}{2}}$  were expected. Concerning the four-fermion background,  $e^+e^- \rightarrow q\bar{q}\ell^+\ell^-$ , it is expected [10] to contribute at most 0.3 events, for  $\ell = \mu$  or  $e$  and under similar cuts.

With the conditions i-iii and  $\rho > 1.8 \text{ GeV}^{\frac{1}{2}}$  applied, no candidate remained from the data sample, whereas 1 event in class (iia) survived from the Monte Carlo hadronic  $Z^0$  sample. This Monte Carlo event was found to be of the type  $Z^0 \rightarrow b\bar{b}$ .

The detection efficiencies found for second family leptoquarks with masses up to  $42 \text{ GeV}/c^2$ , using conditions i-iii,  $\rho > 1.8 \text{ GeV}^{\frac{1}{2}}$  and without asking for lepton identification are given in Table 1. The error quoted is statistical only. Figure 2 gives the distribution of the  $\rho$  values of the isolated particles of the finally selected  $D_0 \bar{D}_0$  events in the Monte Carlo sample with leptoquark mass  $m=42 \text{ GeV}/c^2$ .

Table 1: Detection efficiency (%) for 2<sup>nd</sup> family leptoquarks of various masses (in  $\text{GeV}/c^2$ )

m	20	25	35	42	50	60
case i	6.5	5.5	10.2	7.5	11.1	5.9
case iia	16.4	15.0	19.3	18.9	17.7	12.2
case iib	14.5	12.5	10.2	11.9	8.3	8.3
Total	37.4	33.	39.7	38.3	37.1	26.4
	$\pm 3.3$	$\pm 3.2$	$\pm 2.0$	$\pm 2.2$	$\pm 2.9$	$\pm 2.5$

The detection efficiencies of Table 1 include a reduction by 10% to conservatively account for systematic uncertainties in the Monte Carlo hadronization [11], [13], coming from string fragmentation or independent fragmentation. For the first family,  $D_0 \bar{D}_0 \rightarrow ue^- \bar{u}e^+$ , a conservative estimate was made for the detection efficiencies by reducing further those of the second family by 15%, to account for the lower reconstruction efficiencies for two electrons relatively to two muons, as measured in  $Z^0 \rightarrow e^+e^-$  and  $Z^0 \rightarrow \mu^+\mu^-$  decays, respectively.

The number of  $D_0 \bar{D}_0$  events expected to be seen from the second family alone [12], as well as from the second and first combined (neglecting the mixing between mass-degenerate first and second family D's) are plotted in Figure 3 for the  $E_6$  inspired model of elementary leptoquarks [3]. The same number of events would generally be expected for any leptoquark with charge  $Q=-1/3$  and coupling  $c$  fixed by gauge symmetry (Section 1). In the same plot are given the expectations for a composite leptoquark with  $Q=2/3$  (just a factor of 4 larger). From these plots one finds the mass limits 42.5, 43, 43  $\text{GeV}/c^2$  for the three cases, respectively, at 95% confidence level. Inclusion of soft photon bremsstrahlung from the initial state reduces the expected leptoquark cross section by 30 to 40% (depending on the photon energy cut-off), which amounts to reducing the mass limits by at most 5%.

In Figure 4, the model independent cross section limits are given for any pair-produced spin-0 object of mass  $m$  decaying to a quark and a charged lepton in the mass interval 20-43  $\text{GeV}/c^2$  with  $\text{BR}=1$ , (second family, [12]) as well as the limit for the branching ratio into a quark and a charged lepton, at 95% confidence level (second family, [12]). Using these values, limits can be placed on the coupling times branching ratio  $|g| \cdot \text{BR}(D_0 \rightarrow q\ell)$  for a leptoquark, where  $g$  and  $\text{BR}$  are treated as free parameters (second family, [12]). These are plotted in the same Figure using the scale on the right side.

## 5 Search for a Singly Produced Leptoquark

This search has been extended to leptoquark masses higher than  $m_{Z^0}/2$ , by applying the same selection procedure as for  $D_0 \bar{D}_0$  above, to Monte Carlo samples of events with one real leptoquark with  $m=50$  and  $60 \text{ GeV}/c^2$ . These were generated according to the matrix element of reference [4] inspired by  $E_6$  superstrings and then processed as the real data, after a detailed simulation of the detector. The detection efficiencies, reduced by 10% to account for systematic uncertainties as for the pair production case, are given in Table 1.

In Figure 3, the expected number of events is given[12], neglecting the mixing between leptoquarks of the first and second family, assuming that they are mass degenerate and taking a Yukawa coupling  $\lambda$  of the leptoquark to the lepton and the quark such that  $\alpha_\lambda = \lambda^2/4\pi = \alpha$ .

In Figure 4, the cross section limits for the second family have been included[12] which are found for the single  $D_0$  process based on the detection efficiencies of Table 1. The expected total cross section values [4] at 50 and  $60 \text{ GeV}/c^2$  for each family (including  $D_0, \bar{D}_0, D_0^c$  and  $\bar{D}_0^c$ ), corrected by Monte Carlo for the cuts applied in [4], are 2.2 and 1.1 pb, respectively, for  $\alpha_\lambda = \alpha$ . The upper limits found on the Yukawa coupling  $\lambda$  of the leptoquark to the lepton and the quark, are  $\alpha_\lambda < 0.6\alpha$ , for  $m=50 \text{ GeV}/c^2$  and  $\alpha_\lambda < 1.6\alpha$  for  $m=60 \text{ GeV}/c^2$  at 95% confidence level, if the leptoquarks of the first and second family have the same mass and we neglect mixing.

## 6 Conclusions

We have searched for pair produced scalar leptoquarks in event topologies with two jets and two isolated charged particles. No candidate was found for the first or second family. Model independent cross section limits at 95% confidence level are given ( Figure 4) of the order of 2 pb in the leptoquark mass interval 20-45  $\text{GeV}/c^2$ . Also, branching ratio limits, as well as  $|g|.BR$  have been deduced, when the coupling  $|g|$  and BR are free parameters in the latter case.

Mass limits have been set at about 42-43  $\text{GeV}/c^2$  for  $E_6$  superstring inspired or composite models (Figure 3) for a branching ratio of 2/3 for decay into a quark and a charged lepton. Due to the steep fall-off of the cross section near  $m_{Z^0}/2$  because of the  $\beta^3$  factor, our bound depends only slightly on the particular model considered. Similar results have been obtained by other LEP experiments[14].

We have also searched for one on-mass-shell  $E_6$ -superstring inspired leptoquark, in a mass region above  $m_{Z^0}/2$  where cross section limits are found and upper limits are set on the coupling  $\lambda^2/4\pi$  at 50 and  $60 \text{ GeV}/c^2$  of  $0.6\alpha$  and  $1.6\alpha$ , respectively at 95% confidence level, if mass degeneracy is assumed for the leptoquarks of the first and second family.

## Acknowledgements

We are greatly indebted to our technical staff and collaborators and funding agencies for their support in building and operating the DELPHI detector, and to the members of



the CERN-SL Division for the superb performance of the LEP collider. We are grateful to N. D. Tracas for useful discussions.

## References

- [1] J. Pati and A. Salam, *Phys. Rev.* **D10** (1974) 275;  
L. F. Abbot and E. Fahri, *Phys. Lett.* **B101** (1981) 69; *Nucl. Phys.* **B189** (1981) 547;  
I. Bars, in *Proceed. 1984 Summer Study on the Design and Utilization of the SSC* (Snowmass, Co.) Eds. R. Donaldson and J. Mortin (APF, New York, 1985) p. 38;  
S. Dimopoulos and L. Susskind, *Nucl. Phys.* **B155** (1979) 237;  
S. Pakvasa, *Int. J. Mod. Phys.* **A2** (1987) 1317;  
"Physics at LEP 1", CERN 89-08, Vol.2: Higgs Search and New Physics, and references therein (1989);  
B. Schrempp and F. Schrempp, *Phys. Lett.* **153B** (1985) 101.
- [2] V. Barger, N.G. Deshpande and F.G. Gunion, in *Proc., 1986 Summer Study on the Design and Utilization of the SSC* (Snowmass, Co, 1986);  
V. D. Angelopoulos et al. *Nucl. Phys.* **B292** (1987) 59.
- [3] D. Schaile and P. Zerwas, in *Proceed., Workshop on Physics on Future Accelerators* (La Thuile and Geneva) CERN 87-07, Vol. II, p. 251 (1987);  
J. L. Hewett and T. G. Rizzo, *Phys. Rev.* **D36** (1987) 3367;  
J.L. Hewett and T.G. Rizzo, *Phys. Rep* **183** (1989)195.
- [4] N.D. Tracas and S. D. P. Vlassopoulos, *Phys. Lett* **B220** (1989) 285.
- [5] JADE Collab., W. Bartel et al, *Z. Phys* **C36** (1987) 15.
- [6] UA1 Collab., talk presented by S. Geer at the Intern. Europhysics Conf. on High Energy Physics (Uppsala, Sweden, 1987).
- [7] AMY Collab., G. N. Kim et al., *Phys. Lett* **B240** (1990)243;  
T. Tsuboyama, *Proceedings of the Rencontre de Moriond*, Feb. 1990, p. 199.
- [8] P. Aarnio et al. (DELPHI Collaboration), *Phys. Lett.* **B240** (1990) 271;  
P. Abreu et al. (DELPHI Collaboration), *Phys. Lett.* **B241** (1990) 435;  
DELPHI Collaboration, *Nucl. Inst. Meth.* **A303** (1991) 233.
- [9] T. Sjöstrand, *Comp. Phys. Comm.* 27 (1982) 243, *ibid.*, 28 (1983) 229;  
T. Sjöstrand and M. Bengtsson, *Comp. Phys. Comm.*, 43 (1987) 367.
- [10] E.N. Argyres et al., *Nucl. Phys.* **B354** (1991) 1.
- [11] T. Sjöstrand, private communication.
- [12] The number of events for the first family are obtained from those of the second by multiplying with 0.85. The first family cross section limits are obtained from those of the second by dividing by 0.85 and the BR limits by dividing those of the second family by 0.92.
- [13] The leptoquark events were generated with independent fragmentation. We have used string fragmentation in comparable samples of events with  $m=25,35$  and  $42 \text{ GeV}/c^2$  and a fast detector simulation, and observed a relative decrease in the detection efficiency of at most 10%.
- [14] L3 Collab., B. Adeva et al., L3 Preprint#26 (revised version) March 1991;  
OPAL Collab., G. Alexander et al., CERN-PPE/91-61 (1991).

## Figure captions

Figure 1 Multiplicity distribution for each of the 4 jets in the selected MC events at the leptoquark mass  $m=42 \text{ GeV}/c^2$  (4 entries per event).

Figure 2 Distribution of the isolation parameter  $\rho_1$  vs  $\rho_2$  of the two isolated particles for the selected MC events at  $m=42 \text{ GeV}/c^2$ .

Figure 3 Expected number of events from the hadronic sample of 116 000 produced  $Z^0$  with 2 jets + 2 charged isolated particles,  $\text{BR}=2/3$ .

- a)  $D_0\bar{D}_0 \rightarrow 2 \text{ jets} + 2 \text{ charged isolated particles}$  within the second family for the  $E_6$  inspired model [3],  $Q=-1/3$   $\text{BR}=2/3$ , or, for a composite leptoquark with  $Q=-1/3$ ,  $\text{BR}=2/3$  and coupling  $c$  fixed by gauge symmetry(left scale).
- b) As in a) but combining the two families when the first and second family  $D$ 's are mass degenerate(left scale).
- c) As in a) but for a composite leptoquark with  $Q=2/3$ (left scale).
- d) Expected events with 2 jets and 2 charged isolated particles, when only one  $D_0$  is produced on shell for the  $E_6$  inspired model [4],  $Q=-1/3$ ,  $\text{BR}=2/3$  (first and second family  $D$ 's are mass degenerate),  $\alpha_\lambda = \alpha$ (right scale).
- e) Same as d) but for the second family only (i.e. no mass degeneracy of the  $D$ 's)(right scale).
- f) Line of 95% confidence level.

Figure 4 Limits at 95 % CL:

- a) Model independent cross section limits for a pair-produced scalar object decaying to a quark and a charged isolated particle (second family[12]) Arrows show excluded area in  $20-45 \text{ GeV}/c^2$  (left scale).
- b) Cross section limits in the  $45-60 \text{ GeV}/c^2$  interval for a singly produced scalar leptoquark decaying to a quark and a charged isolated particle (2nd family [12]) using efficiencies based on the kinematics of  $E_6$  inspired class of models [4](left scale).
- c) Model independent branching ratio limits for a scalar leptoquark to decay to a quark and a charged isolated particle (2nd family, right scale) [12].
- d)  $|g| \cdot \text{BR}$  limit for a scalar leptoquark with free coupling  $g$  decaying to a quark and a charged isolated particle (2nd family, right scale) [12].




Fig. 1




Fig. 2




Fig. 3




Fig. 4