

# SUPERSYMMETRIC MAJORON SIGNATURES AND SOLAR NEUTRINO OSCILLATIONS \*

A. Santamaria <sup>†</sup> and J. W. F. Valle<sup>‡</sup>

*Departament de Física Teòrica, Universitat de Valencia*

*46100 Burjassot, València, Spain*

## Abstract

Spontaneous R parity breaking in supergravity solves the solar neutrino problem through matter enhanced neutrino oscillations. The model may be tested in collider experiments and through “dynamical” effects associated with the existence of a weakly interacting Majoron. Apart from astrophysical effects, Majoron emission can produce observable changes in  $\mu$  and  $\tau$  decay spectra for parameter values that substantially reduce the solar neutrino flux. A signature of the model is the possible observation of the decay  $\mu \rightarrow e + \text{Majoron}$ .

---

\*This work was supported in part by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and also by CAYCIT, Spain, under Plan Movilizador de la Física de Altas Energías

<sup>†</sup>Also at IFIC, Universitat de València – CSIC. Address after October 1987, Physics Dept, Carnegie-Mellon University, Pittsburgh, PA.

<sup>‡</sup>Also at Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720, USA



## 1. Introduction

The solar neutrino problem may be resolved either with new physics to change the solar parameters [1] or with non-standard weak interactions to modify neutrino propagation properties. These include neutrino decay [2], neutrino magnetic moment [3] and neutrino oscillations [4,5,6]. While it is possible to build consistent models for fast neutrino decay [7], the required value of the neutrino magnetic moment needed to solve the solar neutrino problem is hard to reconcile with existing limits on neutrino mass. Neutrino decay, however, has also been rendered unlikely by the recent observation of neutrinos from the the SN1987A supernova [8] although, for large mixings, the model may still survive [9]. This may leave oscillations as the most likely explanation of the solar neutrino problem in terms of non-standard neutrino properties. Unfortunately it will be difficult to test experimentally for the oscillation hypothesis. If oscillations occur *in vacuo* then the relevant mass squared differences are too small to be probed [6]. A lot of interest has been recently devoted to the possibility of *matter-enhanced* oscillations [4]. These are possible due to the effect of coherent neutrino scattering in the solar medium even when vacuum mixing angles which are very small. This resonant enhancement is possible, however, only if the neutrino mass difference lies in the range <sup>§</sup>

$$10^{-7} \lesssim \left( \frac{\delta m}{\text{eV}} \right)^2 \lesssim 10^{-4}. \quad (1)$$

Again the oscillation hypothesis indicates mass differences that are too small to be checked in existing experimental setups. Although effort will be pushed in this direction [11], it will be a number of years before one will be able to start probing the range specified in eq (1). From this point of view it is interesting to ask *What are the physics options for having an*

---

<sup>§</sup>Detailed analysis of the range of parameters required for the resonant amplification of neutrino oscillations to take place has been given by many authors, see *eg* ref [10].

*experimentally testable solution to the solar neutrino puzzle in terms of neutrino oscillations?*

and, in addition, *What is the physical origin for the relevant neutrino mass scale?*

The second question may be answered in a variety of ways *eg* by introducing in the theory heavy singlet fermions such as right handed neutrinos, at a mass scale *above* that of electroweak symmetry breaking. This could, for example, be the grand unification scale in SO(10) models [12] or some intermediate scale as present, *eg*, in some superstring inspired  $E_6$  models [13,14]. Alternatively, one may introduce a physical mass scale *below* the Fermi scale and have a see-saw mechanism in which the states relevant in determining the neutrino mass are at the Fermi scale [15]. A natural framework to implement this idea is supergravity, in which case the heavy states necessary in the see-saw mechanism are just the supersymmetric ( SUSY ) partners of gauge and Higgs fields. *In both cases we are still left with the first question.*

There are two generic types of models where the solar neutrino oscillation hypothesis may be experimentally checked. First it can be checked “kinematically” in models where  $m \gg \delta m$ , where  $m$  represents a typical neutrino mass and  $\delta m$  denotes a neutrino mass difference. One way to model this hierarchical difference in the values of  $m$  and  $\delta m$  is to attribute it [5] to the presence of a large intermediate scale in superstring inspired models [13]. The electron neutrino is just one Weyl component of a four-component Quasi-Dirac neutrino [16] of mass  $m \gg \delta m$ . The other component may be another neutrino flavour or, as in ref [5], can be a *sterile* neutrino. If  $\nu_e$  is a Quasi-Dirac particle it can be massive enough as to be detectable in tritium decay experiments and also to be relevant for cosmology, without conflicting any experimental limit. The solar neutrino problem can then be solved by large vacuum oscillations from the *active* to the *sterile* component as  $\nu_e$  propagates from sun to earth.

The alternative possibility is  $m = O(\delta m)$  so that “kinematical” neutrino mass effects will not be detectable but the theory contains some new “dynamical” degree of freedom whose “large” effects can be probed and used, in a sense, to “track” an otherwise undetectably small mass. The prototype of this situation is when B-L is a spontaneously broken symmetry of the Lagrangian. If ungauged, as in the standard model, spontaneous B-L breaking generates a Goldstone boson — a Majoron — which we denote  $J$ .<sup>¶</sup> The Majoron is a true dynamical degree of freedom so it has interactions which are related in a well defined way with the neutrino mass. Its emission generates new mechanisms of stellar energy loss. There are a variety of different Majoron models; here we concentrate on a variant of the idea where the Majoron is the SUSY partner of the neutrino [15]. In this case Majoron emission will also produce small changes in the decay parameters of the muon and the  $\tau$  lepton which could be seen in precision measurements. The present good agreement of the observations with the standard model predictions leads to nontrivial constraints. Interestingly enough, these constraints may allow the possible observation of rare decays such as  $\mu \rightarrow e + J$  which would provide an interesting signature for this scenario. The smallness of the neutrino mass, required by the resonance condition, eq (1), is dictated by constraints on the couplings of the Majoron that follow from a variety of considerations thus giving a dynamical basis for the see-saw mechanism.

## 2. The Spontaneously Broken R Parity (SBRP) Model

The model is described by the minimal supergravity superpotential

$$h_{ij}^u u_i^c Q_j H_u + h_{ij}^d d_i^c Q_j H_d + h_{ij}^e e_i^c \ell_j H_d + \mu H_u H_d \quad (2)$$

where the parameter  $\mu$  is related with electroweak breaking driven by radiative corrections

---

<sup>¶</sup>If the gauge group contains B-L there will be instead an additional gauge boson, coupled to neutrinos [17].

associated with the top quark. The first three terms give rise to masses for up and down-type quarks, and charged leptons, respectively, once the two Higgs fields  $H_u$  and  $H_d$  acquire their vacuum expectation values (vevs)  $v_u$  and  $v_d$ . In general supersymmetry and lepton flavour are broken explicitly in the scalar potential via soft scalar mass terms and possibly cubic scalar self couplings, and also via  $SU(3) \times SU(2) \times U(1)$  invariant gaugino mass terms  $M_i$ ,  $i=1,2,3$ . Gaugino masses break the continuous R invariance of the theory down to a discrete symmetry, called R parity. R parity is *even* for all particles of the standard model (including the Higgs scalars) and *odd* for their SUSY partners. R parity too may be broken, either explicitly [18] or spontaneously by nonzero vevs for the scalar neutrinos [19],

$$v_i = \langle \ell_i^0 \rangle; \quad i = e, \mu, \tau. \quad (3)$$

In the minimal model, spontaneous R parity breaking is very restrictive. Recent analysis [20] indicates that sleptons lighter than about 65 Gev and a top quark heavier than about 70 Gev are required. <sup>||</sup>

Spontaneous breaking of B-L, an ungauged continuous symmetry, generates a Nambu-Goldstone boson — a Majoron — given in ref [15], giving a neutrino mass

$$m = \frac{\mu M \sum_i v_i^2}{2v_u v_d M - M_1 M_2 \mu} \quad (4)$$

where  $2M = g_1^2 M_2 + g_2^2 M_1$  and  $g_i$  are gauge coupling constants. Note that since B-L is broken by *one* unit via the scalar neutrino vev, eq (3), it takes *two* such breakings to generate a (Majorana) mass for the (left handed) neutrino: hence the square in the see-saw formula, eq (4). This contrasts with the non-supersymmetric Majoron model [21] in which a scalar Higgs triplet is introduced to generate neutrino masses directly and therefore *linear* in the lepton number breaking expectation value.

---

<sup>||</sup>This breaking may be far easier to achieve if one adds, eg, terms that break total lepton number explicitly as well.

Another striking difference, which makes the SBRP model much more restrictive, is the fact that *one and only one* neutrino acquires mass, namely, the one which is related by supersymmetry to the Majoron. Its mass is given by eq (4).\*\* This simplifies considerably the structure of the CC weak interaction reducing the parameters relevant for describing the resonant neutrino oscillations in the sun to just *three* parameters: two mixing angles ( the third angle is not a physical parameter, due to the mass degeneracy between the two massless neutrinos ) and one neutrino mass parameter,  $m$ . Moreover, CP is conserved in the charged current. The matrix  $K$  describing the CC weak interaction, can be put in the canonical form [15,22]

$$K = \omega_{23}(\theta_{23})\omega_{13}(\theta_{13}) \quad (5)$$

where  $\omega_{ij}$  is a rotation by an angle  $\theta_{ij}$  in the  $ij$  plane.

For  $\nu_\tau \gg \nu_\mu \gg \nu_e$  the mixing angles in eq (5) are small so that the massive state is mostly  $\nu_\tau$  and resonant solar neutrino conversions occur from  $\nu_e$  into  $\nu_\tau$  [15]. Because of such drastic simplification, there are interesting experimental signatures that make the SBRP model testable. In [15] we showed how the gaugino-Higgsino mass spectrum is restricted by Davis' experimental results and how the allowed range of parameters where resonant amplification of solar neutrino oscillations can occur may be directly explored by searching for charged SUSY partners of gauge and Higgs particles, say, in electron-positron machines. Here we concentrate on low energy tests of the model based on the dynamics of the Majoron.

### 3. Constraints on Majoron Couplings

The Majoron couples to quarks through its Higgs doublet admixture via the first two terms in eq (2). These couplings are always flavour diagonal. The situation is different for

---

\*\*The full tree level neutral lepton mass matrix was given in ref [15] and radiative corrections are negligible for our purposes.

the charged leptons due to their mixing with charginos ( charged SUSY partners of gauge and Higgs bosons ), described by the mass matrix

$$\begin{array}{c|ccc}
& e_j^+ & \widetilde{H}_u^+ & \widetilde{W}^+ \\
\hline
e_i^- & h_{ij}v_d & 0 & g_2v_i \\
\widetilde{H}_d^- & h_{ji}v_i & \mu & g_2v_d \\
\widetilde{W}^- & 0 & g_2v_u & M_2
\end{array} \tag{6}$$

Since fermions of different weak isospin are mixed in eq (6) it follows that in our model the Majoron couplings to charged leptons are *not* flavour diagonal. *This is another important difference between the SBRP model and the triplet Majoron model and which as we will see, may be tested experimentally.* The coupling of the Majoron to physical charged leptons is described by the effective interaction Lagrangian,

$$i \frac{g_2^2}{\sqrt{2}} \frac{vm_j}{m_W^2} \left( \frac{1}{2} \delta_{ij} + fg_{ij} \right) \bar{e}_{Li} e_{Rj} J + \text{H.C.} \tag{7}$$

where the function  $f$  is given by

$$f = \frac{1+x^2}{(1-xy)^2} = M_W^2 \left( \frac{\cos^2 \phi}{\mu_1^2} + \frac{\sin^2 \phi}{\mu_2^2} \right) \tag{8}$$

$\phi$  is a chargino mixing angle and  $\mu_i$  are their masses.  $x$  and  $y$  denote, respectively, the Higgsino mixing parameter  $\mu$  and the supersymmetry breaking SU(2) gaugino mass parameter  $M_2$  in units of the  $W$  mass ( $x = \mu/m_W$ ,  $y = M_2/m_W$ ). For simplicity we took  $v_u \approx v_d$ . The coupling matrix  $g_{ij}$  is a projection matrix, given by

$$g_{ij} = \begin{pmatrix} s_{13}^2 & c_{13}s_{13}s_{23} & c_{13}s_{13}c_{23} \\ c_{13}s_{13}s_{23} & c_{13}^2s_{23}^2 & c_{13}^2c_{23}^2s_{23} \\ c_{13}s_{13}c_{23} & c_{13}^2c_{23}^2s_{23} & c_{13}^2c_{23}^2 \end{pmatrix} \tag{9}$$

The first term in eq (7) comes from the Majoron admixture in the  $H_d$  Higgs scalar while the second comes from lepton-chargino mixing.



In this model stellar energy loss proceeds via *single* Majoron emission in the Compton-like process  $\gamma + e \rightarrow e + J$ . From the cross section given in ref [23] and the coupling of eq (7) we have

$$v \left( \frac{1}{2} + f g_{ee} \right) \lesssim 30 \text{ KeV}. \quad (10)$$

In addition, *double* Majoron emission occurs in  $\gamma + e \rightarrow e + J + J$ . This process has normal gauge strength couplings but is mediated by the heavy charginos. We have calculated the total cross section for this process in the approximation of small photon energies  $E_\gamma \ll m_e$  and obtained the limit

$$f^2 g_{ee}^2 \lesssim 10. \quad (11)$$

The same couplings also induce, mediated by the heavy charginos, double Majoron emission processes  $e_j \rightarrow e_i + J + J$  thus changing the spectrum of  $e_j \rightarrow e_i + \nu + \nu$ . Requiring the fractional change in the Michel parameter to be within the accuracy of present experimental determination gives for the case of  $\mu \rightarrow e + J + J$  decay, the limit [24]

$$f^2 g_{e\mu}^2 \lesssim 10^{-2} \quad (12)$$

while for the corresponding  $\tau$  decays the limits are

$$f^2 g_{e\tau}^2 \lesssim .35 \quad (13)$$

and a similar bound on  $f g_{\mu\tau}$ . Due to the existence of another light scalar in the theory the limits could be up to a factor 2 stronger. Thus, from this point of view, precision measurements of  $\mu$  and  $\tau$  decays are theoretically very interesting.

Due to the couplings of eq (7) *single* Majoron emission in  $\mu$  and  $\tau$  decays can also occur. The branching ratio  $B_{ij}$  for  $e_j \rightarrow e_i + J$ , relative to  $e_j \rightarrow e_i + \nu + \nu$ , is given by

$$B_{ij} = 96\pi^2 \left( \frac{v}{m_j} \right)^2 f^2 g_{ij}^2. \quad (14)$$

For the case of  $\mu \rightarrow e + J$  decay the branching ratio may be as large as

$$B_{e\mu} \approx 3 \times 10^{-7} \times (v/30\text{Kev})^2 \quad (15)$$

thus suggestively close to the present experimental limit of TRIUMF  $B_{e\mu} < 2.6 \times 10^{-6}$  [25]. For  $\tau$  decays, however, the corresponding allowed branching ratios are far below what can be experimentally probed.

#### 4. Discussion

The supersymmetric Majoron is markedly different from the triplet Majoron [21] and also less likely to be excluded by accurate Z width measurement. In this model the tiny neutrino mass needed for the MSW effect [4,10] is accompanied by “large” dynamical effects associated with the existence of the Majoron thus providing a dynamical testing ground for the model. Majoron emission in  $\mu$  and  $\tau$  decays may be at the cutting edge of experimental test, for parameter values which solve the solar neutrino problem by the enhanced oscillation effect. If mixing angles  $s_{23}$  and  $s_{13}$  are small resonant solar neutrino conversions occur from  $\nu_e$  into  $\nu_\tau$  [15]. A reduction in solar neutrino flux below 2.6 SNU implies severe restrictions on the supersymmetric spectrum [15] and potentially large effects in  $\tau$  decays. As an example taking  $v \approx 30\text{Kev}$ ,  $s_{13} \approx s_{23} \approx .1$ , and  $f \approx 6$ , corresponding to lightest chargino mass values consistent with present PETRA limits, we get a deviation in the  $\tau$  decay Michel parameter comparable to present limits, while the change for the  $\mu$  decay is a factor three or so below present experimental limit. For the same parameters the BR for  $\mu \rightarrow e + J$  is  $\approx 3 \times 10^{-7}$ . A larger value for  $s_{23}$  would increase both the deviation in Michel parameter and the BR for single Majoron emission in  $\mu$  decay. These effects are likely to be larger by up to a factor two due to the existence of another light scalar in the theory. On the other hand, if we take a larger value for  $v$  the BR  $B_{e\mu}$  increases without changing the Michel parameters. Note

that for these parameter values, the lightest *chargino* weighs less than half of the  $Z$  mass and thus might be seen at accelerator experiments.

We are thankful to L. Wolfenstein and P. Rosen for the hospitality extended to one of us (J. V.) at Aspen, during the Workshop on Solar and Astrophysical Neutrinos, where part of this work was done. J. V. also thanks the hospitality of the LBL Theory Group where this work was completed, especially I. Hinchliffe and L. Hall for clarifying discussions. The work of A. S. was supported by a fellowship from CSIC.

## REFERENCES

1. G West, talk at XI International Workshop on Weak Interactions, Santa Fe, June 1987; G Gelmini, L Hall, M Lin, Nucl Phys B281 (1987) 726.
2. J N Bahcall, S T Petcov, S Toshev, J W F Valle, Phys Lett B181 (1986) 369
3. L Okun, M Voloshin, M Vytotsky, Sov.J.Nucl.Phys. 44 (1986) 440.
4. P Mikheyev, A Smirnov, Nuovo Cimento Soc Ital Fis 9C (1986) 17; H Bethe, Phys Rev Lett 56 (1986) 1305; L Wolfenstein, Phys Rev D17 (1978) 2369.
5. R Mohapatra, J W F Valle, Phys Lett B177 (1986) 47
6. S Glashow, L Krauss, Phys Lett B190 (1987) 199.
7. J W F Valle, Phys Lett B131 (1983) 87; G Gelmini, J W F Valle, Phys Lett B142 (1984) 181; G Gelmini, D Schramm, J W F Valle, Phys Lett B146 (1984) 311; J W F Valle, Phys Lett B159 (1985) 49; S Nussinov, Phys Lett B185 (1987) 171.
8. K Hirata et al Phys Rev Lett 58 (1987) 1490; R Bionta et al, Phys Rev Lett 58 (1987) 1494.
9. J Frieman, H Haber, K Freese, SLAC PUB 4261 (1987)
10. W Haxton, Phys Rev Lett 57 (1986) 1271; S P Rosen, J Gelb, Phys Rev D34 (1986) 969; V Barger et al Phys Rev D34 (1986) 980.
11. F Boehm, private communication.
12. R Johnson, S Ranfone, J Schechter, Phys Lett B179 (1986) 355; P Langacker et al, Nucl Phys B282 (1987) 589.

13. J W F Valle, Phys Lett B186 (1987) 73; R Mohapatra, J W F Valle, Phys Rev D34 (1986) 1642; J P Derendinger et al, Nucl Phys B267 (1986) 365; S Nandi, U Sarkar, Phys Rev Lett 55 (1986) 566; F del Aguila et al, Nucl Phys B272 (1986) 413.
14. J W F Valle in Proc. of International Symposium on Weak and Electromagnetic Interactions in Nuclei, Heidelberg, 1986, ed H Klapdor, Springer-Verlag, p 927.
15. A Santamaria, J W F Valle, Phys Lett B195 (1987) 423
16. J W F Valle, Phys Rev D27 (1983) 1672; J W F Valle, M Singer, Phys Rev D28 (1983) 540; S Petcov, Phys Lett B110 (1982) 245; L Wolfenstein, Nucl Phys B186 (1981) 147.
17. J W F Valle, Phys Lett B196 (1987) 157
18. L Hall, M Suzuki, Nucl Phys B231 (1984) 419; I H Lee, Nucl Phys B246 (1984) 120.
19. R Mohapatra, C Aulakh, Phys Lett B119 (1983) 136; G G Ross, J W F Valle, Phys Lett B151 (1985) 375; J Ellis, G Gelmini, C Jarlskog, G G Ross, J W F Valle, Phys Lett B150 (1985) 142; S Dawson, Nucl Phys B261 (1985) 297.
20. A Bouquet, P Salati B284 (1987) 557; B Gato, J Leon, J P Mercader, M Quiros, Nucl Phys B260 (1985) 203.
21. G Gelmini, M Roncadelli, Phys Lett B99 (1981) 411.
22. J Schechter, J W F Valle, Phys Rev D21 (1980) 309; Phys Rev D22 (1980) 2227.
23. H Georgi, S Glashow, S Nussinov, Nucl Phys B193 (1981) 297; J Ellis, K Olive, Nucl Phys B223 (1983) 252 and references therein.
24. G Kane, W Rolnick Nucl Phys B232 (1984) 21; W Buchmuller, F Scheck, Phys Lett B145 (1984) 421; A Santamaria, J Bernabeu, A Pich, Phys Rev D36 (1987) 1408.

25. A Jodidio et al, Phys Rev D34 (1986) 1967.