



A Neandertal dietary conundrum: Insights provided by tooth enamel Zn isotopes from Gabasa, Spain

Klervia Jaouen^{a,b,1}, Vanessa Villalba-Mouco^{c,d,e}, Geoff M. Smith^{b,f}, Manuel Trost^b, Jennifer Leichter^{g,h}, Tina Lüdecke^{g,i}, Pauline Méjean^a, Stéphanie Mandrou^a, Jérôme Chmeleff^a, Danaé Guiserix^{a,i}, Nicolas Bourgon^{b,h}, Fernanda Blascol, Jéssica Mendes Cardoso^{a,k}, Camille Duquenoy^a, Zineb Moubtahij^b, Domingo C. Salazar Garcia^{b,l,m,n}, Michael Richards^o, Thomas Tütken^h, Jean-Jacques Hublin^{b,p}, Pilar Utrilla^e, and Lourdes Montes^e

Edited by Suzanne Pilaar Birch, The University of Georgia, Athens, GA; received July 1, 2021; accepted April 14, 2022 by Editorial Board Member Dolores R. Piperno

The characterization of Neandertals' diets has mostly relied on nitrogen isotope analyses of bone and tooth collagen. However, few nitrogen isotope data have been recovered from bones or teeth from Iberia due to poor collagen preservation at Paleolithic sites in the region. Zinc isotopes have been shown to be a reliable method for reconstructing trophic levels in the absence of organic matter preservation. Here, we present the results of zinc (Zn), strontium (Sr), carbon (C), and oxygen (O) isotope and trace element ratio analysis measured in dental enamel on a Pleistocene food web in Gabasa, Spain, to characterize the diet and ecology of a Middle Paleolithic Neandertal individual. Based on the extremely low $\delta^{66}\text{Zn}$ value observed in the Neandertal's tooth enamel, our results support the interpretation of Neandertals as carnivores as already suggested by $\delta^{15}\text{N}$ isotope values of specimens from other regions. Further work could help identify if such isotopic peculiarities (lowest $\delta^{66}\text{Zn}$ and highest $\delta^{15}\text{N}$ of the food web) are due to a metabolic and/or dietary specificity of the Neandertals.

carnivory | Middle Paleolithic | zinc isotope ratios | Iberian Neandertals | hominin

Over the last 30 years, analyses of nitrogen isotopes in collagen ($\delta^{15}\text{N}_{\text{collagen}}$) have provided direct evidence for Neandertal diets across Europe and Asia. These studies all indicate a carnivorous (1–12), or at least a meat-heavy, diet for European Neandertals. However, one peculiarity of Neandertal $\delta^{15}\text{N}_{\text{collagen}}$ remains the subject of an ongoing debate. From the one Siberian and eight western European sites, where both Neandertal and associated fauna have been analyzed, nitrogen isotope ratios in Neandertal collagen are systematically higher than that of other carnivores (3, 6–8, 10, 11, 13, 14). An explanation for such elevated values could be the consumption of herbivores, such as mammoths, which themselves exhibit elevated $\delta^{15}\text{N}$ values due to the consumption of plants growing on arid soils (1, 2, 7). While mammoth remains are usually scarce at Neandertal fossil localities, they were nonetheless occasionally consumed, as suggested by remains with cut marks and other human butchery signatures (15). The absence of mammoth remains at Middle Paleolithic sites could be a result of 1) Neandertals chose to leave large bone elements at the kill site and transport other edible carcass products, mainly meat, back to the habitation site (15), or 2) mammoths were not frequently consumed, and the $\delta^{15}\text{N}$ peculiarity consequently reflects the consumption of other resources enriched in ^{15}N .

Alongside this $\delta^{15}\text{N}$ peculiarity, one major obstacle to our knowledge of Neandertals' subsistence patterns is that the preservation of organic matter limits the application of collagen-bound nitrogen isotope analysis to fossil specimens. Collagen degrades over time at a varying speed depending on climatic and environmental conditions (16). The oldest hominin specimen in which bone protein is preserved is that of Scladina (Belgium), which dates to 90,000 cal BP (calibrated years before the present) (17), but most studied specimens are younger than 50,000 cal BP (1–3, 6–8, 10–13, 18). Furthermore, these specimens are only from sites in northwestern and central Europe and Siberia, where climatic conditions favored collagen preservation. As a result, the variability of Neandertals' diet over time and between regions may not accurately be reflected by the currently available isotope data. In Iberia, where the latest surviving Neandertals have been discovered (19, 20), collagen was successfully extracted for only one site (21). Therefore, our knowledge of Iberian Neandertal diets mostly relies on zooarcheological and dental calculus data, which show some inconsistencies (21–25). For instance, similar to other western European sites, zooarcheological studies emphasize the consumption of terrestrial mammals and birds (21). In contrast, analysis of dental calculus for microremains and ancient DNA metagenomic analysis (26–28)

Significance

Neandertals' diets are a topic of continued debate, especially since their disappearance has been frequently attributed to their subsistence strategy. There is no clear consensus on how variable their diets were in time and space. Isotope studies have helped quantify meat consumption in Neandertals, but usually rely on nitrogen isotope analyses of collagen, a protein rarely preserved in samples older than 50 ka. Moreover, collagen extraction for isotope analyses is rarely successful in Iberian skeletal material. Here, we employ zinc isotope analysis of dental enamel of a Neandertal and associated fauna (Gabasa, Spain), which can be applied to contexts >50 ka. This proxy confirms a high level of carnivory in an Iberian Neandertal.

Author contributions: K.J., V.V.-M., T.T., J.-J.H., P.U., and L.M. designed research; K.J., V.V.-M., G.M.S., M.T., J.L., T.L., P.M., S.M., D.G., N.B., J.M.C., and Z.M. performed research; K.J., V.V.-M., M.T., J.L., J.C., F.B., and C.D. contributed new reagents/analytic tools; K.J., V.V.-M., G.M.S., J.L., T.L., F.B., C.D., T.T., and J.-J.H. analyzed data; and K.J., V.V.-M., G.M.S., J.L., T.L., P.M., D.G., N.B., D.C.S.G., M.R., T.T., J.-J.H., P.U., and L.M. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission. S.P.B. is a guest editor invited by the Editorial Board.

Copyright © 2022 the Author(s). Published by PNAS. This article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

¹To whom correspondence may be addressed. Email: klervia.jaouen@get.omp.eu.

This article contains supporting information online at <http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2109315119/-DCSupplemental>.

Published October 17, 2022.

highlight the frequent consumption of plants and mushrooms. Indeed, Weyrich et al. (26) even suggest that Neandertals at El Sidrón (Fig. 1) rarely consumed meat but often ate mushrooms, which would also result in elevated $\delta^{15}\text{N}$ values (29). The consumption of marine foods is also attested for coastal Neandertals, but its frequency cannot be truly assessed in the absence of isotope studies (21, 23–25, 30). Finally, cannibalism has been documented at two Iberian sites (El Sidrón and Zafaraya) (22, 31) (Fig. 1), though such practices appear limited and most likely occurred only during periods of nutritional stress (32). Therefore, it is challenging to confirm the homogeneity of Neandertals' diets across time and space, calling into question a direct link between their subsistence strategy and disappearance.

This study aims to investigate if the Zn isotope proxy could help elucidate the dietary behaviors of Neandertals and the source of their $\delta^{15}\text{N}$ peculiarity, specifically by studying a Late Pleistocene Iberian food web where the presence of mammoth has not been documented (33). The development of Zn isotope analysis ($^{66}\text{Zn}/^{64}\text{Zn}$, expressed as $\delta^{66}\text{Zn}$) has proven that trophic level information can be retrieved from mammalian tooth enamel ($\delta^{66}\text{Zn}_{\text{enamel}}$) (34, 35), including fossil samples from Pleistocene food webs where organic matter is typically not preserved (36, 37). Previous studies have demonstrated that $\delta^{66}\text{Zn}_{\text{enamel}}$ decreases by ca. 0.30 to 0.60 ‰ with each step in archeological and modern food webs (34–38) and that $\delta^{66}\text{Zn}$ values associated with breastfeeding are higher than postweaning-associated values (39). While the main source of variation of $\delta^{66}\text{Zn}_{\text{enamel}}$ values is diet, local geology can also likely influence the isotope ratio of a given animal (36, 39). To date, three modern assemblages from Koobi Fora (Kenya), Kruger Park, and the western Cape (South Africa) (40), a few animals from a historical site (Rennes, France) (41), and three Late Pleistocene sites (Tam Hay Marklot, Nam Lot, and Tam Pa Ling, Laos) (36, 37) are the only terrestrial food webs for which Zn isotope data in teeth and/or bones have been published (SI Appendix, Fig. S14). In the modern Koobi Fora savannah food web, $\delta^{66}\text{Zn}_{\text{enamel}}$ differences have been observed between browsers and grazers (35), but this pattern was not seen in any of the three Pleistocene Asian forest food webs (36, 37). Among modern and historical human populations,

historically documented diets relying on plants are associated with higher $\delta^{66}\text{Zn}$ values than those that include a substantial quantity of animal products (41–44). Zinc isotopes of ancient hominins have been analyzed only in one Pleistocene *Homo sapiens* individual (37) from Southeast Asia, but not yet in any Neandertal specimen.

This current study contributes significantly to our understanding of Iberian Neandertal diets by providing information on their trophic ecology for a region where traditional nitrogen isotope analyses are unfeasible due to the poor preservation of organic matter. We use the Zn isotopic tool as well as other mobility, ecological, and dietary proxies applied on tooth enamel from hominin and animal remains from the cave site Cueva de los Moros 1 (Gabasa, Pyrenees, Spain; Fig. 1). The site, excavated in the 1980s, is very well documented [for stratigraphic context, see Montes and Utrilla (45) and SI Appendix, Section S1]. All remains come from layers e, f, and g of a single stratigraphic layer directly above layer h dated to 143 ± 43 ka. Numerous carnivore remains were recovered along with Neandertal remains (layers e and f), allowing for comparison of the different meat-eating taxa. Coexisting herbivores from three different types of environmental contexts are homogeneously represented in layers e, f, and g: mountains (Iberian ibex [*Capra pyrenaica*], chamois [*Rupicapra rupicapra*]), forest (cervids including red deer [*Cervus elaphus*]), and open environments (horses [*Equus ferus*], European wild asses [*Equus hydruntinus*]).

Sample Collection

Species, Sample, and Proxy Selection. We analyzed the tooth enamel of 65 samples belonging to 43 different teeth of 12 taxa for $\delta^{66}\text{Zn}_{\text{enamel}}$ (Fig. 2 and SI Appendix, Table S15), 42 samples belonging to 39 different teeth for $\delta^{13}\text{C}_{\text{enamel}}$ and $\delta^{18}\text{O}_{\text{enamel}}$ (SI Appendix, Table S19), 23 samples belonging to 18 different teeth for $^{87}\text{Sr}/^{86}\text{Sr}$ (SI Appendix, Table S7), and 40 samples for trace element ratios (SI Appendix, Table S11). We sampled 12 to 50 mg of tooth enamel (1 to 20 mg for Zn isotopes) of five carnivore taxa (lynx [*Lynx spelaea*], wolf [*Canis lupus*], fox [*Vulpes vulpes*], dhole [*Cuon alpinus*], cave hyena [*Crocuta spelaea*]), one omnivore (cave bear [*Ursus spelaeus*]), and at least




Fig. 1. (A) Location of the Gabasa site as well as other Neandertal sites mentioned in the text. (B) Detailed map of the Gabasa region. San Esteban de Litera and Benabarre are nearby modern cities.

six herbivore species (Iberian ibex [*C. pyrenaica*], chamois [*R. rupicapra*], red deer [*C. elaphus*], rabbit [*Oryctolagus cuniculus*], horse [*E. ferus*], European wild ass [*E. hydruntinus*]), as well as one of the four Neandertal dental remains recovered from the site. Hyenas (*C. spelaea*) are treated as a separate dietary group (bone-eating carnivores) based on the results of the Koobi Fora and Nam Lot food webs (35, 37).

Sampling Strategy Related to Breastfeeding. The single sampled Neandertal tooth is a first molar (M1; Ga1.Rev.303) (46). Previous studies based on tooth wear and trace elements suggest that Neandertals were likely weaned between the ages of 1 and 2.5 y old (47–50), thus possibly overlapping with the crown's formation period for this tooth. To minimize destructive analyses, we sampled an enamel fragment from the lower part of the crown, next to the cervical margin. This part of the tooth probably formed after weaning or at least when maternal milk consumption could have been strongly reduced (51). The specimen was heavily worn, making it difficult to evaluate the enamel formation period of the part that we sampled. Still, we were able to evaluate trophic spacing between species while taking into account the influence of maternal milk consumption, which triggers higher $\delta^{66}\text{Zn}$ values in human M1 (39). To do so, we analyzed $\delta^{66}\text{Zn}_{\text{enamel}}$ on different types of nonhominin teeth (dp4, C, P2, P3, P4, M1, M2, and M3; *SI Appendix, Table S4*) formed at different times during ontogeny and obtained values associated with breastfeeding and/or solid food consumption for all species analyzed from Gabasa. In addition, we analyzed Sr/Ca and Ba/Ca ratios that also show variation between tooth enamel mineralized in utero and during pre- or postweaning periods (52).

Sampling Strategy Related to Neandertal Mobility and Ecology. Other factors such as diagenesis, geographical origin, environmental context, or metabolism can be a source of Zn isotope variability (35, 36, 39, 53). In recent studies, linear mixing models highlighted a possible link between Zn and Sr isotope ratios in teeth, which has been interpreted as resulting from the influence of the local geology on the Zn isotope ratios of its associated food webs (36, 37, 39). To control for this potential bias, we analyzed strontium isotope ratios of Gabasa tooth enamel ($^{87}\text{Sr}/^{86}\text{Sr}_{\text{enamel}}$) for samples with sufficient material (*SI Appendix, Section S5*). We also collected Sr isotope data from rocks, soils, plants, and teeth to assess the regional bedrock isotopic data ($^{87}\text{Sr}/^{86}\text{Sr}_{\text{bedrock}}$) and identify whether some animals could have been derived from areas proximal to Gabasa with different geology (*SI Appendix, Table S18*). We hypothesize that the Neandertal consumed local food sources, whereby its $^{87}\text{Sr}/^{86}\text{Sr}$ should fall within the ranges observed both for the local bedrock and assumed sympatric animals.

We estimated carbon and oxygen isotope compositions ($\delta^{13}\text{C}_{\text{enamel}}$ and $\delta^{18}\text{O}_{\text{enamel}}$, respectively) of the Neandertal and associated fauna. Notably, $\delta^{13}\text{C}_{\text{enamel}}$ and $\delta^{18}\text{O}_{\text{enamel}}$ values can offer insights into ecological proxies, such as the environment from which the Neandertal individual obtained its prey (i.e., open arid vs. closed mesic habitats), and possibly reveal information about which types of prey were consumed (54–56). Moreover, carbon and oxygen isotope data allow us to compare the food web of Gabasa with other food webs where Neandertals have been discovered. For one horse tooth, five samples were taken serially along the growth axis of the tooth to evaluate potential seasonality and intratooth variability effects over time on Sr and Zn isotope ratios.

Sampling Strategy to Assess the Degree of Chemical Tooth Alteration. Although it has been shown that dental enamel preserves biogenic Zn isotope signatures very well in Pleistocene or even older samples (36, 57), we independently confirmed that Zn isotope variability is not related to diagenetic processes. For this purpose, we analyzed trace elements that are abundant in soils but not in tooth enamel (Fe, Al, Mn) and hence are indicative of postmortem trace element uptake (*SI Appendix, Section S3*). Additionally, we examined the carbonate content of dental enamel to eliminate samples with obvious diagenetic alteration (>10 wt% CO_3) (*SI Appendix, Section S3*). To capture the influence of diagenesis on isotope ratios, we also performed measurements on dentin samples from two wolves for Sr, C, and O isotopes and trace elements and one wild ass for Zn isotopes. Finally, four roots of teeth which enamel was sampled (one chamois, two foxes, and one lynx) were used to attempt collagen extraction (*SI Appendix, Tables S9 and S10*), using a high extraction–yield protocol adapted for radiocarbon dating (58, 59) in the hope of getting additional trophic level information and possibly permitting high-precision radiocarbon dating.

Results

Datasets, methods, and additional discussions are given in detail in *SI Appendix*.

Zinc Isotope Data. Zinc isotope values of Gabasa mammalian tooth enamel range from 0.35 to 1.63 ‰ ($n = 65$) and are mainly influenced by the dietary group of the animals (Figs. 2 and 3) (Kruskal–Wallis $\chi^2 = 19.496$, $P = 0.0002159$ and $\chi^2 = 18.298$, $P = 0.0003818$; samples with elevated Al, Mn, and Fe content are excluded), which are assigned based on those used in other $\delta^{66}\text{Zn}_{\text{enamel}}$ isotope studies and zooarcheological data (dietary groups: carnivores, bone-eating carnivores, omnivores, and herbivores) (35, 60, 61). Animal values from different archeological layers fall in the same range (*SI Appendix, Section S4*). As previously observed for modern, historical, and Late Pleistocene tooth enamel, there is no correlation between Zn isotope values and Zn concentration data (35, 36, 39, 41, 42). This is an excellent indicator to assess the preservation of biogenic Zn isotope signatures (62). On average, large herbivores have higher Zn isotope ratios ($\delta^{66}\text{Zn}_{\text{enamel}} = 1.24 \pm 0.17$ ‰ 1σ ; $n = 16$) than carnivores ($\delta^{66}\text{Zn}_{\text{enamel}} = 0.92 \pm 0.18$ ‰ 1σ ; $n = 16$; 0.85 ± 0.14 ‰ if hyenas and the Neandertal are excluded). The Neandertal tooth exhibits the lowest $\delta^{66}\text{Zn}_{\text{enamel}}$ value (0.35 ± 0.00 ‰), substantially lower ($\Delta^{66}\text{Zn}_{\text{Neandertal-carnivores}} = 0.57$ ‰) than that of the lowest carnivore's value. Mirroring patterns in the modern Koobi Fora (East Africa) food web, the difference between herbivore and carnivore values is even more pronounced when hyenas are excluded, as bone consumption is thought to induce higher $\delta^{66}\text{Zn}$ values (35). The omnivorous cave bears ($\delta^{66}\text{Zn}_{\text{enamel}} = 1.15 \pm 0.13$ ‰ 1σ ; $n = 3$) exhibit Zn isotope ratios closer to those of large herbivores and rabbits rather than the carnivores', similar to a pattern often observed in nitrogen isotopes (63–66). The herbivores show the highest $\delta^{66}\text{Zn}$ values (0.96 to 1.60 ‰; $n_{\text{measurements}} = 40$ from $n_{\text{tooth}} = 27$ teeth) with grazers (European wild asses and horses, $\delta^{66}\text{Zn} = 1.35 \pm 0.18$ ‰ 1σ ; $n_{\text{tooth}} = 6$) exhibiting the highest values, another similarity to observations in modern food webs (35, 40).

Strontium Isotope Data. The $^{87}\text{Sr}/^{86}\text{Sr}_{\text{enamel}}$ range from 0.70834 to 0.70916 ($n = 24$). Strontium and Zn isotope ratios show no correlation, and the Neandertal individual shows an $^{87}\text{Sr}/^{86}\text{Sr}_{\text{enamel}}$ overlapping with those of deer and rabbits (Fig. 4A). Strontium




Fig. 2. Enamel $\delta^{66}\text{Zn}$ values per mammalian taxon for different dietary groups from Gabasa, Spain given as boxplots. Note that subsamples taken from one single tooth reflect differences in dietary ^{66}Zn values. See text for detailed discussion about the low Neandertal $\delta^{66}\text{Zn}$ value.

isotope ratios from dental enamel and dentin are compatible with the values expected for the region of Gabasa (*SI Appendix, Sections S1 and S5*). As dentin is likely to be affected by diagenetic alteration and Sr exchange with the soil, these values might reflect soil values and can thus be an indicator of the local bioavailable Sr isotope composition at Gabasa (*SI Appendix, Section S3*).

Carbon and Oxygen Isotope Data. $\delta^{13}\text{C}_{\text{enamel}}$ data range from -14.13 to -8.35 ‰ ($n = 44$), and $\delta^{18}\text{O}_{\text{enamel}}$ data range from -7.46 to 1.64 ‰ ($n = 43$). Mountain herbivores (chamois and Iberian ibex) have distinctly higher $\delta^{13}\text{C}_{\text{enamel}}$ and $\delta^{18}\text{O}_{\text{enamel}}$ compared with other herbivores, which might be due to water evaporation on mountain slopes and consumption of plants from more arid contexts (Fig. 3 and *SI Appendix, Section S6*). The Neandertal's $\delta^{13}\text{C}_{\text{enamel}}$ and $\delta^{18}\text{O}_{\text{enamel}}$ values overlap with those of cave animals (Fig. 4 A and B and *SI Appendix, Fig. S12*). Five carbon ($\delta^{13}\text{C}$) and oxygen ($\delta^{18}\text{O}$) values were possibly impacted by diagenetic carbonate, and were excluded for ecological interpretation (mostly rabbits; *SI Appendix, Sections S3 and S6*).

Trace Element Ratios (Ba/Ca and Sr/Ca). Trace element ratios are mostly discussed in *SI Appendix, Sections S3 and S7*. The $-\log(\text{Ba}/\text{Ca})$ ranges from 3.25 to 4.75 ($n = 40$) and $-\log(\text{Sr}/\text{Ca})$ from 2.77 to 3.62 ($n = 40$). The two ratios measured in dental enamel do not correlate. As expected (67), mixed feeders exhibit lower Sr/Ca than grazers, but carnivores (especially hyenas) show Sr/Ca ratios overlapping with those of grazers (*SI Appendix, Fig. S13*). We report the highest Sr/Ca and Ba/Ca ratios for mountain and forest animals and the lowest Ba/Ca for the Neandertal individual. Here, these ratios do not correlate with tooth developmental stages (i.e., ontogenetic time of crown mineralization).

Correlations and Comparisons between Different Dietary Proxies. Zn isotopes do not correlate with Sr/Ca or Ba/Ca ratios (Fig. 4 and *SI Appendix, Sections S3 and S4*). Equids have higher $\delta^{66}\text{Zn}$, $\delta^{13}\text{C}$, and $^{87}\text{Sr}/^{86}\text{Sr}$ and lower Ba/Ca than the rest of the herbivores (Fig. 4). When performing a principal-component analysis (PCA) based on five proxies ($\delta^{66}\text{Zn}$, $\delta^{13}\text{C}$, $\delta^{18}\text{O}$, Sr/Ca, and Ba/Ca), the PC1 places the Neandertal among cave animals (hyenas and bears). This first component is mostly based on $\delta^{13}\text{C}$, $\delta^{18}\text{O}$, and Ba/Ca values. Strangely, it is precisely when cave animals are removed that a positive correlation appears between $\delta^{13}\text{C}$ and $\delta^{66}\text{Zn}$ ($R^2 = 0.41$, $P = 0.00009$). This correlation could be due to a combination of

a trophic level effect and higher $\delta^{13}\text{C}$ and $\delta^{66}\text{Zn}$ values among grazers. No clear trend has been observed between $\delta^{66}\text{Zn}$ values and the type of tooth sampled (*SI Appendix, Section S4*), but it should be noted that this could result from the low sample size per tooth group and different timing of weaning and dental development between species.

Collagen Extraction. The collagen extraction performed on the dentin from four teeth was unsuccessful, and not enough collagen was extracted for $\delta^{15}\text{N}_{\text{collagen}}$ and/or radiocarbon analyses. This failure is not surprising for samples dating back to about 150 ka (62) (*SI Appendix, Section S2*).

Discussion

Insights into the Diet and Ecology of Iberian Pleistocene Mammals from Zn Isotopes in Tooth Enamel. Our study shows similar trends to those previously reported in modern and fossil food webs: Herbivores exhibit higher $\delta^{66}\text{Zn}$ than carnivores ($\Delta^{66}\text{Zn}_{\text{herbivores-carnivores}} = +0.31$ ‰ and $+0.38$ ‰ excluding hyenas). The trophic level depletion is similar to that of the modern food web of Koobi Fora, Kenya (dental enamel values: $+0.40$ ‰ hyenas included, and $+0.44$ ‰ excluding hyenas), or the marine mammal food webs of Arctic Canada ($+0.32$ ‰ on average in bones) (34, 38) but lower than that of the Laotian fossil sites ($+0.63$ ‰ hyenas excluded). Grazers (horses and European wild asses) tend to show higher $\delta^{66}\text{Zn}$ than browsers, as observed in the savannah of Koobi Fora (35) but not in the tropical rain forests of Tam Hay Marklot and Nam Lot (36, 37). Hyenas have more elevated $\delta^{66}\text{Zn}_{\text{enamel}}$ values than other carnivores (Fig. 2), as systematically observed in other sites (35, 37, 40). These higher values are likely due to bone consumption (35), although hyenas have more extended breastfeeding time than other carnivores (68) and milk would also similarly drive $\delta^{66}\text{Zn}_{\text{enamel}}$ up. However, high Sr/Ca and low Ba/Ca ratios observed in the different teeth of all hyenas support the interpretation of the signature of solid food consumption and consequently of bones as a likely driving factor behind their elevated $\delta^{66}\text{Zn}_{\text{enamel}}$ compared with other sympatric carnivores (*SI Appendix, Fig. S13 and Table S4*). Cave bears exhibit $\delta^{66}\text{Zn}_{\text{enamel}}$ overlapping with rabbits, deer, and chamois (Fig. 2), which confirms the frequent observations based on low $\delta^{15}\text{N}_{\text{collagen}}$ indicating they were mostly herbivorous (12, 63, 66).




Fig. 3. Average dental enamel isotope and trace element ratios for the different dietary categories at Gabasa, Spain. (A) $\delta^{66}\text{Zn}_{\text{enamel}}$. (B) $^{87}\text{Sr}/^{86}\text{Sr}$. (C) $\delta^{13}\text{C}_{\text{enamel}}$. (D) $\delta^{18}\text{O}_{\text{enamel}}$. (E) $\log(\text{Ba}/\text{Ca})$. Omnivores correspond to one single taxon (cave bears) as well as carnivores with significant bone consumption (cave hyenas).

The $\delta^{66}\text{Zn}_{\text{enamel}}$ values of the entire food web of Gabasa are the most elevated values ever measured in dental enamel [mean values about 0.2 ‰ higher than in the modern food web of Koobi Fora and 0.7 ‰ higher than in the Late Pleistocene food webs of Marklot, Nam Lot, and Tam Pa Ling (35, 36, 41, 42); *SI Appendix, Section S4 and Fig. S14*]. This is probably not due to the geology because limestones, known to show the highest $\delta^{66}\text{Zn}$ (69), are present at both Gabasa and the Laotian sites. The Gabasa food web's uniqueness comes from its environmental context as it is the coldest terrestrial biotope ever studied for faunal $\delta^{66}\text{Zn}$. Pollen data suggest an arid and cold climate (70) with periods characterized by steppe vegetation and some forest patches mainly composed of conifers (45, 70, 71). Carbon and oxygen isotopes exhibit patterns similar to those observed for mammals from various Middle

Paleolithic western European sites (54, 55, 72) and are compatible with midmountain mosaic landscape (*SI Appendix, Sections S1 and S6*). Tree leaves usually exhibit relatively low $\delta^{66}\text{Zn}$ and were probably scarce in Gabasa, as opposed to the forest of Tam Hay Marklot and the fynbos of the western Cape (40), where the lowest $\delta^{66}\text{Zn}$ in herbivore teeth was measured.

The Diet of a Neandertal at Gabasa. Our results demonstrate that the Neandertal individual from Gabasa shows a Zn isotope signature of a top-level carnivore, similar to that observed for nitrogen isotopes for other sites with Neandertal occupation. Carbon and oxygen isotope and trace element data suggest that the individual inhabited the local area around the cave (*SI Appendix, Sections S6 and S7*). Of all the animal taxa analyzed in Gabasa, the Neandertal specimen easily exhibits the




Fig. 4. Zinc isotope values ($\delta^{66}\text{Zn}$, average dental enamel values per tooth) cross-plotted versus (A) carbon ($\delta^{13}\text{C}$), (B) oxygen ($\delta^{18}\text{O}$), and (C) strontium ($^{87}\text{Sr}/^{86}\text{Sr}$) isotope composition as well as (D) Ba/Ca ratios measured in faunal and Neandertal tooth enamel at Gabasa, Spain. Each point corresponds to a single tooth, except for the horse tooth, for which five samples were taken (samples are connected with a line). Error bars correspond to the SD between analytical duplicates for C and O isotopes (0.02 to 0.36 ‰; one sample with higher SD has been removed) and for Zn (analytical duplicates: 0.00 to 0.04 ‰; SD for separate samples taken from a single tooth: 0.04 to 0.21 ‰). SEs for strontium isotope ratios are 0.000003 to 0.000010 and are too small to display.

lowest Zn isotope ratio. As the Sr isotope ratio of the Neandertal tooth enamel is similar to those of other animals (Fig. 4), the low $\delta^{66}\text{Zn}_{\text{enamel}}$ value is unlikely the result of a different geographic origin (and consequently different bedrock and isotope baseline) of this individual but reflects diet instead. Previous work on animal food webs documented a trophic spacing for $\delta^{66}\text{Zn}_{\text{enamel}}$ of 0.3 to 0.6 ‰ (34–36, 38). In Gabasa, the average difference between carnivores and herbivores is 0.31 ‰, but the Neandertal shows a $\delta^{66}\text{Zn}$ value 0.57 ‰ lower than that of the carnivores—0.50 ‰ if hyenas are excluded—and 0.85 ‰ lower than the herbivores, which would be about three trophic levels higher than that of the herbivores if we consider a trophic spacing of 0.31 ‰. This mirrors dietary reconstructions based on nitrogen isotopes from individuals from other European sites such as Goyet (18) (Belgium), Jonzac (13) (France), or Les Cottés (7) (France), where Neandertals exhibit higher $\delta^{15}\text{N}_{\text{collagen}}$ values than sympatric carnivores.

In order to explain the low $\delta^{66}\text{Zn}_{\text{enamel}}$ value of the Neandertal tooth, we here consider 1) a signature of breastfeeding, 2) specific diets (the consumption of either light Zn-bearing foods or the absence of consumption of heavy Zn-bearing foods), and 3) a metabolic origin.

The breastfeeding signature hypothesis can be ruled out because it leads to high $\delta^{66}\text{Zn}_{\text{enamel}}$. In a previous study, teeth formed during the breastfeeding period of two human populations from very different geographical and archeological contexts exhibit average $\delta^{66}\text{Zn}_{\text{enamel}}$ values that are 0.3 ‰ higher than the teeth whose dental enamel forms after weaning (39).

Furthermore, the low Ba/Ca of the Neandertal dental enamel suggests that it was formed after the breastfeeding period (49, 52, 73) (SI Appendix, Table S20). Moreover, the region of the tooth sampled was very close to the tooth's cervical margin because of tooth wear, which usually corresponds to formation ages of 2 to 3 y old (74). The turnover of Zn in the body occurs over the course of just a few months, and weaning age (or at least an age with substantial consumption of solid food) among Neandertals is assumed to be around 1 y old (49). Even if milk consumption extended to the period from which enamel was sampled from the tooth, it is unlikely that it was a major source of Zn in the diet at this stage. Therefore, the value observed in the Neandertal tooth enamel is likely a reflection of an adult diet. An early weaning age also prevails in recent humans and has also been documented with Zn isotopes in a Late Pleistocene *H. sapiens* (37), strongly contrasting with the conditions observed among apes, *Australopithecus* (75), and earlier representatives of the genus *Homo* (76). This adaptation results from the necessity for the breastfeeding mother to share the energetic burden of her child's large brain with other adults able to contribute to solid food. It is central to the human pattern of “cooperative breeding” and has critical social and behavioral implications.

Some dietary explanations for the low $\delta^{66}\text{Zn}_{\text{enamel}}$ of the Neandertal individual involving light Zn-bearing foods are also excluded:

- 1) Adult vs. juvenile prey: In principle, if [as in humans (39)] milk consumption leads to higher $\delta^{66}\text{Zn}$, preferential

consumption of postweaning juveniles or adults might lead to lower $\delta^{66}\text{Zn}$. However, this explanation seems unlikely for two reasons. First, such an isotopic disparity has not been observed in herbivore teeth in relationship to weaning in this study or elsewhere (36), so the age of the prey does not appear to matter. Second, Neandertals targeted younger deer and horses (61) (*SI Appendix, Section S1*) while other carnivores targeted adult ibexes. As such, if prey age mattered, Neandertal Zn isotope compositions should be biased toward higher $\delta^{66}\text{Zn}$, not lower. Although we cannot rule out the possibility that the formation period of the Neandertal tooth is asynchronous with the site's period of occupancy, there is currently no evidence to support this.

- 2) Cannibalism or hypercarnivorism: Cannibalism or carnivore meat consumption could be associated with low Zn isotope ratios, but the absence of cut marks on hominin and carnivore bones rules out this hypothesis.
- 3) Food with unusually low $\delta^{66}\text{Zn}$ values: Archeological evidence for some isotopically light foods that Neandertals might have specifically targeted may be missing [e.g., animal livers (53, 77, 78), birds, mushrooms, fruits, and leaves (79–81), insects (82, 83), or aquatic resources (43)]. However, $\delta^{66}\text{Zn}_{\text{enamel}}$ of the consumers of those foods has been measured and is not associated with lower values than that of sympatric carnivores, both in Gabasa (e.g., lynx are known to consume liver, deer consume leaves, while foxes, bears, and rabbits have been known to consume mushrooms) and other food webs [apes and tapirs for insects and fruits (36)].
- 4) Aquatic foods: The consumption of aquatic resources has been demonstrated for some Iberian Neandertals, though not at Gabasa (23, 25, 30). Still, the consumption of common trout (*Salmo trutta*) has been suggested for a Mousterian site at the Grotte du Noisetier (84), located in the French Pyrenees (Fig. 1). The trophic level of this species could potentially explain low $\delta^{66}\text{Zn}_{\text{enamel}}$ values, but would imply that the main source of Zn in the diet comes from freshwater fish. Trout Zn content is about 10 times less than that of liver or muscle from mammals (85), requiring Neandertals to have consumed trout almost exclusively to shift their $\delta^{66}\text{Zn}_{\text{enamel}}$ to the observed low value. As no fishbone or other evidence for aquatic resource use was found in the Gabasa region, we can disregard this hypothesis as unlikely.
- 5) Food processing: Elevated $\delta^{15}\text{N}$ values among Neandertals have been suggested to result from food processing such as fermentation or cooking. However, Zn fractionates only at a temperature above $\sim 900^\circ\text{C}$ (86), which is not reached in open fires, and even if fractionation did occur, it would enrich the Neandertal food in heavy Zn isotopes (86). Biotic fractionation during meat processing is not documented, but could happen within muscle tissues if the reactions were incomplete during fermentation (87) or other curing processes. However, this would only influence the final $\delta^{66}\text{Zn}$ of the consumed food if both reagents and products of the curing process would not be contained in the final food product.

A combination of dietary practices might explain the low $\delta^{66}\text{Zn}_{\text{enamel}}$ value of the Neandertal. Zinc isotopes are highly fractionated in mammalian tissues (77, 78, 88, 89), and different mammalian species have different isotope compositions. Thus, a low $\delta^{66}\text{Zn}$ value might reflect consumption of muscle and liver (low $\delta^{66}\text{Zn}$) from deer and rabbits (lower $\delta^{66}\text{Zn}$ than other herbivores; Fig. 3) (also supported by $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ for which Neandertals exhibit values close to these animals; Figs. 3 and 4 and *SI Appendix, Fig. S10*), while excluding bones and

blood (high $\delta^{66}\text{Zn}$). Although cut marks suggest Neandertals hunted deer and horses (61, 90), rabbits are the most abundant taxon of the site (*SI Appendix, Fig. S2*), and some rabbit bones show cut marks (*SI Appendix, Fig. S3*). Disarticulation of rabbits requires minimal use of tools (61), so they could have been heavily consumed without physical evidence. Temporal asynchrony between cave deposits and tooth growth might explain a discrepancy between zooarcheological and isotopic dietary conclusions if the deposits represent a period when Neandertals preferentially ate horses and deer, whereas the portion of the tooth corresponds to a period when rabbits were preferentially eaten. Higher $\delta^{66}\text{Zn}$ among other sympatric carnivores hunting the same species as Neandertal (*SI Appendix, Table S3*) (61, 91) might partially reflect bone consumption: Hyenas tend to have elevated $\delta^{66}\text{Zn}_{\text{enamel}}$ values, likely due to bone consumption (35, 37). Red foxes and wolves gnaw on bones [although they ingest much less bone than hyenas (92)], while lynxes partially digest the bones of rabbits, which make up 80 to 100% of their typical diets (91). In addition, all these carnivores may consume substantial blood, possibly helping to explain higher $\delta^{66}\text{Zn}$ values than the Neandertal.

Metabolic specificity (different diet–tissue isotopic offsets) for Neandertals compared with other animals might also explain the unusually low $\delta^{66}\text{Zn}$ value and possibly the higher $\delta^{15}\text{N}_{\text{collagen}}$ values that have been measured elsewhere. It is indeed striking how similarly the patterns between $\delta^{66}\text{Zn}_{\text{enamel}}$ and $\delta^{15}\text{N}_{\text{collagen}}$ mirror each other: In both proxies, Neandertals typically exhibit isotope ratios that would appear to imply a higher trophic level than sympatric carnivores (1, 3–5, 7, 11, 63). Although such distinct compositions have been interpreted to reflect aquatic food resources for some Paleolithic humans (11), compound-specific isotope analysis of amino acids shows that anatomically modern humans from Buran Kaya (Crimea) relied on terrestrial resources, even though they exhibit the highest $\delta^{15}\text{N}_{\text{collagen}}$ known for the Paleolithic (93). These authors suggest preferential consumption of mammoths (absent from Gabasa), but a metabolic origin might yield similarly high $\delta^{15}\text{N}_{\text{collagen}}$ values. We acknowledge that some studies show minimal offsets in $\delta^{15}\text{N}_{\text{collagen}}$ between humans and sympatric herbivores (94). However, the trophic spacing between the Paleolithic and Neolithic periods did not decrease to the extent expected for a transition from hunter-gatherer to a more cereal-based diet (95). Furthermore, for Zn isotopes, a medieval population with expected high meat and marine fish consumption shows strikingly lower $\delta^{66}\text{Zn}_{\text{enamel}}$ values than for sympatric herbivores (0.6 ‰ lower) and for a dog, cat, and pig (0.3 ‰ lower), while $\delta^{15}\text{N}_{\text{collagen}}$ values were strikingly higher than those of the herbivores (4 ‰) and dog, cat, and pig (1 ‰) (41). These observations closely resemble patterns for Neandertals at Gabasa (for Zn isotopes) and elsewhere (for N isotopes), and suggest a metabolic effect. Controlled feeding experiments on animals with an omnivorous diet are necessary to assess this further. The reason for this metabolic difference remains unknown, but it should be noted that the Neandertal is the most carnivorous species among primates, and the only meat eater not from the order *Carnivora* within the Gabasa faunal assemblage. Therefore, Neandertals may have had some metabolic peculiarities compared with other carnivores and primates regarding adaptations of their gastrointestinal tract and their dietary preferences (96).

Concluding Remarks

Zinc isotope analysis of tooth enamel can successfully characterize past animal and human diets in the absence of collagen

preservation for $\delta^{15}\text{N}$ analysis. Whichever method is used to characterize Neandertals' diets ($\delta^{66}\text{Zn}_{\text{enamel}}$ and $\delta^{15}\text{N}_{\text{collagen}}$), isotope values represent an extreme in carnivore trophic levels, and Iberian Neandertals appear to be no exception. Meat consumption is supported by both the zooarcheological data at Gabasa and $\delta^{66}\text{Zn}_{\text{enamel}}$ analysis. Furthermore, the low Zn isotope ratio observed in the single measured Gabasa Neandertal specimen suggests that this individual might have had a distinct diet compared with other carnivores (possibly avoiding the consumption of bones and blood) and/or may not have fractionated Zn isotopes like other sympatric carnivores. Our successful analysis of Zn isotopes from a Pleistocene hominin-tooth paves the road for a refined understanding of Neandertal diet. Further work should focus on understanding whether such unusual compositions result from specific diets or metabolic fractionation in these extinct hominins.

Materials and Methods

Materials. Details of the context of Gabasa cave deposits, geology, dating, zooarcheology, paleoenvironment, and sampling strategy are available in *SI Appendix*. The material used in this study is associated with the permit "Resolution" by the Gobierno de Aragon on 10 February 2016 and 31 August 2018.

Methods. Tooth enamel fragments and powders were sampled at the Department of Human Evolution of the Max Planck Institute for Evolutionary Anthropology (MPI-EVA) using a drill with a diamond saw and at the Géosciences Environnement Toulouse of the Observatoire Midi Pyrénées (GET/OMP) using a MicroMill. For Zn isotope analyses, 1- to 20-mg enamel fragments were dissolved in 1 mL 1 N HCl. Samples were cleaned with double-distilled H_2O in an ultrasonic bath and dried at 50 °C. Zinc was then extracted by column chromatography using the modified protocol of Moynier et al. (97) described by Jaouen et al. (34). The $\delta^{66}\text{Zn}_{\text{enamel}}$ analyses were conducted on three different multicollector inductively-coupled plasma mass spectrometers (MC-ICP-MS) instruments: a Nu 500 (Ecole Normale Supérieure), a Neptune (MPI-EVA), and a Neptune Plus (GET/OMP) using a Cu doping technique (*SI Appendix, Section S4*). The $\delta^{66}\text{Zn}_{\text{enamel}}$ values were corrected to take into account the offset (+0.27 ‰) existing between our in-house standard, bracketing standard (AA-MPI, which stands for Alfa Aesar- Max Planck Institute), and JMC-Lyon solution. Zinc isotope data are thus reported toward the JMC-Lyon values, and this correction does not impact the relative offsets or our interpretations. Strontium isotope analyses were conducted on 5 to 20 mg of enamel powder, after performing a purification step based on a modified protocol of Deniel and Pin (98) described by Richards et al. (13), using a Triton Plus thermal ionization mass spectrometer (TIMS) and a Neptune Plus at GET/OMP Toulouse. O and C isotope data were obtained from untreated tooth enamel powder (<100 μg) using the "cold trap" method with a Thermo Scientific Delta V IRMS in the Vonhof laboratory at the Max Planck Institute

for Chemistry (MPIC) after Vonhof et al. (99). The reference materials NIST SRM 1400 (for Zn), NIST SRM 1486 (for Sr), IAEA-603, NBS18, NIST SRM 120c, and three internal house standards—a carbonate standard (VICS) and a tooth enamel structural carbonate (AG-Lox) for C and O and a dentin sample (AZE) for Zn—were analyzed in the same runs as the samples and gave consistent results for each laboratory and compared with previous studies (*SI Appendix, Tables S3–S5*). Trace element analyses were conducted at the GET using an ICap (Thermo Fisher) triple quadrupole mass spectrometer (TO-ICP-MS). Finally, collagen was extracted from four tooth roots for $\delta^{15}\text{N}$ analyses using the protocol of Talamo and Richards (58). All trace elements and isotope results are available in *SI Appendix*.

Data Availability. All study data are included in the article and/or *SI Appendix*.

ACKNOWLEDGMENTS. This study was funded by Deutsche Forschungsgemeinschaft Project PALEODIET (378496604), European Research Council (ERC) Project ARCHEIS (Grant 803676), ERC Project VERTEBRATE HERBIVORY (Grant 681450), and the Max Planck Society. G.M.S. received funding from the European Union's Horizon Europe Research and Innovation Programme under Marie Skłodowska-Curie Grant Agreement 101027850. We thank Maria Alonso Lescán from the Museum of Huesca for her help during the collection sampling, Adeline Le Cabec for her help in understanding of tooth development, Sven Steinbrenner for technical help, Mary Kita for her help in the improvement of the figures, Jeremy McCormack for helpful discussions, and Hubert Vonhof for access to the stable isotope lab at MPIC Mainz. D.C.S.G. acknowledges funding by the Generalitat Valenciana (CIDEGEN/2019/061), and L.M. and P.U. from the Ministerio de Ciencia e Innovación (MICINN) (PID2020-116598GB-I00 GAPS AND DATES. DINAMICAS CULTURALES EN LA PREHISTORIA DE LA CUENCA DEL EBRO). We are grateful to the two reviewers who helped to significantly improve the manuscript.

Author affiliations: ^aGéosciences Environnement Toulouse, Observatoire Midi Pyrénées, UMR 5563, CNRS, 31400 Toulouse, France; ^bDepartment of Human Evolution, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany; ^cInstitute of Evolutionary Biology, Consejo Superior de Investigaciones Científicas-Universitat Pompeu Fabra, 08003 Barcelona, Spain; ^dDepartment of Archaeogenetics, Max Planck Institute for the Science of Human History, 07745 Jena, Germany; ^eResearch Group "Primeros Pobladores y Patrimonio Arqueológico del Valle del Ebro" Department of Sciences of Antiquity-Prehistory, University of Zaragoza, 50009 Zaragoza, Spain; ^fSchool of Anthropology and Conservation, University of Kent, Kent, CT2 7NZ, United Kingdom; ^gEmmy Noether Group for Hominin Meat Consumption, Max Planck Institute for Chemistry, 55128 Mainz, Germany; ^hApplied and Analytical Palaeontology, Institute of Geosciences, Johannes Gutenberg University, 55128 Mainz, Germany; ⁱLaboratoire de Géologie de Lyon: Terre, Planètes, Environnement, Ecole Normale Supérieure de Lyon, UMR 5276, CNRS, 69007 Lyon, France; ^jServicio de Cultura, Ayuntamiento de Zaragoza, 50071 Zaragoza, Spain; ^kMuseum of Archeology and Ethnology, University of São Paulo, 05508-070 São Paulo, Brazil; ^lDepartament de Prehistòria, Arqueologia i Història Antiga, Universitat de València, 46010 València, Spain; ^mGrupo de Investigación en Prehistoria IT-1223-19, University of the Basque Country (UPV-EHU) and Basque Foundation for Science (Ikerbasque), 01006 Vitoria, Spain; ⁿDepartment of Geological Sciences, University of Cape Town, 7700 Cape Town, South Africa; ^oDepartment of Archaeology, Simon Fraser University, Burnaby, BC V5A 1S6, Canada; and ^pChaire de Paléanthropologie, Collège de France, 75005 Paris, France

1. H. Bocherens, D. G. Drucker, D. Billiou, M. Patou-Mathis, B. Vandermeersch, Isotopic evidence for diet and subsistence pattern of the Saint-Césaire I Neandertal: Review and use of a multi-source mixing model. *J. Hum. Evol.* **49**, 71–87 (2005).
2. H. Bocherens, "Neandertal dietary habits: Review of the isotopic evidence" in *The Evolution of Hominin Diets*, J.-J. Hublin, M. P. Richards, Eds. (Springer, 2009), pp. 241–250.
3. H. Bocherens et al., New isotopic evidence for dietary habits of Neandertals from Belgium. *J. Hum. Evol.* **40**, 497–505 (2001).
4. C. Wißing et al., Isotopic evidence for dietary ecology of late Neandertals in north-western Europe. *Quat. Int.* **411**, 327–345 (2016).
5. M. P. Richards et al., Neandertal diet at Vindija and Neandertal predation: The evidence from stable isotopes. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 7663–7666 (2000).
6. M. P. Richards, R. W. Schmitz, Isotope evidence for the diet of the Neandertal type specimen. *Antiquity* **82**, 553–559 (2008).
7. K. Jaouen et al., Exceptionally high $\delta^{15}\text{N}$ values in collagen single amino acids confirm Neandertals as high-trophic level carnivores. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 4928–4933 (2019).
8. F. Welker et al., Palaeoproteomic evidence identifies archaic hominins associated with the Châtelperronian at the Grotte du Renne. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 11162–11167 (2016).
9. H. Bocherens et al., Isotopic biogeochemistry (^{13}C , ^{15}N) of fossil vertebrate collagen: Application to the study of a past food web including Neandertal man. *J. Hum. Evol.* **20**, 481–492 (1991).
10. Y. I. Naito et al., Ecological niche of Neandertals from Spy Cave revealed by nitrogen isotopes of individual amino acids in collagen. *J. Hum. Evol.* **93**, 82–90 (2016).
11. M. P. Richards, E. Trinkaus, Out of Africa: Modern human origins special feature: Isotopic evidence for the diets of European Neandertals and early modern humans. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 16034–16039 (2009).
12. H. Bocherens et al., Palaeoenvironmental and palaeodietary implications of isotopic biogeochemistry of Last Interglacial Neandertal and mammal bones in Sladina Cave (Belgium). *J. Archaeol. Sci.* **26**, 599–607 (1999).
13. M. P. Richards et al., Isotopic dietary analysis of a Neandertal and associated fauna from the site of Jonzac (Charente-Maritime), France. *J. Hum. Evol.* **55**, 179–185 (2008).
14. D. C. Salazar-García et al., Dietary evidence from Central Asian Neandertals: A combined isotope and plant microremains approach at Chagyrskaya Cave (Altai, Russia). *J. Hum. Evol.* **156**, 102985 (2021).
15. G. M. Smith, Neandertal megafaunal exploitation in western Europe and its dietary implications: A contextual reassessment of La Cotte de St Brelade (Jersey). *J. Hum. Evol.* **78**, 181–201 (2015).
16. G. J. van Klinken, Bone collagen quality indicators for palaeodietary and radiocarbon measurements. *J. Archaeol. Sci.* **26**, 687–695 (1999).
17. S. Pirson, M. Toussaint, D. Bonjean, K. Di Modica, "Spy and Sladina caves: A Neandertal's story" in *Landscapes and Landforms of Belgium and Luxembourg*, A. Demoulin, Ed. (Springer, 2018), pp. 357–383.
18. C. Wißing et al., Stable isotopes reveal patterns of diet and mobility in the last Neandertals and first modern humans in Europe. *Sci. Rep.* **9**, 4433 (2019).
19. R. Jennings, C. Finlayson, D. Fa, G. Finlayson, Southern Iberia as a refuge for the last Neandertal populations. *J. Biogeogr.* **38**, 1873–1885 (2011).
20. B. Galván et al., New evidence of early Neandertal disappearance in the Iberian Peninsula. *J. Hum. Evol.* **75**, 16–27 (2014).
21. D. C. Salazar-García et al., Neandertal diets in central and southeastern Mediterranean Iberia. *Quat. Int.* **318**, 3–18 (2013).
22. A. Rosas et al., Paleobiology and comparative morphology of a late Neandertal sample from El Sidron, Asturias, Spain. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 19266–19271 (2006).

23. J. Zilhão *et al.*, Last Interglacial Iberian Neanderthals as fisher-hunter-gatherers. *Science* **367**, eaaz7943 (2020).
24. M. Nabais, J. Zilhão, The consumption of tortoise among Last Interglacial Iberian Neanderthals. *Quat. Sci. Rev.* **217**, 225–246 (2019).
25. M. Cortés-Sánchez *et al.*, Earliest known use of marine resources by Neanderthals. *PLoS One* **6**, e24026 (2011).
26. L. S. Weyrich *et al.*, Neanderthal behaviour, diet, and disease inferred from ancient DNA in dental calculus. *Nature* **544**, 357–361 (2017).
27. R. C. Power *et al.*, Dental calculus indicates widespread plant use within the stable Neanderthal dietary niche. *J. Hum. Evol.* **119**, 27–41 (2018).
28. K. Hardy *et al.*, Neanderthal medics? Evidence for food, cooking, and medicinal plants entrapped in dental calculus. *Naturwissenschaften* **99**, 617–626 (2012).
29. H. J. O'Regan, A. L. Lamb, D. M. Wilkinson, The missing mushrooms: Searching for fungi in ancient human dietary analysis. *J. Archaeol. Sci.* **75**, 139–143 (2016).
30. C. B. Stringer *et al.*, Neanderthal exploitation of marine mammals in Gibraltar. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 14319–14324 (2008).
31. V. Michel, D. Delanghe-Sabatier, E. Bard, C. Barroso Ruiz, U-series, ESR and ¹⁴C studies of the fossil remains from the Mousterian levels of Zafarraya Cave (Spain): A revised chronology of Neanderthal presence. *Quat. Geochronol.* **15**, 20–33 (2013).
32. M. Yustos, J. Y. S. de los Ferreros, Cannibalism in the Neanderthal world: An exhaustive revision. *J. Taphon.* **13**, 33–52 (2015).
33. D. J. Álvarez-Lao, N. García, Chronological distribution of Pleistocene cold-adapted large mammal faunas in the Iberian Peninsula. *Quat. Int.* **212**, 120–128 (2010).
34. K. Jaouen, P. Szpak, M. P. Richards, Zinc isotope ratios as indicators of diet and trophic level in Arctic marine mammals. *PLoS One* **11**, e0152299 (2016).
35. K. Jaouen, M. Beasley, M. Schoeninger, J.-J. Hublin, M. P. Richards, Zinc isotope ratios of bones and teeth as new dietary indicators: Results from a modern food web (Koobi Fora, Kenya). *Sci. Rep.* **6**, 26281 (2016).
36. N. Bourgon *et al.*, Zinc isotopes in Late Pleistocene fossil teeth from a Southeast Asian cave setting preserve paleodietary information. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 4675–4681 (2020).
37. N. Bourgon *et al.*, Trophic ecology of a Late Pleistocene early modern human from tropical Southeast Asia inferred from zinc isotopes. *J. Hum. Evol.* **161**, 103075 (2021).
38. J. McCormack *et al.*, Zinc isotopes from archaeological bones provide reliable trophic level information for marine mammals. *Commun. Biol.* **4**, 683 (2021).
39. K. Jaouen *et al.*, Zinc isotope variations in archaeological human teeth (Lapa do Santo, Brazil) reveal dietary transitions in childhood and no contamination from gloves. *PLoS One* **15**, e0232379 (2020).
40. K. Jaouen, M.-L. Pons, V. Balter, Iron, copper and zinc isotopic fractionation up mammal trophic chains. *Earth Planet. Sci. Lett.* **374**, 164–172 (2013).
41. K. Jaouen *et al.*, Tracing intensive fish and meat consumption using Zn isotope ratios: Evidence from a historical Breton population (Rennes, France). *Sci. Rep.* **8**, 5077 (2018).
42. K. Jaouen, E. Herscher, V. Balter, Copper and zinc isotope ratios in human bone and enamel. *Am. J. Phys. Anthropol.* **162**, 491–500 (2017).
43. M. Costas-Rodríguez, L. Van Heghe, F. Vanhaecke, Evidence for a possible dietary effect on the isotopic composition of Zn in blood via isotopic analysis of food products by multi-collector ICP-mass spectrometry. *Metallomics* **6**, 139–146 (2014).
44. L. Van Heghe, E. Engström, I. Rodushkin, C. Cloquet, F. Vanhaecke, Isotopic analysis of the metabolically relevant transition metals Cu, Fe and Zn in human blood from vegetarians and omnivores using multi-collector ICP-mass spectrometry. *J. Anal. At. Spectrom.* **27**, 1327–1334 (2012).
45. L. Montes, P. Utrilla, "The cave of Los Moros-1 at Gabasa (Huesca)" in *Pleistocene and Holocene Hunters-Gatherers in Iberia and the Gibraltar Strait. The Current Archaeological Record*, R. Sala Ramos, Ed. (Universidad de Burgos-Fundación Atapuerca, 2014), pp. 181–188.
46. J. I. Lorenzo, L. Montes, Restes Néandertaliens de la Grotte de "Los Moros de Gabasa" (Huesca, Espagne). *Trabalhos de Arqueologia* **17**, 77–86 (2001).
47. M. Skinner, Developmental stress in immature hominines from Late Pleistocene Eurasia: Evidence from enamel hypoplasia. *J. Archaeol. Sci.* **23**, 833–852 (1996).
48. P. B. Pettitt, Neanderthal lifecycles: Developmental and social phases in the lives of the last archaics. *World Archaeol.* **31**, 351–366 (2000).
49. C. Austin *et al.*, Barium distributions in teeth reveal early-life dietary transitions in primates. *Nature* **498**, 216–219 (2013).
50. T. M. Smith *et al.*, Dental evidence for ontogenetic differences between modern humans and Neanderthals. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 20923–20928 (2010).
51. R. Macchiarelli *et al.*, How Neanderthal molar teeth grew. *Nature* **444**, 748–751 (2006).
52. T. Tsutaya, M. Yoneda, Reconstruction of breastfeeding and weaning practices using stable isotope and trace element analyses: A review. *Am. J. Phys. Anthropol.* **156** (suppl. 59), 2–21 (2015).
53. K. Jaouen *et al.*, Dynamic homeostasis modeling of Zn isotope ratios in the human body. *Metallomics* **11**, 1049–1059 (2019).
54. H. Bocherens *et al.*, Direct isotopic evidence for subsistence variability in Middle Pleistocene Neanderthals (Payre, southeastern France). *Quat. Sci. Rev.* **154**, 226–236 (2016).
55. M. Ecker *et al.*, Middle Pleistocene ecology and Neanderthal subsistence: Insights from stable isotope analyses in Payre (Ardèche, southeastern France). *J. Hum. Evol.* **65**, 363–373 (2013).
56. R. E. Stevens, R. E. M. Hedges, Carbon and nitrogen stable isotope analysis of northwest European horse bone and tooth collagen, 40,000 BP–present: Palaeoclimatic interpretations. *Quat. Sci. Rev.* **23**, 977–991 (2004).
57. J. McCormack *et al.*, Trophic position of *Otodus megalodon* and great white sharks through time revealed by zinc isotopes. *Nat. Commun.* **13**, 2980 (2022).
58. S. Talamo, M. Richards, A comparison of bone pretreatment methods for AMS dating of samples >30,000 BP. *Radiocarbon* **53**, 443–449 (2011).
59. S. Talamo, H. Fewlass, R. Maria, K. Jaouen, "Here we go again": The inspection of collagen extraction protocols for ¹⁴C dating and palaeodietary analysis. *Sci. Technol. Archaeol. Res.* **7**, 62–77 (2021).
60. M. F. Blasco, In the pursuit of game: The Mousterian cave site of Gabasa 1 in the Spanish Pyrenees. *J. Anthropol. Res.* **53**, 177–217 (1997).
61. F. Blasco Sancho, *Hombres, Fieras y Presas: Estudio Arqueozoológico y Tafonómico del Yacimiento del Paleolítico Medio de la Cueva de Gabasa 1* (Universidad de Zaragoza, Huesca, Spain, 1995).
62. K. Jaouen, What is our toolbox of analytical chemistry for exploring ancient hominin diets in the absence of organic preservation? *Quat. Sci. Rev.* **197**, 307–318 (2018).
63. Y. I. Naito *et al.*, Evidence for herbivorous cave bears (*Ursus spelaeus*) in Goyet Cave, Belgium: Implications for palaeodietary reconstruction of fossil bears using amino acid $\delta^{15}\text{N}$ approaches. *J. Quat. Sci.* **31**, 598–606 (2016).
64. S. C. Münzel *et al.*, Behavioural ecology of Late Pleistocene bears (*Ursus spelaeus*, *Ursus ingressus*): Insight from stable isotopes (C, N, O) and tooth microwear. *Quat. Int.* **339**, 148–163 (2014).
65. H. Bocherens, M. Fizet, A. Mariotti, Diet, physiology and ecology of fossil mammals as inferred from stable carbon and nitrogen isotope biogeochemistry: Implications for Pleistocene bears. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **107**, 213–225 (1994).
66. M. P. Richards *et al.*, Isotopic evidence for omnivory among European cave bears: Late Pleistocene *Ursus spelaeus* from the Peștera cu Oase, Romania. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 600–604 (2008).
67. M. Sponheimer, D. de Ruiter, J. Lee-Thorp, A. Späth, Sr/Ca and early hominin diets revisited: New data from modern and fossil tooth enamel. *J. Hum. Evol.* **48**, 147–156 (2005).
68. W. J. Binder, B. Van Valkenburgh, Development of bite strength and feeding behaviour in juvenile spotted hyenas (*Crocuta crocuta*). *J. Zool.* **252**, 273–283 (2000).
69. F. Moynier, D. Vance, T. Fujii, P. Savage, The isotope geochemistry of zinc and copper. *Rev. Mineral. Geochem.* **82**, 543–600 (2017).
70. P. González-Sampériz, L. Montes, P. Utrilla, Pollen in hyena coprolites from Gabasa Cave (northern Spain). *Rev. Palaeobot. Palynol.* **126**, 7–15 (2003).
71. P. Utrilla, L. Montes, B. Fernanda, T. J. Torres Pérez-Hidalgo, J. E. Ortiz Menéndez, La cueva de Gabasa revisada 15 años después: Un cubil para las hienas y un cazadero para los Neandertales. *Zona Arqueológica* **13**, 376–389 (2010).
72. H. Bocherens, M. L. Fogel, N. Tuross, M. Zeder, Trophic structure and climatic information from isotopic signatures in Pleistocene cave fauna of southern England. *J. Archaeol. Sci.* **22**, 327–340 (1995).
73. Q. Li *et al.*, Spatially-resolved Ca isotopic and trace element variations in human deciduous teeth record diet and physiological change. *Environ. Archaeol.* **10**.1080/14614103.2020.1758988 (2020).
74. D. Guatelli-Steinberg, Recent studies of dental development in Neanderthals: Implications for Neanderthal life histories. *Evol. Anthropol.* **18**, 9–20 (2009).
75. R. Joannes-Boyau *et al.*, Elemental signatures of *Australopithecus africanus* teeth reveal seasonal dietary stress. *Nature* **572**, 112–115 (2019).
76. M. C. Dean, B. H. Smith, "Growth and development of the Nariokotome youth, KNM-WT 15000" in *The First Humans—Origin and Early Evolution of the Genus Homo*, F. E. Grine, J. G. Fleagle, R. E. Leakey, Eds. (Springer, 2009), pp. 101–120.
77. F. Moynier, T. Fujii, A. S. Shaw, M. Le Borgne, Heterogeneous distribution of natural zinc isotopes in mice. *Metallomics* **5**, 693–699 (2013).
78. V. Balter *et al.*, Contrasting Cu, Fe, and Zn isotopic patterns in organs and body fluids of mice and sheep, with emphasis on cellular fractionation. *Metallomics* **5**, 1470–1482 (2013).
79. J. Viers *et al.*, Evidence of Zn isotopic fractionation in a soil-plant system of a pristine tropical watershed (Nsimi, Cameroon). *Chem. Geol.* **239**, 124–137 (2007).
80. C. Cloquet, J. Carignan, M. F. Lehmann, F. Vanhaecke, Variation in the isotopic composition of zinc in the natural environment and the use of zinc isotopes in biogeochemistry: A review. *Anal. Bioanal. Chem.* **390**, 451–463 (2008).
81. C. Caldeas, D. J. Weiss, Zinc homeostasis and isotopic fractionation in plants: A review. *Plant Soil* **411**, 17–46 (2017).
82. R. D. Evans, W. Wang, H. E. Evans, R. B. Georg, Variation in Zn, C, and N isotope ratios in three stream insects. *Facets* **1**, 205–216 (2016).
83. K. N. Nitzsche *et al.*, Magnesium and zinc stable isotopes as a new tool to understand Mg and Zn sources in stream food webs. *Ecosphere* **11**, e03197 (2020).
84. V. Mourre *et al.*, "Exploitation du milieu montagnard dans le Moustérien final: La Grotte du Noisetier à Fréchet-Aure (Pyrénées Centrales Françaises)" in *Proceedings of the XV World Congress UISPP (Lisbon, 4e9 September 2006). Mountain Environments in Prehistoric Europe: Settlement and Mobility Strategies*, S. Grimaldi, Th. Perrin, J. Guilaine, Eds. (Archaeopress, Oxford, UK, 2008), pp. 1–10.
85. H. Scherz, E. Kirchoff, Trace elements in foods: Zinc contents of raw foods—A comparison of data originating from different geographical regions of the world. *J. Food Compos. Anal.* **19**, 420–433 (2006).
86. G. Monge *et al.*, Earliest evidence of pollution by heavy metals in archaeological sites. *Sci. Rep.* **5**, 14252 (2015).
87. A. Coutaud, M. Meheut, J. Viers, J.-L. Rols, O. S. Pokrovsky, Zn isotope fractionation during interaction with phototrophic biofilm. *Chem. Geol.* **390**, 46–60 (2014).
88. B. Mahan, F. Moynier, A. L. Jørgensen, M. Habekost, J. Siebert, Examining the homeostatic distribution of metals and Zn isotopes in Göttingen minipigs. *Metallomics* **10**, 1264–1281 (2018).
89. K. Jaouen, V. Balter, Menopause effect on blood Fe and Cu isotope compositions. *Am. J. Phys. Anthropol.* **153**, 280–285 (2014).
90. F. Blasco, L. Montes, P. Utrilla, "Deux modèles de stratégie occupationnelle dans le Moustérien tardif de la vallée de l'Ebre: Les grottes de Peña Miel et Gabasa" in *The Last Neanderthals, the First Anatomically Modern Humans: A Tale about the Human Diversity*, E. Carbonell, M. Vaquero, Eds. (Universitat Rovira i Virgili, Tarragona, Spain, 1996), pp. 385–434.
91. L. Lloveras, M. Moreno-García, J. Nadal, Taphonomic analysis of leporid remains obtained from modern Iberian lynx (*Lynx pardinus*) scats. *J. Archaeol. Sci.* **35**, 1–13 (2008).
92. G. Haynes, A guide for differentiating mammalian *Carnivore* taxa responsible for gnaw damage to herbivore limb bones. *Paleobiology* **9**, 164–172 (1983).
93. D. G. Drucker *et al.*, Isotopic analyses suggest mammoth and plant in the diet of the oldest anatomically modern humans from far southeast Europe. *Sci. Rep.* **7**, 6833 (2017).
94. S. Lösch, G. Grupe, J. Peters, Stable isotopes and dietary adaptations in humans and animals at pre-pottery Neolithic Nevalli Cori, southeast Anatolia. *Am. J. Phys. Anthropol.* **131**, 181–193 (2006).
95. R. E. M. Hedges, L. M. Reynard, Nitrogen isotopes and the trophic level of humans in archaeology. *J. Archaeol. Sci.* **34**, 1240–1251 (2007).
96. P. Langer, M. Claus, Morphological adaptation of the eutherian gastrointestinal tract to diet. *Vertebr. Zool.* **68**, 237–252 (2018).
97. F. Moynier, F. Albarède, G. F. Herzog, Isotopic composition of zinc, copper, and iron in lunar samples. *Geochim. Cosmochim. Acta* **70**, 6103–6117 (2006).
98. C. Deniel, C. Pin, Single-stage method for the simultaneous isolation of lead and strontium from silicate samples for isotopic measurements. *Anal. Chim. Acta* **426**, 95–103 (2001).
99. H. B. Vonhof *et al.*, High-precision stable isotope analysis of <5 μg CaCO_3 samples by continuous-flow mass spectrometry. *Rapid Commun. Mass Spectrom.* **34**, e8878 (2020).