

Journal section: Oral Medicine and Pathology
 Publication Types: Review

doi:10.4317/jced.59830
<https://doi.org/10.4317/jced.59830>

Antimicrobial resistance in patients with odontogenic infections: A systematic scoping review of prospective and experimental studies

Carlos-M. Ardila ^{1,2}, Jader-Alexander Bedoya-García ^{2,3}

¹ DDS. Periodontist. Ph. D in Epidemiology. Postdoc in Bioethics. Titular Professor. Universidad de Antioquia U de A, Medellín, Colombia

² Biomedical Stomatology Research Group, Universidad de Antioquia U de A, Medellín, Colombia

³ DDS. Periodontist. Professor. Universidad de Antioquia U de A, Medellín, Colombia

Correspondence:
 Calle 70 No. 52-21
 Medellín, Colombia
martin.ardila@udea.edu.co

Ardila CM, Bedoya-García JA. Antimicrobial resistance in patients with odontogenic infections: A systematic scoping review of prospective and experimental studies. J Clin Exp Dent. 2022;14(10):e834-45.

Received: 25/06/2022
 Accepted: 16/08/2022

Article Number: 59830 <http://www.medicinaoral.com/odo/indice.htm>
 © Medicina Oral S. L. C.I.F. B 96689336 - eISSN: 1989-5488
 eMail: jced@jced.es
Indexed in:
 Pubmed
 Pubmed Central® (PMC)
 Scopus
 DOI® System

Abstract

Background: Patients with odontogenic infections are commonly prescribed antimicrobials on an experiential base without knowing the precise microorganisms implicated. The aim of this systematic scoping review is to evaluate the prevalence and proportions of antimicrobial-resistant species in patients with odontogenic infections.

Material and Methods: A systematic scoping review of scientific evidence was accomplished involving different databases.

Results: Eight randomized clinical trials and 13 prospective observational studies were included. These investigations analyzed 1506 patients. The species that showed higher levels of resistance included aerobic and facultative anaerobe such as *Staphylococcus aureus*, *Streptococcus viridans*, *Klebsiella pneumoniae*, *Streptococcus milleri*, *Enterococcus spp.*, *Pseudomonas aeruginosa*, *Proteus mirabilis*, and *Staphylococcus coagulases-negative*. In obligate anaerobes sampled were *Peptostreptococcus spp.*, *Bacteroides spp.*, and *Prevotella spp.* *Staphylococcus* showed resistance to ampicillin, piperacillin, clindamycin, amoxicillin, metronidazole, and penicillin. *Streptococcus* had resistance to metronidazole, clindamycin, doxycycline, penicillin, and amoxicillin. *Peptostreptococcus spp.* presented resistance to penicillin, amoxicillin, erythromycin, and cefalexin. Gram-negative microorganisms had resistance to tetracycline, ciprofloxacin, azithromycin, amoxicillin, erythromycin, and penicillin. *Bacteroides spp.* exhibited resistance to penicillin, erythromycin, and gentamicin. *Prevotella spp.* showed resistance to penicillin, amoxicillin, erythromycin, clindamycin, levofloxacin, and imipenem. Finally, *Klebsiella spp.* displayed resistance to ampicillin, amoxicillin, moxifloxacin, and cefalexin. Interestingly, one clinical trial showed that after therapy there was a reduction in sensitivity of 18% for azithromycin and 26% for spiramycin.

Conclusions: Most of the microorganisms had resistance to diverse groups of antimicrobials. Suitable antimicrobials must be prescribed founded on the microbial samples, culture susceptibility, and clinical progression of the odontogenic infection. Furthermore, it was observed high levels of resistance to antimicrobials that have been used in local and systemic therapy of oral cavity infections. A preponderance of anaerobic microorganisms over aerobic ones was observed.

Key words: Antibiotic resistance, odontogenic infections, efficacy, microorganisms, scoping review.

Introduction

Odontogenic infection is the most commonly appearing infection in the orofacial area. These infections comprise from periapical abscesses to mild and profound infections in the neck and are frequently caused by periodontitis and dental caries as well as pericoronitis and complications during dental procedures (1).

It has been recognized that the treatment of odontogenic orofacial and neck infections is mainly oriented to the clinical alleviation of suppuration. Nonetheless, antimicrobials adjunct to that therapy is relevant, particularly when there is systemic compromise (2). The empirical choice of appropriate antimicrobials for the management of these infections is supported by their clinical efficacy, low prices, few adverse events, and good availability (3).

This empirical management has generated complications related to the use of antimicrobials, an issue that in turn has allowed investigating of regular prescription practices by dentists (4,5). The selection of antimicrobial for the treatment of odontogenic infections preferably requires the performance of a microbial culture to carry out susceptibility tests. Nevertheless, it has been indicated that 46% of dentists from different countries disregarded this conduct before the recommendation of antimicrobials, albeit 83% of the total clinicians interrogated were conscious of the growth in antimicrobial resistance (5). Therefore, patients with these odontogenic infections are commonly prescribed antimicrobials on an experiential base without knowing the precise microorganisms implicated. This antimicrobial management could or could not generate satisfactory effects due to diverse reasons such as bacterial specificity and antimicrobial resistance (6).

On the other hand, geographical differentiation, the occurrence of resistant microorganisms, and native antibacterial prescribing policies generate variability in the antimicrobial profile of pathogens between communities (7). Since the development of antimicrobial resistance during antimicrobial management in dental practice is a matter of concern, it is relevant to carry out a scoping review that allows for evaluating the antimicrobial resistance patterns by phenotypic identification of the microorganisms most commonly isolated from odontogenic infections. To achieve this objective, it was proposed to answer some questions related to antimicrobial resistance, in terms of the prevalence and proportions of antimicrobial-resistant species in odontogenic infections. Furthermore, the antimicrobials to which the odontogenic pathogens present resistance were also investigated.

Material and Methods

This review of prospective and experimental studies in humans was carried out considering the PRISMA (Preferred Reporting Items for Systematic Reviews and

Meta-analyses) extension for scoping reviews (8). The scoping structure involved different databases such as PubMed/MEDLINE, SCOPUS, SCIELO, and LILACS, including the gray literature. MeSH terms and keywords were used to investigate publications in all languages until March 2022, integrating the terms odontogenic, infections, antibacterial drug resistance, dental infection, antibiotic resistance, antibiotics, alveolar abscess, dentoalveolar abscesses, antibacterial susceptibility breakpoint determination, bacterial sensitivity tests, and prospective and experimental studies. Then, a searching process was implemented to explore databases using Boolean operators (AND, OR): “odontogenic” OR “infections” OR “antibacterial drug resistance” OR “dental infections” OR “antibiotic resistance” OR “antibiotics” AND “dentoalveolar abscesses” AND “antibiotic resistance” OR “antimicrobials” OR “alveolar abscess” OR “microorganisms” OR “antibacterial susceptibility breakpoint determination”, OR “bacterial sensitivity tests”.

-Resources selection

Only prospective and experimental studies involving persons diagnosed at the beginning of the study with moderate to severe orofacial/dentoalveolar infection of odontogenic origin, and studies containing phenotypic analysis and antimicrobial susceptibility results were selected. Furthermore, lactating and pregnant women, patients in whom it was not feasible to acquire a proper pus sample, or if systemic antimicrobial was not necessary, or presenting a coexisting systemic illness, were not contemplated for this review. Duplicate publications and analyses applied to animals were also not incorporated.

-Questions

This scoping review aims to answer the following questions: What is the prevalence of antimicrobial-resistant species in patients with odontogenic infections? What is the proportion of antimicrobial-resistant species in patients with odontogenic infections? To which antibiotics did the microorganisms show resistance?

-Review process

Both investigators assessed the titles and abstracts and chose prospective and experimental studies to consider the full text for probable suitability. In case of discrepancy among authors, research eligibility was defined by agreement. The Kappa test was implemented to calculate the score of agreement among researchers (>85).

-Data collection

A table was considered to include the most pertinent information from the chosen reports. This procedure was completed individually by each of the investigators. Successively, the records were compared. Documented information contained authors' names, date of publication, amount of patients and quantity of isolates assessed, the occurrence of antimicrobial-resistant bacteria, the percentage of antimicrobial-resistant microorganisms, and antimicrobials in which resistance was observed.

-Risk of Bias

Both authors of this scoping review, independently assessed the methodological quality of the included investigations, using a previously described instrument (9). The instrument contains 16 conditions. A value from 0 to 3 is given to each criterion (0=it does not provide the level of detail needed to generate a decision for a criterion; 1=slightly provided; 2=moderately provided; 3=completely provided). The sum of these criteria gives a total result for the body of evidence, stated as a proportion of the maximum probable score.

Results

The electronic exploration conceded 560 investigations. After evaluating the titles and abstracts, 79 studies were eliminated for their unimportance, and 4 duplicate publications were also ignored. Reading the full text occasioned the omission of 456 additional investigations because they did not meet some selection criteria. Finally, 8 randomized clinical trials (RCTs) (10-17) and 13 (18-30) prospective observational studies were included in this scoping review (Fig. 1).

The features of the incorporated studies are shown in Table 1-1 cont.-3. These researches were published between 1987 (17) and 2021 (24). These investigations assessed 1506 participants with a minimum sample of 21(10) patients and a maximum of 142 (26).

Antibiotic resistance to a wide variety of antimicrobials was explored, including cephalosporins, metronidazole, penicillin, amoxicillin, amoxicillin-clavulanic acid, tetracycline, doxycycline, clindamycin, ampicillin, ciprofloxacin, gentamycin, erythromycin, azithromycin, imipenem, spiramycin, linezolid, vancomycin, bacitracin,

amikacin, piperacillin, moxifloxacin, and levofloxacin. However, the most studied antibiotics were penicillin, clindamycin, metronidazole, amoxicillin and, amoxicillin/clavulanic acid.

On the other hand, concerning the prevalence of antimicrobial-resistant species, it was observed that a great variety of microorganisms were isolated (Table 1); nevertheless, the species that showed higher levels of resistance included *Staphylococcus*, *Streptococcus spp.*, *Peptostreptococcus spp.*, *Prevotella spp.*, and *Bacteroides spp.* Among aerobic and facultative anaerobe prevailed, *Staphylococcus aureus* (12,18,19,25,28), *Streptococcus viridans* (15,17,26,27), *Klebsiella pneumoniae* (12,19,22,27), *Streptococcus milleri* (22), *Enterococcus spp.* (25), *Pseudomonas aeruginosa* (28), *Proteus mirabilis* (19), and *Staphylococcus coagulases-negative* (12). In obligate anaerobes sampled were *Peptostreptococcus spp.* (12,15,17,18,20,21,23,24,26,30), *Bacteroides spp.* (12,14,18,21,24,26,28-30), and *Prevotella spp.* (11,18,20,21,23,26,30).

All the investigations implemented different protocols for the identification of the microorganisms studied. Nonetheless, most researchers used the disk diffusion protocol to check susceptibility to antimicrobials and interpreted following recognized guides.

Table 1 also details the different proportions of antimicrobial-resistant species. The studies found that *Staphylococcus* showed resistance to ampicillin (19), piperacillin (19), clindamycin (10), amoxicillin (12), metronidazole (18,21), and penicillin (17,24,25,28-30). *Streptococcus* had resistance to metronidazole (23), clindamycin (10), doxycycline (11), penicillin (28,29), and amoxicillin (26,27,31). *Peptostreptococcus spp.* presen-




Fig. 1: Flowchart of the studies selection method.

Table 1: Features of the studies evaluated.

Authors Publication date	Patients/samples	Age	Prevalence of isolated bacteria	Proportions of antimicrobial-resistant species
Umeshappa <i>et al.</i> 2021 (24)	100/115	14-65 years	<i>S. aureus</i> and <i>S. viridans</i> were the most predominant isolates (50%), followed by Peptostreptococcus spp. (23.75%) Bacteroides spp. and Prevotella spp.	Global resistance to penicillin was 41.5% among obligate anaerobes due to beta-lactamase production, while amoxicillin/clavulanic acid showed absolute susceptibility. Seventy-three percent of microorganisms had resistance to erythromycin. Cefotaxime and ciprofloxacin showed good efficacy (83% each). Bacteroides spp. presented resistance to erythromycin and gentamicin. Metronidazole showed efficacy only against obligate anaerobes.
Uppada & Sinha 2020 (25)	124/144	21-40 years	Staphylococcus spp. (44%), Enterococcus spp. (23%), Streptococcus spp. (19%).	<i>S. aureus</i> presented resistance to penicillin (29%) while ciprofloxacin, cephalosporin, and clindamycin showed 100% of efficacy against this microorganism. Most of the Gram-negative bacilli and anaerobes had susceptibility to metronidazole.
Sebastian <i>et al.</i> 2019 (26)	142/125	35 years on average	Peptostreptococcus (62%), <i>S. viridans</i> (35%), Bacteroides (27.78%).	There was resistance to amoxicillin in 97% of aerobic bacteria, 86% of anaerobic microorganisms, and 86% of mixed pathogens. The entire aerobic, anaerobic, and mixed group of bacteria had susceptibility to linezolid (100%). All the anaerobic microorganisms had susceptibility to metronidazole (100%) and 65% of mixed bacteria presented susceptibility to metronidazole. Thirty-five percent of mixed microorganisms showed resistance to metronidazole. All the aerobic pathogens were susceptible to clindamycin (100%), and 83% of anaerobic groups were susceptible to clindamycin. A total of 63% of mixed bacteria were susceptible to clindamycin. Resistance to clindamycin was 17% in anaerobes and 37% in mixed pathogens. Vancomycin and bacitracin were highly resistant. In the macrolide group, microorganisms presented high resistance to erythromycin (97%). Resistance to azithromycin was also observed in 80% of aerobic bacteria, 78% of anaerobic microorganisms, and 70% of mixed pathogens.
Shakia <i>et al.</i> 2018 (18)	125/167	37 years on average	In the aerobic/microaerophilic group, 17 different species were isolated. Streptococcus spp. was the most common, and <i>S. viridans</i> was the prevalent isolated microorganism (n=48) followed by <i>S. aureus</i> (n=20) and <i>E. faecalis</i> (n=8). Eighteen diverse species of anaerobes were isolated (n=65). The most numerous isolates were <i>F. nucleatum</i> (17%; n=11), followed by Prevotella spp. (11%; n = 7).	Amoxicillin/clavulanic-acid, penicillin and clindamycin presented good efficacy against the aerobic isolates (100%, 97% and 99%, respectively). Cefotaxime displayed substantial efficacy against the aerobic samples (91%). <i>S. aureus</i> showed resistance against ciprofloxacin (12.5%). The resistance of aerobic samples to metronidazole is recognized information. Metronidazole, ciprofloxacin, amoxicillin/clavulanic acid, and clindamycin had good efficacy against all the anaerobic isolates (100%, 91%, and 86%, correspondingly). Penicillin had low efficacy against anaerobes (25%).

Table 1 cont.: Features of the studies evaluated.

Jagadish <i>et al.</i> 2017 (19)	37/31	41 years on average	Twentyone isolates (51.2%) were Gram-positive cocci and 20 (48.8%) were Gram-negative bacilli. Enterobacteriales (41%) were dominant followed by Bacillales (32%), Lactobacillales (20%), and Pseudomonadales (7%), correspondingly.	Thirty-eight percent of Gram-positive aerobes showed resistance to piperacillin, being <i>S. aureus</i> resistant 50% to piperacillin, and 100% sensitive to gentamicin. Most isolates of Staphylococcus were resistant to ampicillin. Staphylococcus isolates were susceptible to cefotaxime, azithromycin and ciprofloxacin in 92%, 77%, and 77%, respectively. Of all Streptococcus isolates, 100 % were sensitive to ampicillin followed by 83.3 % to ciprofloxacin, gentamicin, and ceftazidime. Gram-negative microorganisms showed susceptibility to tetracycline, ciprofloxacin, and azithromycin (71%, 70%, and 65%, respectively). <i>E. coli</i> was 100% sensitive to ampicillin. <i>Klebsiella quasipneumoniae</i> , <i>K. pneumoniae</i> , <i>Proteus mirabilis</i> , and <i>Shigella flexneri</i> were 100 % resistant to ampicillin. <i>E. coli</i> , <i>P. aeruginosa</i> , and <i>C. koseri</i> were 100 % sensitive to ciproflaxin. <i>E. cloacae</i> was 100% sensitive to azithromycin while <i>P. mirabilis</i> showed resistance.
Shah <i>et al.</i> 2016 (27)	100/100	36 years on average	Aerobic Gram-positive (73%), aerobic Gram-negative (18%). <i>S. viridans</i> (47%), <i>S. aureus</i> (16%) <i>Klebsiella pneumoniae</i> (11%).	<i>S. viridans</i> presented 34% of resistance to amoxicillin while showed susceptibility to amoxicillin/clavulanic acid (68%), ceftriaxone (89%), carbenicillin, amikacin, and imipenem (100%) Moxifloxacin presented intermediate susceptibility (64%). <i>S. aureus</i> presented 31% of resistance to amoxicillin while amoxicillin/clavulanic acid had 100% of susceptibility. <i>K. pneumoniae</i> presented resistance to amoxicillin (64%), and moxifloxacin (36%). This microorganism was 100% susceptible to ceftriaxone, carbenicillin, amikacin, and imipenem.
Gómez-Arámbula <i>et al.</i> 2015 (10)	21/43	42 years on average	There was a preponderance of facultative and moderate anaerobe bacteria, including streptococci (23%), aerococci (21%), and staphylococci (12%).	All the isolated microorganisms showed low resistance to moxifloxacin and ceftriaxone (3% and 8%, respectively), while 35% were resistant to clindamycin.
Walia <i>et al.</i> 2014 (28)	42/40	38 years on average	Seventy percent were aerobes, 64% were Gram-positive aerobes and 36% were Gram-negative aerobes. <i>S. aureus</i> was the principally sampled pathogen (18%). The Gram-negative aerobes were sampled in 25% of patients. The prevalent anaerobic sampled was Peptostreptococcus (10%), <i>Bacteroides melaninogenicus</i> (5%), and <i>Bacteriodes fragilis</i> in (5%). The Gram-negative aerobes were presented in 25% of patients (<i>Klebsiella</i> , 10%; <i>E. coli</i> , 10%; and <i>Pseudomonas aeuroginosa</i> , 5%).	Thirty-nine percent of Gram-positive aerobes presented resistance to penicillin, of which, <i>S. aureus</i> had resistance 71% to penicillin and erythromycin, whereas 100 % of susceptibility was observed to gentamicin, ciprofloxacin, and cefotaxime. Few samples of staphylococcus were susceptible to penicillin. <i>E. coli</i> and <i>Klebsiella</i> had 100% of susceptibility to amikacin, whereas pseudomonas showed 100% of resistance to amikacin, but had susceptibility to cefotaxime, cefuroxime, and ciprofloxacin.

Table 1 cont.-1: Features of the studies evaluated.

Fating <i>et al.</i> 2014 (29)	26/26	36 years on average	Gram-positive cocci (74%) followed by Gram-negative bacilli (19%). The most prevalent microorganisms were Alpha hemolytic streptococci (70%), Beta hemolytic streptococci (10%), <i>S. aureus</i> (10%), Acinetobacter (5%), and klebsiella species (5%). Thirty-eight samples of anaerobic microorganisms were observed (anaerobic streptococci, 63%; Bacteroides 13%, and Fusobacterium 8%).	All the aerobic pathogens had susceptibility to gentamycin, vancomycin, imipenem, and linezolid (100%). Eighty percent of the strains had susceptibility to penicillin G, amoxicillin, and amoxicillin/clavulanic acid. Twenty percent of the sampled strains had resistance to penicillin G, amoxicillin/clavulanic acid, and amoxicillin, while 10% of them presented resistance to doxycycline and cefixime.
Singh <i>et al.</i> 2014 (30)	30/30	32 years on average	Strict anaerobes (43%), aerobes (39%), and mixed growth 19%. Among aerobes, alpha-hemolytic <i>S. aureus</i> (37%) and Peptostreptococcus (37%) as anaerobes were the most prevalent followed by Bacteroides and Prevotella (7%).	Aerobes presented resistance to penicillin at 22% while amoxicillin/clavulanic acid was 100% effective. It was observed low susceptibility to the macrolide group (37% to erythromycin). Cefuroxime showed an efficacy of 47% while cefotaxime and ciprofloxacin presented 83%. Amikacin showed efficacy on all microorganisms tested.
Sobottka <i>et al.</i> 2012 (11)	71/205	NR	The most predominant microorganisms were Prevotella species (n=56), <i>Streptococcus mitis</i> (n=53), other <i>viridans</i> group streptococci (n=24), and Neisseria spp. (n=19).	Ninety-eight percent, 96%, 85%, 60%, and 50% of all odontogenic microorganisms were susceptible to moxifloxacin, amoxicillin/clavulanic acid, levofloxacin, penicillin, clindamycin, and doxycycline, respectively. <i>S. mitis</i> presented high resistance to doxycycline (75%). Clindamycin lacked efficacy against several species of Neisseria. Penicillin presented rates of susceptibility from 0% for diverse species of Neisseria to 100% for <i>Streptococcus anginosus</i> group/hemolytic streptococci. Proportions of susceptibility to levofloxacin fluctuated from 100% for Neisseria spp. and <i>P. intermedia</i> , to 73% for <i>P. oralis</i> and other anaerobes. Anaerobes and Neisseria spp. were 100% susceptible to amoxicillin-clavulanic acid, while 83% of samples of other <i>viridans</i> group streptococci had a susceptibility.
Matijević <i>et al.</i> 2009 (12)	90/90	45 years on average	A total of 111 microorganisms strains were studied. The most frequent were Gram-positive facultative anaerobic microorganisms (81%), principally <i>S. viridans</i> (75%).	Resistance of isolated microorganisms to amoxicillin and cefalexin was 24%, and 11%, respectively. <i>S. viridans</i> had high susceptibility to amoxicillin and cefalexin (98% and 95%, respectively). <i>S. coagulans</i> -negative and <i>S. aureus</i> were resistant to amoxicillin (89% and 100%, respectively). Klebsiella spp. and Serratia spp. showed resistance to amoxicillin (100%), and cefalexin (75%). Peptostreptococcus also presented 50% of resistance to both antimicrobials.
Chardin <i>et al.</i> 2009 (13)	81/81	32 years on average	Oral streptococci	The percentage of streptococci with diminished susceptibility to amoxicillin varied depending on the day of evaluation (days 0, 9, and 30).

Table 1 cont.-2: Features of the studies evaluated.

Al-Nawas & Maeurer. 2008 (20)	30/30	NR	The prevalent microorganisms were Prevotella spp. (n=17), Pepto-streptococcus spp., (n=15) and Propionibacterium spp. (n=5).	A total of 87% of the samples were susceptible to penicillin while 97% of the anaerobes were susceptible to amoxicillin/clavulanic acid, imipenem plus cilastatin, and clindamycin. A total of 83% were susceptible to metronidazol. <i>F. nucleatum</i> and <i>Prevotella disiens</i> were not fully susceptible to imipenem plus cilastatin.
Boyanova et al. 2006 (21)	118/118	NR	Anaerobic bacteria were observed in 75% of the samples, while anaerobes were shown in 20% of the isolates. The prevalent microorganisms were Prevotella (28%), Fusobacterium (13%), Actinomyces spp. (21), anaerobic cocci (12%) and Eubacterium spp. (10%). <i>Bacteroides fragilis</i> strains were isolated from 7 specimens.	Gram-negative anaerobes were resistant to amoxicillin, clindamycin, and metronidazole in 27%, 5%, and 3%, respectively. Gram-positive species showed resistance to clindamycin and metronidazole (5% and 58%, respectively).
Flynn et al. 2006 (22)	37/37	35 years on average	A total of 8% of the cases had aerobic microorganisms only, 17% showed anaerobes only, and 67% presented a mixed flora.	Nineteen percent of the isolated strains showed resistance to penicillin. Four clindamycin-resistant bacteria were observed, one each of <i>Streptococcus milleri</i> , <i>Eikenella corrodens</i> , and <i>Streptococcus mitis</i> , and one strain of <i>K. pneumoniae</i> that also showed resistance to penicillin. Resistance to clindamycin was also observed in 17% of the cases.
Kuriyama et al. 2005 (23)	112/112	37 years on average	The most common microorganisms were Prevotella, Peptostreptococcus, streptococci, and Fusobacterium species.	A total of 30% of Prevotella species showed resistance to penicillin. All strains of Eikonella species, and Veillonella species presented resistance to penicillin. Fusobacterium species, Eikonella species and Veillonella species presented decreased antimicrobial susceptibility to erythromycin. All streptococcal samples presented resistance to metronidazole; however, all samples of Prevotella, Peptostreptococcus, and Fusobacterium species showed susceptibility to this antibiotic.
Gilmore et al. 1998 (14)	55/55	NR	Seventy-four percent of the isolates had a mixture of facultative and obligate anaerobic microorganisms, 20% presented only anaerobic bacteria, and 6% showed only aerobic pathogens. <i>S. viridans</i> was observed in 6% of all samples while Bacteroides species (not fragilis) were found in 75% of the samples.	Nine percent of the aerobic and facultative samples presented resistance to penicillin, and 12% had resistance to clindamycin, while 9% of the anaerobic cultures showed resistance to penicillin and 2% to clindamycin.
Lo et al. 1993 (15)	60/60	NR	<i>S. viridans</i> and Peptostreptococcus spp. were the most prevalent bacteria.	It was observed a diminution in susceptibility to azithromycin and spiramycin after therapy.

Table 1 cont.-3: Features of the studies evaluated.

Gorbach <i>et al.</i> 1991 (16)	55/157	NR	Mixed cultures of facultative and anaerobic microorganisms were observed in 74% of patients, anaerobes in 20%, and facultative bacteria only in 3 patients. Streptococci isolates were observed in over 90 % of positive cultures. Among the strict anaerobic microorganisms, <i>Bacteroides</i> spp. (not <i>Bacteroides fragilis</i>) were presented in 75% of the cultures.	The general resistance to penicillin was observed in 9% of the samples, while 2% were resistant to clindamycin. No one <i>Bacteroides</i> showed resistance to clindamycin.
Quayle <i>et al.</i> 1987 (17)	50/44	NR	<i>S. viridans</i> and Peptostreptococcus spp. were the most prevalent bacteria.	Seventeen anaerobes and 48% of aerobes showed resistance to penicillin.

ted resistance to penicillin (24,30), amoxicillin (12,26), erythromycin (30), and cefalexin (12,30). *Prevotella* spp. showed resistance to penicillin (18,23), amoxicillin (21), erythromycin (23), clindamycin (26), levofloxacin (11), and imipenem (20). Gram-negative microorganisms had resistance to tetracycline, ciprofloxacin, azithromycin (19,26), amoxicillin (21,26), erythromycin (23,24), and penicillin (18,23,24). *Bacteroides* spp. displayed resistance to penicillin (24), erythromycin (24,30), and gentamicin (24). Finally, *Klebsiella* spp. exhibited resistance to ampicillin (19), amoxicillin (12,27), moxifloxacin (27), and cefalexin (12). Interestingly, in one RCT, at baseline 75% of *S. viridans* and Peptostreptococcus spp. were susceptible to azithromycin and 63% to spiramycin. However, after therapy, 57% had susceptibility to azithromycin and 37% to spiramycin, with a reduction in sensitivity of 18% for azithromycin and 26% for spiramycin (15). Similarly, in another RCT, the percentage of streptococci with diminished susceptibility to amoxicillin ranged from 1.3 % of the total streptococci on day 0 to 23% on day 9, and 7.7% on day 30 (13).

On the other hand, it was found that *Staphylococcus* had good susceptibility to cefotaxime (19,25,28), ceftriaxone (10), azithromycin (19), clindamycin (25,26), amoxicillin/clavulanic acid (24,27), ciprofloxacin (25,28), and moxifloxacin (11). *Streptococcus* presented good susceptibility to ampicillin, ciprofloxacin, gentamicin, ceftazidime (19), ceftriaxone, moxifloxacin (10), amoxicillin/clavulanic-acid (24), linezolid (26), clindamycin (26), and amoxicillin (12,13). Peptostreptococcus spp. showed susceptibility to amoxicillin-clavulanic acid (20,24), imipenem plus cilastatin (20), clindamycin (20), linezolid (26), and metronidazole (23). Gram-negative microorganisms displayed susceptibility to levofloxacin, amoxicillin-clavulanic acid (11), and metronidazole (23,26). Lastly, *Escherichia coli* was sensitive to amikacin (28), ampicillin and ciprofloxacin (19). *Klebsiella* spp. showed susceptibility to ceftriaxone (27), carbenicillin (27), amikacin (27,28), and imipenem (27). *P.*

aeruginosa and *Citrobacter koseri* also were susceptible to ciprofloxacin (19,28), while *Enterobacter cloacae* had susceptibility to azithromycin (19).

Responding to the third question of this scoping review, it was found that the antimicrobials to which the microorganisms studied presented greater resistance were penicillin (17,18,23-25,28-30), amoxicillin (12,21,26,27,29), erythromycin (24,28,30), and metronidazole (18,21,23,26).

All studies included in this review fully met at least 75% of the defined quality criteria (9), therefore, they were classified as of good quality (Table 2). Nevertheless, it is important to highlight that the studies included in this review presented great heterogeneity in their designs, reflected in the exploration of different classes of antimicrobials, great variability in the characteristics of the patients studied, and variability in the microbiological identification and the microorganisms studied, among other characteristics.

Discussion

To the best of the authors' knowledge, this scoping review is the first to consider the prevalence and proportions of antimicrobial-resistant species in patients with odontogenic infections. Whereas adjunctive antimicrobials are helpful implements in the therapy of some oral cavity infections, latent concerns occur regarding variations in the oral species as a consequence of their management (31,32). These issues involve the three questions proposed in this scoping review.

It is important to note that in this review, only studies containing phenotypic analysis were evaluated, understanding that the manifestation of antibiotic resistance genes does not essentially reveal the antimicrobial resistance of the microorganisms (33). On the other hand, only prospective studies and RCTs were included in this review, considering that retrospective studies are subject to bias including missing data, classification and interpretation bias in clinical records, and inconsistencies in treatment methods, among others (22).

Table 2: Quality of the selected studies (9).

Study	Fully met criteria	Score in the percentage of compliance
Umeshappa <i>et al.</i> 2021 (24)	14	87.5%
Uppada & Sinha 2020 (25)	14	87.5%
Sebastian <i>et al.</i> 2019 (26)	14	87.5%
Shah <i>et al.</i> 2016 (27)	14	87.5%
Shakia <i>et al.</i> 2018 (18)	12	75%
Jagadish <i>et al.</i> 2017 (19)	14	87.5%
Gómez-Arámbula <i>et al.</i> 2015 (10)	14	87.5%
Walia <i>et al.</i> 2014 (28)	14	87.5%
Fating <i>et al.</i> 2014 (29)	15	94%
Singh <i>et al.</i> 2014 (30)	14	87.5%
Sobottka <i>et al.</i> 2012 (11)	14	87.5%
Matijević <i>et al.</i> 2009 (12)	14	87.5%
Chardin <i>et al.</i> 2009 (13)	15	94%
Al-Nawas & Maeurer. 2008 (20)	14	87.5%
Boyanova <i>et al.</i> 2006 (21)	14	87.5%
Flynn <i>et al.</i> 2006 (22)	15	94%
Kuriyama <i>et al.</i> 2005 (23)	14	87.5%
Gilmore <i>et al.</i> 1998 (14)	14	87.5%
Lo <i>et al.</i> 1993 (15)	14	87.5%
Gorbach <i>et al.</i> 1991 (16)	12	75%
Quayle <i>et al.</i> 1987 (17)	12	75%

Although incision and drainage is the first treatment option for odontogenic infections, an adequate knowledge of the microorganisms involved in these infections, in addition to their susceptibility to antimicrobials will allow for establishing an adequate therapeutic regimen (1,19). On many occasions after adequate surgical therapy, patients do not improve. One of the relevant reasons is the presence of bacterial resistance and the selection of the inappropriate antimicrobial (24). Unfortunately, while awaiting laboratory results containing information on the identified microorganisms and their antimicrobial susceptibility, clinicians make an empirical selection of antimicrobials (23).

Microbiological samples from odontogenic infections are characterized by being constituted by a complexity of species, which can vary from aerobes and anaerobes to a mixture of aerobes and anaerobes (24). The proportion of these microorganisms varies between studies due to dissimilar techniques and resources implemented. Herein, regarding the prevalence of antimicrobial-resistant species, the most resistant bacteria were *Staphylococcus*, *Streptococcus* spp., *Peptostreptococcus* spp., *Bacteroides* spp., and *Prevotella* spp. Among aerobic and facultative anaerobe prevailed, *S. aureus* (12,18,19,25,28), *S. viridans* (15,17,26,27), *K. pneumoniae* (12,19,22,27),

S. milleri (22), *Enterococcus* spp. (25), *P. aeruginosa* (28), *P. mirabilis* (19), and *S. coagulans-negative* (12). In obligate anaerobes sampled were *Peptostreptococcus* spp. (12,15,17,18,20,21,23,24,26,30), *Bacteroides* spp. (12,14,18,21,24,26,28-30), and *Prevotella* spp. (11,18,20,21,23,26,30). These microorganisms were resistant to different antimicrobials in dissimilar proportions (Table 1).

Penicillin is the antimicrobial traditionally used for odontogenic infections. Unfortunately, and due to its widespread use, it has developed the appearance of resistant microorganisms (34). The device comprises beta-lactamase labor that has been validated in anaerobic Gram-negative bacilli. The occurrence of orofacial odontogenic infections including beta-lactamase generating microorganisms fluctuates from 13% to 39% (24). In this regard, it has been informed that the proportion of β -lactam penicillinase resistance in *S. aureus* samples from hospitals and dental clinics observed in 2020 are comparable to methicillin-resistant *S. aureus* percentages described in 2018 (34). Resistance to penicillin has also been frequent in anaerobes caused by the production of beta-lactamase (24).

Amoxicillin has also been one of the antimicrobials that have been prescribed empirically for the management of

odontogenic infections (26). Nonetheless, as found in this review, aerobic microorganisms (12,26), anaerobic bacteria (12,21,26), and mixed pathogens (12,26) presented resistance to this antibiotic. On the other hand, amoxicillin/clavulanic acid has shown good efficacy against aerobic and anaerobic microorganisms, showing superiority in activity to amoxicillin alone (11,20,24,27). The supplement with clavulanic acid broadens the spectrum against *Staphylococcus* spp. and other anaerobes by conceding beta-lactamase resistance (18). It has also been indicated that clindamycin may be an alternative in cases of inefficacy to amoxicillin (35). Clindamycin has good efficacy in aerobic Gram-positive cocci, including *S. aureus*, *Streptococcus* spp., and most anaerobes, counting penicillin-resistant species such as *Bacteroides* spp., *Prevotella* spp., and *Porphyromonas* (20,25,26). Furthermore, the efficacy of clindamycin and amoxicillin/clavulanic acid in odontogenic infections is comparable (36).

As found in this review, it is widely known that metronidazole does not show efficacy against aerobes (18,21,23), but it does against obligate anaerobes (20,23-26). The combination of metronidazole with penicillin has been recommended because it covers the microbial flora of odontogenic infections, compensating for the limited action of penicillin against beta-lactam anaerobes. The combination of amoxicillin/clavulanic acid plus metronidazole has also shown efficacy against strict anaerobes and facultative anaerobes (24).

First and second-generation cephalosporins have presented efficacy against aerobes and anaerobic Gram-positive cocci, corroborating the results of this scoping review (10,12,19,24,27-30). However, their efficacy against anaerobic Gram-negative rods is unpredictable. In this regard, it has been reported that cefotaxime (third-generation cephalosporin) has demonstrated in vitro efficacy against anaerobic bacteria of the mixed flora of odontogenic infections (24), also confirming the results described here (18,19).

Regarding the group of macrolides, it was observed high resistance to erythromycin (21,23,24,26,28,30); however, the efficacy of azithromycin for the treatment of odontogenic infections shows controversial results (15,19,26). While two studies described good efficacy against *Staphylococcus* spp. (15,19), other research informed high resistance to aerobic bacteria (80%), anaerobic microorganisms (78%), and 70% of mixed pathogens (26). These high values of resistance to macrolides have also been previously referenced (37).

Controversial susceptibility results were also observed in the quinolone group. Ciprofloxacin demonstrated a good efficacy against *S. aureus* (24,25,28), *Staphylococcus* spp., and *Streptococcus* spp. (19), Gram-negative microorganisms, *E. coli*, and *P. aeruginosa* (19). One RCT showed that 98% of pathogens (*S. viridans*, *Prevo-*

tella spp., *Neisseria* spp., *Streptococcus anginosus*, and other anaerobes) were susceptible to moxifloxacin (11). Similarly, another RCT indicated that anaerobes, *Streptococcus*, and *Staphylococcus* spp. showed low resistance to moxifloxacin (10), while one prospective study described that *S. viridans* and *K. pneumoniae* displayed intermediate susceptibility and high resistance to this antimicrobial, respectively (27). As has been described, methodological and geographical differences in research evaluating bacterial resistance may support these results (7,38).

Interestingly, in this review a considerable number of studies found a changing tendency in terms of the preponderance of anaerobic microorganisms over aerobic ones (14,20,21,24,30). Therefore, it has been recommended that prompt identification and careful management of odontogenic infections by surgical drainage and adjunct antimicrobials are essential to avoid the risk of expansion into adjoining fascial spaces (24). Thus, the elevated proportion of anaerobic microorganisms in the current review underlines the relevance of prospective studies in this field.

In short, different investigations recommend that the combination of amoxicillin plus clavulanic acid is the first line of antimicrobial selection, showing efficacy against most microorganisms involved in odontogenic infections (18,20,24,27,29,30). However, more prospective clinical studies and RCTs are required to evaluate antimicrobial resistance in patients with odontogenic infections in different parts of the world. In this regard, a review involving seven reports that assessed 374 patients from diverse nations worldwide, divulged that antimicrobial resistance frequencies varied rendering to the preceding utilization of antimicrobials (39).

The results described by this scoping review may support clinicians and leaders of public health organizations to create important decisions, as well as to obtain a better consciousness of the relevance of the reasonable management of antimicrobials.

Conclusions

In summary, most of the microorganisms had resistance to diverse groups of antimicrobials. Suitable antimicrobials must be prescribed founded on the microbial samples, culture susceptibility, and clinical progression of the odontogenic infections. Furthermore, it was observed high levels of resistance to antimicrobials that have been used in local and systemic therapy of oral cavity infections. An issue of concern is the preponderance of anaerobic microorganisms over aerobic ones.

References

1. Bahl R, Sandhu S, Singh K, Sahai N, Gupta M. Odontogenic infections: Microbiology and management. *Contemp Clin Dent*. 2014;5:307-11.
2. He D, Qian Y, Zhou L, Qi H, Liu Y. Multifunctional Irrigation-As-

- sisted Vacuum Drainage versus Traditional Drainage in the Treatment of Odontogenic Deep Fascial Infection: A Retrospective Cohort Study. *Infect Drug Resist.* 2021;14:3571-80.
3. Swift JQ, Gulden WS. Antibiotic therapy-managing odontogenic infections. *Dent Clin North Am.* 2002;46:623-33.
 4. Lockhart PB, Thornhill MH, Zhao J, Baddour LM, Gilbert GH, Mc-Knight PE, et al. National Dental PBRN Collaborative Group. Factors that affect dentists' use of antibiotic prophylaxis: Findings from a The National Dental Practice-Based Research Network questionnaire. *J Am Dent Assoc.* 2022;153:552-562.
 5. Karobari MI, Khijmatgar S, Bhandary R, Krishna Nayak US, Del Fabbro M, Horn R, et al. A Multicultural Demographic Study to Analyze Antibiotic Prescription Practices and the Need for Continuing Education in Dentistry. *Biomed Res Int.* 2021;2021:5599724.
 6. Prabhu SR, Nirmalkumar ES. Acute Fascial Space Infections of the Neck: 1034 cases in 17 years follow up. *Ann Maxillofac Surg.* 2019;9:118-23.
 7. Ardila CM, Bedoya-García JA. Antimicrobial resistance of Aggregatibacter actinomycetemcomitans, Porphyromonas gingivalis and Tannerella forsythia in periodontitis patients. *J Glob Antimicrob Resist.* 2020;22:215-218.
 8. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med.* 2018;169:467-73.
 9. Sirriyeh R, Lawton R, Gardner P, Armitage G. Reviewing studies with diverse designs: the development and evaluation of a new tool. *J Eval Clin Pract.* 2012;18:746-52.
 10. Gómez-Arámbula H, Hidalgo-Hurtado A, Rodríguez-Flores R, González-Amaro AM, Garrocho-Rangel A, Pozos-Guillén A. Moxifloxacin versus Clindamycin/Ceftriaxone in the management of odontogenic maxillofacial infectious processes: A preliminary, intrahospital, controlled clinical trial. *J Clin Exp Dent.* 2015;7:e634-9.
 11. Sobotta I, Wegscheider K, Balzer L, Böger RH, Hallier O, Giersdorf I, et al. Microbiological analysis of a prospective, randomized, double-blind trial comparing moxifloxacin and clindamycin in the treatment of odontogenic infiltrates and abscesses. *Antimicrob Agents Chemother.* 2012;56:2565-9.
 12. Matijević S, Lazić Z, Kuljić-Kapulica N, Nonković Z. Empirical antimicrobial therapy of acute dentoalveolar abscess. *Vojnosanit Pregl.* 2009;66:544-50.
 13. Chardin H, Yasukawa K, Nouacer N, Plainvert C, Aucouturier P, Ergani A, et al. Reduced susceptibility to amoxicillin of oral streptococci following amoxicillin exposure. *J Med Microbiol.* 2009;58:1092-7.
 14. Gilmore WC, Jacobus NV, Gorbach SL, Doku HC, Tally FP. A prospective double-blind evaluation of penicillin versus clindamycin in the treatment of odontogenic infections. *J Oral Maxillofac Surg.* 1988;46:1065-70.
 15. Lo Bue AM, Sammartino R, Chisari G, Gismondo MR, Nicoletti G. Efficacy of azithromycin compared with spiramycin in the treatment of odontogenic infections. *J Antimicrob Chemother.* 1993;31:119-27.
 16. Gorbach SL, Gilmore WC, Jacobus NV, Doku HC, Tally FP. Microbiology and antibiotic resistance in odontogenic infections. *Ann Otol Rhinol Laryngol Suppl.* 1991;154:40-2.
 17. Quayle AA, Russell C, Hearn B. Organisms isolated from severe odontogenic soft tissue infections: their sensitivities to cefotetan and seven other antibiotics, and implications for therapy and prophylaxis. *Br J Oral Maxillofac Surg.* 1987;25:34-44.
 18. Shakya N, Sharma D, Newaskar V, Agrawal D, Shrivastava S, Yadav R. Epidemiology, Microbiology and Antibiotic Sensitivity of Odontogenic Space Infections in Central India. *J Maxillofac Oral Surg.* 2018;17:324-31.
 19. Jagadish Chandra H, Sripathi Rao BH, Muhammed Manzoor AP, Arun AB. Characterization and Antibiotic Sensitivity Profile of Bacteria in Orofacial Abscesses of Odontogenic Origin. *J Maxillofac Oral Surg.* 2017;16:445-52.
 20. Al-Nawas B, Maeurer M. Severe versus local odontogenic bacterial infections: comparison of microbial isolates. *Eur Surg Res.* 2008;40:220-4.
 21. Boyanova L, Kolarov R, Gergova G, Deliverska E, Madjarov J, Marinov M, et al. Anaerobic bacteria in 118 patients with deep-space head and neck infections from the University Hospital of Maxillofacial Surgery, Sofia, Bulgaria. *J Med Microbiol.* 2006;55:1285-9.
 22. Flynn TR, Shanti RM, Levi MH, Adamo AK, Kraut RA, Trieger N. Severe odontogenic infections, part I: prospective report. *J Oral Maxillofac Surg.* 2006;64:1093-103.
 23. Kuriyama T, Absi EG, Williams DW, Lewis MA. An outcome audit of the treatment of acute dentoalveolar infection: impact of penicillin resistance. *Br Dent J.* 2005;198:759-63.
 24. Umeshappa H, Shetty A, Kavati K, Vivek GK, Vaibhav N, Mohammed I. Microbiological profile of aerobic and anaerobic bacteria and its clinical significance in antibiotic sensitivity of odontogenic space infection: A prospective study of 5 years. *Natl J Maxillofac Surg.* 2021;12:372-9.
 25. Uppada UK, Sinha R. Outcome of Odontogenic Infections in Rural Setup: Our Experience in Management. *J Maxillofac Oral Surg.* 2020;19:113-8.
 26. Sebastian A, Antony PG, Jose M, Babu A, Sebastian J, Kunnilathu A. Institutional microbial analysis of odontogenic infections and their empirical antibiotic sensitivity. *J Oral Biol Craniofac Res.* 2019;9:133-8.
 27. Shah A, Ramola V, Nautiyal V. Aerobic microbiology and culture sensitivity of head and neck space infection of odontogenic origin. *Natl J Maxillofac Surg.* 2016;7:56-61.
 28. Walia IS, Borle RM, Mehendiratta D, Yadav AO. Microbiology and antibiotic sensitivity of head and neck space infections of odontogenic origin. *J Maxillofac Oral Surg.* 2014;13:16-21.
 29. Fating NS, Saikrishna D, Vijay Kumar GS, Shetty SK, Raghavendra Rao M. Detection of Bacterial Flora in Orofacial Space Infections and Their Antibiotic Sensitivity Profile. *J Maxillofac Oral Surg.* 2014;13:525-32.
 30. Singh M, Kambalimath DH, Gupta KC. Management of odontogenic space infection with microbiology study. *J Maxillofac Oral Surg.* 2014;13:133-9.
 31. Feres M, Haffajee AD, Allard K, Som S, Goodson JM, Socransky SS. Antibiotic resistance of subgingival species during and after antibiotic therapy. *J Clin Periodontol.* 2002;29:724-375.
 32. Ardila CM, Bedoya-García JA. Clinical and Microbiological Efficacy of Adjunctive Systemic Quinolones to Mechanical Therapy in Periodontitis: A Systematic Review of the Literature. *Int J Dent.* 2022;2022:4334269.
 33. Jungermann GB, Burns K, Nandakumar R, Tolba M, Venezia RA, Fouad AF. Antibiotic resistance in primary and persistent endodontic infections. *J Endod.* 2011;37:1337-44.
 34. Meinen A, Reuss A, Willrich N, Feig M, Noll I, Eckmanns T, et al. Antimicrobial Resistance and the Spectrum of Pathogens in Dental and Oral-Maxillofacial Infections in Hospitals and Dental Practices in Germany. *Front Microbiol.* 2021;12:676108.
 35. von Konow L, Köndell PA, Nord CE, Heimdahl A. Clindamycin versus phenoxymethylpenicillin in the treatment of acute orofacial infections. *Eur J Clin Microbiol Infect Dis.* 1992;11:1129-35.
 36. Tancawan AL, Pato MN, Abidin KZ, Asari AS, Thong TX, Kochhar P, et al. Amoxicillin/Clavulanic Acid for the Treatment of Odontogenic Infections: A Randomised Study Comparing Efficacy and Tolerability versus Clindamycin. *Int J Dent.* 2015;2015:472470.
 37. Poeschl PW, Spusta L, Russmueller G, Seemann R, Hirschl A, Poeschl E, et al. Antibiotic susceptibility and resistance of the odontogenic microbiological spectrum and its clinical impact on severe deep space head and neck infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2010;110:151-6.
 38. Helstad AG, Kimball JL, Maki DG. Recovery of anaerobic, facultative, and aerobic bacteria from clinical specimens in three anaerobic transport systems. *J Clin Microbiol.* 1977;5:564-9.
 39. Lang PM, Jacinto RC, Dal Pizzol TS, Ferreira MB, Montagner F. Resistance profiles to antimicrobial agents in bacteria isolated from acute endodontic infections: systematic review and meta-analysis. *Int J Antimicrob Agents.* 2016;48:467-74.

Acknowledgments

To the School of Dentistry, Universidad de Antioquia, Medellin, Colombia.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authorship

Both authors have made substantial contributions to the conception and design of the study, acquisition of data, analysis and interpretation of data, drafting the article and revising it critically for important intellectual content, and final approval of the version to be submitted.

Data Availability

Records were obtained from the included investigations.

Conflict of interest

None.