

Journal section: Medically compromised patients in Dentistry
 Publication Types: Review

doi:10.4317/medoral.21862
<http://dx.doi.org/doi:10.4317/medoral.21862>

New horizons in anticoagulation: Direct oral anticoagulants and their implications in oral surgery

Víctor Serrano-Sánchez¹, Jorge Ripollés-de Ramón², Luis Collado-Yurrita³, Iris Vaello-Checa¹, Constantino Colmenero-Ruiz⁴, Alexandra Helm⁵, Maria-José Ciudad-Cabañas⁶, Victoriano Serrano-Cuenca⁷

¹ DDS. Collaborator in the Department of Medicine at the Faculty of Medicine. Faculty of Dentistry of Complutense University of Madrid

² DDS, PhD. Collaborator in Department of Oral Medicine & Orofacial Surgery. Faculty of Dentistry of Complutense University of Madrid. Collaborate Professor at the Master of Oral Surgery and Implantology UC

³ Director of the Department of Medicine. Faculty of Medicine of Complutense University of Madrid

⁴ DDS. Faculty of Dentistry of Complutense University of Madrid. Collaborator in the Master in Oral Surgery at the University Hospital of Príncipe de Asturias (UAH)

⁵ DDS. Faculty of Dentistry of Complutense University of Madrid

⁶ Department of Medicine. Faculty of Medicine of Complutense University of Madrid

⁷ MD, DDS, PHD. Professor of Department of Oral Medicine & Orofacial Surgery. Faculty of Dentistry of Complutense University of Madrid

Correspondence:

C/ Profesor Waksman 13 2º Izda
 28036 Madrid
 Vicser02@ucm.es

Serrano-Sánchez V, Ripollés-de Ramón J, Collado-Yurrita L, Vaello-Checa I, Colmenero-Ruiz C, Helm A, Ciudad-Cabañas MJ, Serrano-Cuenca V. New horizons in anticoagulation: Direct oral anticoagulants and their implications in oral surgery. Med Oral Patol Oral Cir Bucal. 2017 Sep 1;22 (5):e601-8.

<http://www.medicinaoral.com/medoralfree01/v22i5/medoralv22i5p601.pdf>

Received: 14/02/2017
 Accepted: 27/05/2017

Article Number: 21862 <http://www.medicinaoral.com/>
 © Medicina Oral S. L. C.I.F. B 96689336 - pISSN 1698-4447 - eISSN: 1698-6946
 eMail: medicina@medicinaoral.com
Indexed in:
 Science Citation Index Expanded
 Journal Citation Reports
 Index Medicus, MEDLINE, PubMed
 Scopus, Embase and Emcare
 Índice Médico Español

Abstract

Background: Thrombotic disorders remain a leading cause of death in the Western World. For decades, vitamin K antagonists used in the prevention of this pathology, such as warfarin or sintrom, were the only oral agents available for long-term anticoagulation, in spite of their disadvantages.

Material and Methods: An electronic database search was carried out on MedLine and The Cochrane Library Plus, without restrictions on the type of study nor dates, in English and Spanish. Abstracts were reviewed, and complete articles if necessary, considering all articles that included recommendations on DOACs and oral surgery.

Results: In recent years, the so-called “new oral anticoagulants” have been introduced in clinical practice to treat those patients whose medical conditions require long-term anticoagulant treatment, replacing traditional oral anticoagulants.

Conclusions: The new oral anticoagulants represent new therapeutic options, with a number of advantages such as poor interaction with food, minor drug interactions, and do not require periodic dose adjustments or routine controls. The purpose of this review is to establish an update on the new oral anticoagulants: Dabigatran, Rivaroxaban, Apixaban and Edoxaban.

Key words: Novel oral anticoagulants, Dabigatran, Rivaroxaban, Apixaban, Edoxaban, bleeding management, oral surgery, Anti-IIa, Anti Xa.

Introduction

Atrial fibrillation (AF) is the most frequent sustained arrhythmia in humans, affecting 1-2% of the world population. It affects 3 to 6 million people in the United States (1), whereas in Europe, it is estimated that in 2010, 8.8 million adults over the age of 55 suffered from this arrhythmia, and these figures are expected to double by 2060 (2).

The prevalence of AF increases with age, where studies report varying estimates ranging from 2% in people under 80 years of age, to 5-15% in those older than 80 (3). Therefore, AF represents a modern-day epidemic that we must face in our daily clinical practice.

For the past decades, patients with atrial fibrillation or venous thromboembolism have been managed exclusively with vitamin K antagonists. However, they possess a Narrow Therapeutic Index (NTI), where slight changes in plasma levels may lead to treatment failure (subtherapeutic concentrations) or to various adverse effects (supra-therapeutic concentrations) such as an excessive bleeding risk.

As described above, these types of drugs have some limitations, such as ongoing monitoring and dosage adjustment. In addition, there are a number of other drawbacks, such as drug and food interactions, and even with viral diseases, as well as a relatively slow onset of action (4).

Therefore, over the past few years the so-called "new oral anticoagulants", or "direct oral anticoagulants" (DOAC) were developed. These drugs have been introduced in clinical practice to treat various diseases and medical conditions that require the use of extended anticoagulant drug therapy, such as the prophylaxis and treatment of pulmonary and venous thromboembolism, including thromboprophylaxis following orthopedic surgery; prophylaxis and treatment of thromboembolic complications associated with atrial fibrillation and/or prosthetic replacement of cardiac valves; reducing the risk of death and new thromboembolic events such as stroke or reinfarction (5).

Compared with the coumarin derivatives described above, these drugs have very specific targets in the coagulation cascade. There are currently three DOACs approved for use in the US and in several European countries, such as dabigatran etexilate (direct thrombin inhibitor), rivaroxaban and apixaban, all factor Xa inhibitors.

In addition to the three previously mentioned drugs, a fourth drug belonging to the group of factor Xa inhibitors, edoxaban, has been recently approved by the European Medicines Agency.

Material and Methods

An electronic database search was carried out on MedLine and The Cochrane Library Plus, without restric-

tions on the type of study nor dates, in English and Spanish. Abstracts were reviewed, and complete articles if necessary, considering all articles that included recommendations on DOACs and oral surgery.

New oral anticoagulants (noac)

•Dabigatran

Mechanism of action

Dabigatran etexilate (Pradaxa®, Boehringer Ingelheim, Spain) is the first direct and reversible thrombin oral anticoagulant approved for use. It is a pro-drug that is rapidly converted, via plasma and hepatic esterases, to dabigatran. Dabigatran is a potent, direct, competitive inhibitor of thrombin (IIa Factor). Since thrombin allows the conversion of fibrinogen to fibrin in the coagulation cascade, its inhibition prevents thrombus formation. (6)

RE-LY study

The efficacy of dabigatran was first assessed by the RE-LY (7) study, a randomized, multicenter trial designed to compare two fixed doses of dabigatran with warfarin in patients who had AF and were at increased risk for stroke.

Following the results presented by the RE-LY study, we can conclude that, compared with warfarin, dabigatran administered at a dose of 110 mg has the same therapeutic effect as warfarin, but with a lower bleeding rate, whereas dabigatran administered at a dose of 150 mg has a greater therapeutic effect than warfarin in the prevention of stroke and thromboembolism, at a similar bleeding rate (8).

-Rivaroxaban

Mechanism of action

Rivaroxaban (Xarelto®, Bayer HealthCare and Johnson & Johnson Pharmaceutical Research & Development) is an oxazoline derivative, and is the first oral direct activated factor X (Xa) inhibitor to be approved for use. It is a selective, reversible and direct factor Xa inhibitor, both free and bound to the prothrombinase complex, which interferes in both the intrinsic and extrinsic pathways of the coagulation cascade (9).

ROCKET AF Study

The efficacy of rivaroxaban compared with warfarin was assessed in the ROCKET AF (10) study, a multicenter, double-blind, randomized trial. It was concluded that rivaroxaban was noninferior to warfarin in the prevention of stroke or thromboembolism, but presented no differences in bleeding risk, although intracranial haemorrhage and fatal bleeding were reported to a lesser extent in the rivaroxaban group (11).

•Apixaban

Mechanism of action

Apixaban (Eliquis®, Bristol-Myers Squibb) is a potent, reversible, direct and highly selective factor Xa inhibitor. Apixaban inhibits free and clot-bound factor Xa, as

well as prothrombinase activity. By inhibiting factor Xa, both thrombin formation and thrombus formation is impeded. (12)

ARISTOTLE study

The efficacy of apixaban against warfarin was assessed in the ARISTOTLE study (13), a multicenter, double-blind, randomized study. It was concluded that warfarin is highly effective in terms of stroke prevention in patients with AF, but is associated with a variable response, has drug and food interactions, requires regular monitoring and carries a risk of bleeding (including intracranial hemorrhage).

Alternative therapy with apixaban does not require monitoring and is not only more effective than warfarin in the prevention of stroke, but also provides a lower bleeding risk. In summary, in patients with AF, apixaban was superior to warfarin in the prevention of stroke and thromboembolism, and causes less bleeding, resulting in a lower death rate (14).

•Edoxaban

Mechanism of action

Edoxaban (Lixiana ® Bristol-Myers Sqibb) is a highly selective, direct and reversible factor Xa inhibitor. Edoxaban inhibits both free factor Xa and prothrombinase activity. Inhibition of factor Xa in the coagulation cascade reduces thrombin production, extends coagulation time, and reduces the risk of thrombus formation (15).

ENGAGE AF-TIMI 48 study

The efficacy of edoxaban against warfarin was assessed in the ENGAGE AF-TIMI 48 study (16). It is a multicenter, double-blind, randomized study.

It was concluded that edoxaban was noninferior to warfarin in the prevention of stroke or thromboembolism, along with a significantly lower associated risk of hemorrhage or death from cardiovascular causes (17), (Table 1).

Doac interactions

DOACs are also susceptible to drug interactions. Some of the recognized interactions with the new anticoagulants are shown in table 2.

Heparins, thrombolytic and antiplatelet agents potenti-

ate their activity and may alter haemostasis, increasing bleeding risk. All oral anticoagulants interact with each other potentiating their effects, therefore, in their management and modification, the duration of their effect must be considered (18,19) (Table 2).

Clinical approach to DOACs

There are particular three areas of uncertainty, such as laboratory tests to monitor the degree of anticoagulation, preoperative management of the patient and management of bleeding.

-Monitoring

Although routine monitoring is not required, in certain situations it may be useful to assess the degree of anticoagulation.

Routine coagulation tests, such as INR and activated partial thromboplastin time (APTT) do not accurately reproduce the degree of anticoagulation in patients treated with DOAC. Therefore, INR is not used to monitor patients treated with direct anticoagulants, whereas APTT is increased in patients taking dabigatran (and more recently proven in patients treated with edoxaban), but the relationship between this increase and the received dose is yet unclear (20).

Specialized tests, including thrombin time (TT), dilute thrombin time (TTd), and ecarin time (TE) are some of the alternatives that are discussed for DOAC monitoring.

TT is a very accurate test in the assessment of dabigatran activity, and may be useful in determining when the drug has been completely cleared prior to highly aggressive surgery (21). Furthermore, TTd and TE provide the dose-response ratio that could not be determined by APTT, but these tests are not yet available in most institutions because of their high cost. Finally, Hemoclot is a TTd variant specifically calibrated for dabigatran, being a good way to identify those patients with an increased risk of bleeding (T > 60 s), which is carried out exclusively in cases of severe bleeding or urgent surgery (22).

As for rivaroxaban and apixaban, as with dabigatran, INR is also not valid, although TP and APTT can be used (through specific calibration curves). TT and TE

Table 1. Summary of the main properties of NOACs.

	DABIGATRAN	RIVAROXABAN	APIXABAN	EDOXYBAN
Trade name	Pradaxa ®	Xarelto ®	Eliquis ®	Lixiana ®
Mechanism of action	Anti IIa	Anti Xa	Anti Xa	Anti Xa
Time to peak plasma concentration	1 to 1.5 hours	2 to 4 horas	1 to 3 horas	1 to 2 horas
Hepatic metabolism	20%	68%	74%	50%
Renal elimination	80%	35%	25%	50%
Half life	12 to 17 hours	5 to 13 hours	8 to 15 hours	10 to 14 hours

Table 2. Interactions of NOACs with other drugs.

NOACs	DRUGS AND TYPE OF INTERACTION
DABIGATRAN	POTENTIATES: ASA, NSAIDS, Amiodarone, Ciclosporin, Clarithromycin, Clopidogrel, Ketoconazole, Quinidine, Thrombolytics. INHIBITS: Carbamazepine, Dexamethasone, Estrogens Phenytoin, Phenobarbital, St. John’s wort
RIVAROXABAN	POTENTIATES: ASA, Fusidic acid, NSAIDS, Clarithromycin, Clopidogrel, Chloramphenicol, Erythromycin, Ketoconazole, Thrombolytics INHIBITS: Carbamazepine, Dexamethasone, Estrogens, Phenytoin, Phenobarbital, Primidone, St. John’s wort
APIXABÁN	POTENTIATES: ASA, Fusidic acid, NSAIDS, Clopidogrel, Chloramphenicol, Ketoconazole, Nicardipine, Thrombolytics INHIBITS: Carbamazepine, Dexamethasone, Estrogens, Phenytoin, Phenobarbital, Primidone, St. John’s wort
EDOXABÁN	POTENCIAN: ASA, NSAIDS, Verapamil, Quinidine, Ketoconazole, Amiodarone, Dronedarone, Thrombolytics. INHIBEN: Rifampicina, St. John’s wort, Carbamazepine, Phenytoin, Phenobarbital.

are not useful since thrombin is not affected. A specific test that can be used in the case of factor Xa inhibitors is Heptest, which measures anti-Xa activity. Finally, in case of emergency, chromogenic methods of measuring anti-Xa activity can be used, which provide a linear dose-effect ratio (23). In the case of edoxaban, INR is not valid; TP and APTT are prolonged, but without a known link to bleeding risk; TTd is not effective; TE is unaffected, and chromogenic analysis of anti-Xa activity provides us with only quantitative data, since there are no threshold values for hemorrhage or thrombosis.

-Preoperative management

In general, interventions with a low bleeding risk can be safely carried out without the interruption of DOAC (20). In the case of surgeries involving a moderate or severe bleeding risk, it should be weighed against the risk of thromboembolism, where the decision to suspend or delay DOAC should be individualized.

In order to objectify the risk of thromboembolism, the CHADS2 criterion (24), which classifies the general population depending on stroke risk, has been in place since 2001 (Table 3).

However, this criterion has been modified to improve discrimination between low and moderate risk populations. Thus, a new criterion emerges in 2010, named CHA2DS2-VASc (25) (Table 4), in which three more variables are included, such as the female sex, age range from 65 to 74 years, and cardiovascular events.

Once the risk of thromboembolism has been established, the risk of bleeding should be determined. Therefore, the HAS-BLED scale has been proposed (26) (Table 4). Once the risk of suffering a thromboembolic event and bleeding has been determined, the following guideline is generically proposed: (Fig. 1).

Discontinuation of medication prior to surgery is determined by factors such as drug half-life, specific inter-

Table 3: CHADS₂ score for stroke risk.

CHADS ₂	Criteria	Score
C (Congestive heart failure)	Recent congestive heart failure history	1
H (Hypertension)	HT history	1
A (Age)	Age > 75 años	1
D (Diabetes Mellitus)	Diabetes mellitus history	1
S ₂ (Stroke)	Stroke/TIA history	2
Maximum score		6

Table 4: CHA₂DS₂VAS_C and HAS-BLED Scores

Risk-based approach expressed as a scoring system with the acronym CHA ₂ DS ₂ VAS _C		Clinical characteristics of the HAS-BLED bleeding risk score		
Risk factor	Score	Letter	Clinical characteristic	Score
CHF or LVD	1	H	Hipertensión	1
Hypertension	1	A	Abnormal renal and liver function (one point each)	1 or 2
Age ≥ 75 years	2	S	ACV	1
Diabetes Mellitus	1	B	Bleeding	1
Stroke, TIA or thromboembolism	2	L	Labile INR	1
Vascular disease	1	E	Elderly age (>65 years)	1
Age 65-74 years	1	D	Drug therapy or alcohol intake (one point each)	1 or 2
Sex (Female)	1	Maximum score		9
Maximum score	9			




Fig. 1: Action protocol based on bleeding and thrombotic risk.
 • From González Fernández-Tresguerres F. *et al.*

vention bleeding risks, and the patient’s renal function. In general, factor Xa inhibitors should be discontinued at least 24-48 hours prior to intervention in cases of moderate bleeding risk, and 48-72 hours in cases of high risk. Dabigatran should be discontinued at least 72 hours in advance. In the case of renal dysfunction, these times should be prolonged.

Medication shall be resumed once a correct haemostasis has been achieved. In procedures with a moderate risk, it will be resumed 24 hours following the intervention, provided that correct haemostasis is achieved. In surgeries that carry a higher risk, it will be resumed 48 hours afterwards. Unlike warfarin, the onset of these

drugs is very rapid, within a few hours of administration, and is therefore considered a lower risk therapy compared with vitamin K antagonists (20,27-29).

-Preoperative management in oral surgery

The preoperative management of the patient to be surgically intervened in the oral cavity follows the previously described guidelines for direct oral anticoagulation (30-34) (Fig. 1).

Traditionally, oral surgery has been considered a type of surgery involving a low bleeding risk. However, there are certain situations and interventions that imply an increased risk of bleeding, and may be considered as a moderate and even high risk.

In this manner, we can generically categorise dental treatments within three groups, depending on their relative bleeding risk: (30).

- Low risk: local anaesthetic infiltration, single dental extraction, soft tissue biopsy less than 1 cm in size, supragingival prophylaxis, placement of rubber dam, restorative procedures, crown preparation, root canal therapy, prosthetic rehabilitation of implants, and removal of orthodontic brackets and bands.
- Moderate risk: local anaesthesia nerve block, multiple simple extractions (less than 5 teeth), soft tissue biopsies ranging from 1 to 2.5 cm in size, placement of single implants, infragingival prophylaxis (6-12 teeth), and localized gingival surgery (less than 5 teeth).
- High risk: Multiple extractions of more than 5 teeth, surgical extractions requiring raising of mucoperiosteal flap and bone removal, soft tissue biopsies larger than 2.5cm in size, osseous biopsies, removal of torus, placement of multiple implants, complete periodontal treatment of all the oral cavity, gingival surgery of more than 5 teeth, endodontic surgery which involves osseous manipulation, removal of cysts and tumours, and bone regeneration procedures that involve major surgeries.

This classification serves purely as a guideline and should be individualized according to the conditions and pathologies of each patient, as well as the specific characteristics of the intervention, for which we can use the previously described CHA₂DS₂VASC and HAS-BLED scales.

Once the thromboembolic and haemorrhagic risk has been assessed, a decision will be made to maintain or

discontinue anticoagulant therapy (no scoring system can replace the clinician’s clinical judgment), where a consultation with the haematologist or cardiologist responsible for the patient’s medication should always be performed.

-Bleeding management

Regarding the management of a bleeding event, it is important to assess whether it is life-threatening (Fig. 2). To address these events, several strategies have been proposed, such as:

- Altering the pharmacokinetics of DOACs by reducing their absorption, or carrying out their elimination. In the case of dabigatran and apixaban, active carbon can be administered (35). There are no specific studies on its usefulness in the case of rivaroxaban and edoxaban, although it is believed that it may also be useful. In addition, in the case of dabigatran, hemodialysis could be performed, due to its low percentage of binding to plasma proteins (36).
- Antifibrinolytic agents, such as tranexamic acid and aminocaproic acid. Their peri-procedural use have proven a better haemostasis, presenting a higher safety profile than aprotinin (37).
- Plasma Factor Therapy: concentrated prothrombin complexes (CCPs), fresh frozen plasma and cryoprecipitates. The use of these factors may be helpful in the case of life-threatening bleeding, although more studies are needed to support this (38).
- Specific antidotes for each DOAC: Ciraparantag (Phase II), Andexanet Alfa (Phase II) and Idarucizumab. Currently, idarucizumab (Praxbind ®), approved




Fig. 2: Action protocol based on bleeding risk.

at the end of 2015, is the only antidote to have been approved for use in the European Union, making dabigatran the first and only DOAC to have a reversing agent (38) (Fig. 2).

Regarding the measures that we must adopt before surgical interventions on the oral cavity to prevent bleeding, the following should be highlighted:

- Unless contraindicated, a local anaesthetic such as 2% lidocaine will be used with 1: 80,000 or 1:100,000 epinephrine.
- Osteotomy must be reduced to the minimum during extractions, where tooth sectioning may be indicated instead.
- Mucoperiosteal flaps should be raised as atraumatically as possible, avoiding the dissection of planes that favour a pathway for hematomas to break through.
- Thorough curettage of the cavity is essential to avoid secondary infections (an important cause of postoperative bleeding).
- As for sutures, resorbable polyglactin 910 (Vicryl ®) is preferred.
- The use of gelatin sponges, thrombin, collagen (either synthetic or porcine), and oxycellulose. Oxycellulose soaked in tranexamic acid (Kin Exogel ®) has been especially useful.

Conclusions

- DOACs offer a safe and effective therapeutic alternative to traditional oral anticoagulants.
 - Most bleeding episodes are easily controllable with local haemostatic measures.
 - Before performing a dental procedure that involves the risk of associated bleeding, it is necessary to agree with the physician on the best course of action (withdrawal / non-withdrawal, or postponement of daily dose).
- The authors declare no conflicts of interest.

References

1. Go AS, Hylek EM, Phillips KA, Chang Y, Henault LE, Selby JV, et al. Prevalence of diagnosed atrial fibrillation in adults: National implications for rhythm management and stroke prevention: The anticoagulation and risk factors in atrial fibrillation (ATRIA) study. *JAMA*. 2001;285:2370-5.
2. Krijthe BP, Kunst A, Benjamin EJ, Lip GYH, Franco OH, Hofman A, et al. Projections on the number of individuals with atrial fibrillation in the European Union, from 2000 to 2060. *Eur Heart J*. 2013;34:2746-51.
3. Wynn GJ, Todd DM, Webber M, Bonnett L, McShane J, Kirchhof P, et al. The European Heart Rhythm Association symptom classification for atrial fibrillation: validation and improvement through a simple modification. *Europace*. The Oxford University Press. 2014;16:965-72.
4. Imberti D, Gallerani M, Manfredini R. Therapeutic potential of apixaban in the prevention of venous thromboembolism in patients undergoing total knee replacement surgery. *J Thromb Thrombolys*. 2012;34:208-13.
5. Firriolo FJ, Hupp WS. Beyond warfarin: the new generation of oral anticoagulants and their implications for the management of dental patients. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2012;113:431-41.

6. Levy JH, Key NS, Azran MS. Novel oral anticoagulants: implications in the perioperative setting. *Anesthesiology*. 2010;113:726-45.
7. Ezekowitz MD, Connolly S, Parekh A, Reilly PA, Varrone J, Wang S, et al. Rationale and design of RE-LY: randomized evaluation of long-term anticoagulant therapy, warfarin, compared with dabigatran. *Am Heart J*. 2009;157:805-10.
8. Connolly SJ, Ezekowitz MD, Yusuf S, Eikelboom J, Oldgren J, Parekh A, et al. Dabigatran versus Warfarin in patients with Atrial Fibrillation. *N Engl J Med*. 2009;361:1139-51.
9. ROCKET AF Study Investigators. Rivaroxaban — once daily, oral, direct factor Xa inhibition compared with vitamin K antagonism for prevention of stroke and Embolism Trial in Atrial Fibrillation: rationale and design of the ROCKET AF study. *Am Heart J*. 2010;159:340 .e1-347. e1.
10. Patel MR, Mahaffey KW, Garg J, Pan G, Singer DE, Hacke W, et al. Rivaroxaban versus Warfarin in Nonvalvular Atrial Fibrillation. *N Engl J Med*. 2011;365:883-91.
11. Connolly SJ, Eikelboom J, Joyner C, Diener HC, Hart R, Golitsyn S, et al. Apixaban in patients with atrial fibrillation. *N Engl J Med*. 2011;364:806-17.
12. Lopes RD, Alexander JH, Al-Khatib SM, Ansell J, Diaz R, Easton JD et al. Apixaban for reduction in stroke and other Thromboembolic events in atrial fibrillation (ARISTOTLE) trial: design and rationale. *Am Heart J*. 2010;159:331-9.
13. Granger CB, Alexander JH, McMurray JJV, Lopes RD, Hylek EM, Hanna M, et al. Apixaban versus Warfarin in patients with Atrial Fibrillation. *N Eng J Med*. 2011;365:981-92.
14. Matsushima N, Lee F, Sato T, Weiss D, Mendell J. Bioavailability and safety of the factor Xa inhibitor edoxaban and the effects of quinidine in healthy subjects. *Clin Pharm Drug Dev*. 2013;2:358-66.
15. Ruff CT, Giugliano RP, Antman EM, Crugnale Se, Bocanegra T, Mercuri M, et al. Evaluation of the novel factor Xa inhibitor edoxaban compared with warfarin in patients with atrial fibrillation: design and rationale for the Effective aNticoagulation with factor xA next GEneration in Atrial Fibrillation-Thrombolysis In Myocardial Infarction study 48 (ENGAGE AF-TIMI 48). *Am Heart J*. 2010;160:635-41.
16. Giugliano RP, Ruff CT, Braunwald E, Murphy SA, Wiviott SD, Halperin JL, et al. Edoxaban versus Warfarin in Patients with Atrial Fibrillation. *N Eng J Med*. 2013;269:2093-104.
17. Arcelus JJ, Cairols M, Granero X, Jiménez D, Llau JV, Monreal M, et al. Nuevos anticoagulantes orales: una visión multidisciplinaria. *Med Clin (Barc)*. 2009;133:508-12.
18. Heidbuchel H, Verhamme P, Alings M, Antz M, Hacke W, Oldgren J, et al. European Heart Rhythm Association Practical Guide on the use of new oral anticoagulants in patients with non-valvular atrial fibrillation. *Europace*. 2013;15:625-51.
19. Weitz JI, Pollack CV Jr. Practical management of bleeding in patients receiving non-vitamin K antagonist oral anticoagulants. *Thromb Haemost*. 2015;114:1113-26.
20. Wysokinski WE, McBane R D. Periprocedural bridging management of anticoagulation. *Circulation*. 2012;126:486-90.
21. Du S, Weiss C, Christina G, Krämer S, Wehling M, Krämer R, et al. Determination of dabigatran in plasma, serum, and urine samples: comparison of six methods. *Clin Chem Lab Med*. 2015;53:1237-47.
22. Harenberg J, Du S, Wehling M, Zolfaghari S, Weiss C, Krämer R, et al. Measurement of dabigatran, rivaroxaban and apixaban in samples of plasma, serum and urine, under real life conditions. An international study. *Clin Chem Lab Med*. 2016;54:275-83.
23. Barrios V, Calderón A, Escobar C, De la Figuera M. Pacientes con fibrilación auricular asistidos en consultas de atención primaria. *Estudio Val-FAAP*. *Rev Esp Cardiol*. 2012;65:47-53.
24. Chao T, Lin YJ, Tsao HM, Tsai CF, Lin WS, Chang SL, et al. CHADS2 and CHA2DS2-VASc Scores in the Prediction of Clinical Outcomes in Patients With Atrial Fibrillation After Catheter Ablation. *J Am Coll Cardiol*. 2011;58:2380-5.
25. Pisters R, Lane DA, Nieuwlaat R, De Vos RNC, Crijns HJG, Lip GYH. A Novel User-Friendly Score (HAS-BLED) To Assess 1-Year Risk of Major Bleeding in Patients With Atrial Fibrillation. *Chest*. 2010;138:1093-100.

26. Ment J. Direct oral anticoagulants: key considerations for use to prevent stroke in patients with nonvalvular atrial brillation. *Vasc Health Risk Manag.* 2015;11:317-32.
27. Ebright J, Mousa SA. Oral anticoagulants and status of antidotes for the reversal of bleeding risk. *Clin Appl Thromb Hemost.* 2015;21:105-14.
28. Liotta EM, Levasseur-Franklin KE, Naidech AM. Reversal of the novel oral anticoagulants dabigatran, rivoraxaban, and apixaban. *Curr Opin Crit Care.* 2015;21:127-33.
29. Constantinides F, Rizzo R, Pascazio L, Maglione M. Managing patients taking novel oral anticoagulants (NOAs) in dentistry: a discussion paper on clinical implications. *Oral Health.* 2016;16:5.
30. Elad S, Marshall J, Meyerowitz C, Connolly G. Novel anticoagulants: general overview and practical considerations for dental practitioners. *Oral Diseases.* 2016;22:23-32.
31. González Fernández-Tresguerres F, Álvarez Sirvent A, Torres García-Denche J, Fernández-Tresguerres Hernández-Gil I. Nuevos anticoagulantes orales: repercusión odontológica. *Cient Dent.* 2016;13;2:139-48.
32. Mauprivez C, Khonsari R, Razouk O, Goudot P, Lesclous P, Descroix V. Management of dental extraction in patients undergoing anticoagulant oral direct treatment. A pilot study. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2016;122:e146-e155.
33. Hong CH, Islam I. AntiThrombotic Therapy: Implications for Invasive Outpatient Procedures in Dentistry. *J Blood Disorders Transf.* 2013;4:166.
34. Wang X, Mondal S, Wang J, Tirucherai G, Zhang D, Boyd RA, et al. Effect of activated charcoal on apixaban pharmacokinetics in healthy subjects. *Am J Cardiovasc Drugs.* 2014;14:147-54.
35. Crowther M, Crowther MA. Antidotes for novel oral anticoagulants: current status and future potential. *Arterioscler Thromb Vasc Biol.* 2015;35:1736-45.
36. Weitz JI, Pollack CV Jr. Practical management of bleeding in patients receiving non-vitamin K antagonist oral anticoagulants. *Thromb Haemost.* 2015;114:1113-26.
37. Pabinger I, Brenner B, Kalina U, Knaub S, Nagy A, Ostermann H. Prothrombin complex concentrate (Beriplex P/N) for emergency anticoagulation reversal: a prospective multinational clinical trial. *J Thromb Haemost.* 2008;6:622-31.
38. Hu TY, Vaidya VR, Asirvatham SJ. Reversing anticoagulant effects of novel oral anticoagulants: role of ciraparantag, andexanet alfa, and idarucizumab. *Vasc Health Risk Manag.* 2016;12:35-44.

Conflict of Interest

The authors declare no conflict of interest.