

Influence of pressure pre-treatments on liquid whole egg thermal pasteurization – microbiological, physicochemical and functional properties



Ana C. Ribeiro^a, Francisco J. Barba^b, Xavier Barber^c, J.A. Lopes da Silva^a, Jorge A. Saraiva^{a,*}

^a Department of Chemistry, Associate Laboratory LAQV-REQUIMTE, University of Aveiro, Aveiro 3810-193, Portugal

^b Faculty of Pharmacy, University of Valencia, Burjassot, Valencia 46100, Spain

^c Departamento de Estadística, Centro de Investigación Operativa, Universidad Miguel Hernández de Elche-Alicante, Elche, Alicante, Spain

ARTICLE INFO

Keywords:

Liquid whole egg
Pressure treatment
Pasteurization
Salmonella Senftenberg 775/W
Functionality

ABSTRACT

As a possible alternative to commercial thermal pasteurization¹ (TP, 60 °C/3.5 min) of liquid whole egg² (LWE), a pasteurization based on the application of pressure pre-treatments (50 – 250 MPa/5 min) before a shorter TP (60 °C/1.75 min) was investigated. These combined treatments inactivated 3.35 to at least 6.09 log₁₀ cycles of *S. Senftenberg* 775/W, achieving comparable to greater inactivation as commercial TP (at pressures ≥ 200 MPa).

The treated samples presented lower soluble protein and emulsifying activity than non-treated LWE, yet higher viscosity and improved foaming capacity. In contrast to commercial TP, the LWE treated by moderate pressure³ (MP) before a shorter TP (MP-TP) showed higher soluble protein (7%) and viscosity (49%), better emulsifying properties (27 – 67%), and lower total carotenoids content (9%). Concerning sensory analysis, egg tarts prepared with non-treated and treated LWE obtained similar sensory acceptability.

Therefore, the heat sensitization effect induced by pressure pre-treatments on the *S. Senftenberg* 775/W population allows the reduction of TP time, a lethality effect equivalent to commercial TP and similar or improved functionality, being a very promising alternative to produce safe and better functional quality LWE.

1. Introduction

Eggs are considered by World Health Organization as a standard food protein of high biological value (one egg provides ≈ 14% of the recommended protein daily intake) (Food and Agriculture Organization, 2015; Franco et al., 2020). Yet, eggs are an excellent substrate for spoilage food-borne pathogens, and even stored under refrigeration are a highly perishable product (De Souza et al., 2015). *Salmonella* spp. and *Listeria monocytogenes* are pathogens of public health concern in liquid whole egg (LWE), hence thermal pasteurization (TP) is necessary to maintain product safety (European Commission, 2007). Commercially, LWE is pasteurized at 60 – 70 °C for 1.5 – 4.5 min (Food and Drug Administration, 2002; Monfort et al., 2012), nevertheless, this has a negative impact on egg products characteristics (De Souza et al., 2015; De Souza & Fernández, 2013; Herald & Smith, 1989).

In response to consumer demands for natural and minimally processed food products, and in an attempt to overcome the detrimental ef-

fects on eggs quality of conventional TP, the combination of non-thermal technologies such as high pressure (HP), pulsed electric fields (PEF), or ultrasound with a less intense TP have attracted increasing interest, as they may improve food safety and reduce losses of nutritional and sensory quality (Rahman, 2015). This possibility has been confirmed by the few published studies, in which microbial cells underwent sublethal damage during HP (200 – 300 MPa/3 – 30 min) or PEF (25 kV/cm/48 – 250 μs) treatments, thereby becoming more sensitive to subsequent TP (52 °C/3.5 min and 55 °C/2 min), with observed synergistic effects. However, these treatments only reduced about 1.5 – 3.5 log₁₀ cycles of *Salmonella* spp. and *Escherichia coli* K12 DH 5α (*Salmonella* Enteritidis surrogate) in LWE, which is comparatively lower than that achieved by commercial TP at 60 °C/3.5 min and 64 °C/2.5 min (5 – 9 log₁₀ cycles reductions of the most frequent *Salmonella* serotypes) (Jin et al., 2009; Monfort et al., 2011, 2012). So, in an effort to increase microbial lethality, Monfort et al. (2011) and (2012) combined the previous treatments with additives (triethyl citrate (TC) and EDTA), resulting in

* Corresponding author.

E-mail address: jorgesaraiva@ua.pt (J.A. Saraiva).

¹ Thermal pasteurization (TP)

² Liquid whole egg (LWE)

³ Moderate pressure (MP)

a reduction more than 5 log₁₀ cycles of *S. Enteritidis* and *E. coli* K12 DH 5α. Nevertheless, changes observed in the treated samples (reduced soluble protein, increased viscosity, and improved foaming properties) were mainly attributed to the presence of the additives (Monfort et al., 2012, 2012).

Therefore, in the search for processing methodologies that could lead to improved pasteurized LWE, a combined process using a pressure pre-treatment at 50 – 250 MPa/5 min before a TP at the commercial pasteurization temperature (60 °C), but with a shorter pasteurization time (3 min) was investigated. The purpose of this work was to optimize the conditions of pressure pre-treatments followed by a shorter TP in LWE, using *Salmonella enterica* serovar Senftenberg (ATCC 43845) (referred throughout this paper as *S. Senftenberg* 775/W) as a reference, due to its greater heat resistance compared to the serovars associated with most human infections (*S. Typhimurium* and *S. Enteritidis*) (European Food Safety Authority & European Centre for Disease Prevention and Control, 2021). In order to study this possibility in more detail, the impact on LWE quality properties, as well as the development of two microorganisms of interest in LWE (gram-positive and gram-negative bacteria), under post-TP refrigeration storage were also assessed.

2. Materials and methods

2.1. Sample preparation

Fresh shell eggs were acquired from a local supermarket the day before they were used and held overnight under refrigeration. The eggshells were cleaned completely with 70% ethanol, left to dry at room temperature inside a laminar flow cabinet, then cracked aseptically and the whole egg was carefully passed through a colander, while the vitellin membrane was disrupted with a scalpel blade. The whole egg was collected in a sterile cup and gently homogenized for 20 min at 1250 rpm using a magnetic stirrer (MS-3000, Biosan, Latvia).

2.2. Microorganisms

The strains used in this study were *S. Senftenberg* 775/W (ATCC 43845) and *Listeria innocua* (ATCC 33090). One glass bead from a deep-frozen culture was transferred to 250 mL of sterile Tryptic Soy Broth (TSB, Merck, Germany) and incubated at 37 °C with rotational stirring (150 rpm) overnight (Incubating orbital shaker, VWR, Portugal). Thereafter, 1 mL of this culture was transferred to 250 mL of sterile TSB and incubated again at 37 °C with rotational stirring (150 rpm) for 12 h (until the stationary growth phase was reached). The liquid culture was allowed to settle at room temperature for 15 min, and glycerol was added to a final 10% (v/v) concentration. Then, the samples were manually homogenized, transferred to Eppendorf tubes, immediately frozen under liquid nitrogen and stored at – 80 °C for later use.

2.3. Samples inoculation and packaging

The microorganism suspension prepared before (Section 2.2) was thawed at room temperature and used immediately afterwards. This suspension was added to LWE and stirred for 5 min at 1250 rpm with magnetic stirring (MS-3000, Biosan, Latvia). Then, non-inoculated and inoculated samples were packed under aseptic conditions in polyamide-polyethylene bags (Plásticos Macar Lda., Portugal), which had been previously sterilized with UV radiation, and manually thermally sealed to minimize the amount of air inside the bags. The initial concentrations of *S. Senftenberg* 775/W and *L. innocua* were approximately 10⁶ – 10⁷ CFU/mL.

2.4. Pressure treatments and thermal pasteurization

The pressure treatments were performed in a High-Pressure industrial equipment (Modelo 55, Hyperbaric, Spain), with a 55 L pressure

vessel at 50 – 250 MPa for 5 min at 20 °C, using water as the pressure-transmitting medium. For TP, the samples were immersed in a circulating water bath (Circulator Bath, FALC, Italy) at 60 °C/1.75 min (shorter TP) and 60 °C/3.5 min (commercial TP), with the temperature monitored by using a K-type thermocouple (Thermometer 305, Roline, Switzerland). The time started to count after a come-up time of 40 s (previously verified at the geometric centre of the bags (4×3×0.5 cm)). For the sequentially combined treatments, samples that had been submitted to pressure treatments were subsequently thermally treated, as described above. The maximum time between pressure treatments and TP was approximately 20 min. After treatments, the sample bags were immediately placed on ice and stored at 4 °C until analysis.

2.5. Microbiological analyses

The samples were analyzed in triplicated for *S. Senftenberg* 775/W and *L. innocua* counts, and the appropriate serial dilutions were prepared in Ringer's solution. The microorganisms were determined by the spread plate method, by plating 100 μL of the appropriate dilution, in duplicate, onto Xylose Lysine Deoxycholate Agar (XLD Agar, Himedia, India) (ISO 6579:2017) and *Listeria* Palcam Agar (Liofilchem, Italy) (ISO 11290:2017). The plates were incubated at 37 °C for 24 h and 48 h, for *S. Senftenberg* 775/W and *L. innocua*, respectively. Plates with 10 to 300 colonies were selected for counting and the results were expressed as log CFU/mL (microbial load). The results are presented as microbial log load variation (log (N/N₀)), which is calculated by the log load difference between the microbial load at each treatment (N) and the initial microbial load (without any treatment) (N₀).

2.6. Post thermal pasteurization refrigeration storage

LWE pasteurized by the optimized combined treatment was stored for 61 days at 4 °C and compared with commercial TP, to study the microbial development of *S. Senftenberg* 775/W and *L. innocua*. For this purpose, microbiological analysis was conducted before and after treatments, as well as on days 2, 8, 14, 22, 30, 42 and 61 of storage. The experiment was carried out in 3 replicates at each time point.

2.7. Physicochemical and functional analyses

2.7.1. pH and color

The pH was measured at 20 °C by directly submerging a calibrated glass electrode (pH electrode 50 14, Crison Instruments, S. A., Spain) in homogenized egg samples.

Color of LWE samples was analyzed using a Konica Minolta CM 2300d spectrophotometer (Konica Minolta, Japan). The color parameters L* - lightness (0, dark; 100, light), a* - redness (+, red; - green) and b* - yellowness (+, yellow; - blue) were obtained at room temperature with the CIELab system. The CIELab parameters were determined using the original SpectraMagic™ NX Software (Konica Minolta, Japan), according to the International Commission on Illumination regulations. For measurements, 6 mL of sample was homogenized and placed in a sample container. The total color difference (ΔE*) was calculated by Eq. (1) (Koç et al., 2011):

$$\Delta E^* = \left[(L^* - L_0^*)^2 + (a^* - a_0^*)^2 + (b^* - b_0^*)^2 \right]^{1/2} \quad (1)$$

where ΔE* is the total color difference between the sample and the control (raw sample), L* and L₀* are the lightness of treated and raw samples, respectively; a* and a₀* are the redness of treated and raw samples, respectively; and b* and b₀* are the yellowness of treated and raw samples, respectively.

2.7.2. Soluble protein content

The LWE samples were diluted at 10% (m/v) in distilled water, and then centrifuged at 10,000 g for 15 min at 4 °C (Heraeus Biofuge Stratos,

Thermo Electron Corporation, USA) (Sheng et al., 2018). The microassay Bradford method was used to determine the soluble protein content in the supernatant (Kruger, 2002). A standard curve was obtained using bovine serum albumin (Sigma-Aldrich, Portugal) in the range of 0.02 – 0.40 g/L. Measurements were performed in triplicate, and the results are expressed in wt.% (g of protein/100 g of whole egg).

2.7.3. Viscosity

Viscosity was measured using a rotational rheometer (Kinexus PRO, Malvern Analytical, Malvern, United Kingdom) with an attached cone-and-plate geometry (stainless steel cone, 4 ° and 40 mm diameter). The experimental temperature was controlled at 20.0 ± 0.1 °C by a Peltier system at the bottom plate. After transferring the sample to the rheometer bottom plate and allowing for a 5 min equilibration time, the sample was subjected to a 3 min increasing shear rate ramp from 0.1 to 100 s^{-1} , followed by a 3 min decreasing ramp, from 100 to 0.1 s^{-1} . Determination of the apparent viscosity and analysis of the flow curves were performed using the rSpace software (Malvern Analytical version 1.76s). Apparent viscosity measured at an intermediate shear rate of 50 s^{-1} was used for comparison among samples. Measurements were performed in triplicate of sample and duplicate of analysis.

2.7.4. Secondary lipid oxidation

Lipid oxidation was evaluated by quantification of secondary lipid oxidation products using the thiobarbituric acid-reactive substances (TBARS) method, as described by Vyncke (1970) with slight modifications. Two grams of egg were homogenized with 2 mL of 7.5% trichloroacetic acid, and the resulting suspension was centrifuged at 6000 rpm during 15 min at 4 °C (Centurion Scientific Ltd, Scansci, Portugal). 1 mL of the supernatant was added to 1 mL of 2-thiobarbituric acid (20 mM TBA in 99% acetic acid glacial) and the mixture was gently shaken. The mixture was immersed in a 100 °C water bath for 40 min before being cooled on ice. The absorbance was measured at 538 nm (Multiskan GO Microplate Spectrophotometer, Thermo Fisher Scientific Inc., USA). A standard curve of malondialdehyde (MDA) was obtained with 1,1,3,3-tetramethoxypropane, in the range of 0.5 – 10 µM MDA. The measurements were carried out in triplicate, and the results were expressed as µg MDA/100 g of whole egg.

2.7.5. Total carotenoids content

Total carotenoids were quantified based on the method described by Nagata and Yamashita (1992). Briefly, 1.0 g of LWE was dissolved in 20 mL of a (4:6, v/v) acetone-hexane mixture, and the absorbance was measured at 453, 505, 645 and 663 nm (Multiskan GO Microplate Spectrophotometer, Thermo Fisher Scientific Inc., USA). The calculations were conducted using the formulas given by the mentioned authors [24]. The measurements were performed in triplicate of sample and duplicate of analysis, and the results were expressed in mg/100 g of whole egg.

2.7.6. Foaming properties

Foaming properties were measured according to the method used by Sheng et al. (2018). Foams were produced using 7 mL whole egg sample by whipping for 2.5 min with a laboratory homogenizer (Ultra-Turrax, T25 basic, IKA®-Werke, Germany) at a speed of 9500 rpm and room temperature in a 20 mL graduated cylinder. Foaming capacity (FC) was calculated as percentage of increase in foam volume using the Eq. (2):

$$FC (\%) = \frac{V_1}{V_0} \times 100 \quad (2)$$

The foaming stability (FS) was measured by loss of foam volume after 30 min of standing at room temperature. FS was calculated by Eq. (3):

$$FS (\%) = \frac{V_2}{V_1} \times 100 \quad (3)$$

where V_0 is the initial volume of egg (mL); V_1 is the foam volume after whipping (mL); V_2 is the foam volume after 30 min (mL). The foaming

properties was evaluated in triplicate of sample and duplicate of analysis.

2.7.7. Emulsifying properties

As previously described by Zhang et al. (2019), the emulsifying activity index (EAI) and emulsion stability index (ESI) were measured at room temperature. To prepare emulsions, 3 mL of LWE were stirred by a homogenizer (Ultra-Turrax, T25 basic, IKA®-Werke, Germany) with 2 mL of sunflower oil, at 9500 rpm for 90 s. An aliquot of emulsion was collected from the bottom of the homogenized emulsion immediately (0 min) or 10 min after homogenization and diluted 300 times in a 0.1% (w/v) SDS solution. After dilution, the emulsion was shaken and the absorbance was read at 500 nm using a UV-Vis spectrophotometer (UVmini-1240, Shimadzu, Japan). Measurements were performed in triplicate of sample and duplicate of analysis and the EAI and ESI were calculated by Eqs. (4) and (5), respectively (Zhang et al., 2019):

$$EAI \left(\frac{m^2}{g} \right) = \frac{2 \times 2.303 \times A_0 \times DF}{c \times \Phi \times 10^4} \quad (4)$$

$$ESI \text{ (min)} = \frac{A_0}{(A_0 - A_{10}) \times \Delta t} \quad (5)$$

where A_0 and A_{10} are the absorbance of the diluted emulsions at 0 and 10 min, respectively, DF is the dilution factor, c is the initial concentration of protein (g/mL); Φ is the fraction of oil used to form the emulsion (0.4), and Δt is the time interval between taking the first and second aliquots of emulsion.

2.7.8. Gelling properties

Heating set gels were prepared by poring each LWE sample into a plastic casing, sealed at the both ends and heated at 85 °C for 35 min in a programmable circulating water bath (FALC, Italy). The LWE gels were quickly cooled in a water-ice bath and stored at 4 °C for 18 h. Before texture analysis, the gels were brought to room temperature, removed from the plastic and cut with a wire knife into cylinders of 22 mm diameter × 19 mm height. Textural analysis was conducted using a TA.XT.plusC Texture Analyzer (Stable Micro Systems, United Kingdom) equipped with a 30 kg load cell and a 50 mm diameter cylinder aluminium probe (P/50). The uniaxial compression tests up to rupture were performed at 2.00 mm/s. The Exponent Connect software (version 8.0.3.0, Stable Micro Systems, United Kingdom) was used to construct force-time curves and to obtain the following mechanical parameters: maximum force (N), area under the curve (N.sec) and slope at 10% height. For the analysis, at least six cylindrical gels of each treatment were used.

The water holding capacity (WHC) of the LWE gels was determined as reported by Marco-Molés et al. (2011), with some modifications. Centrifuge tubes filled with LWE (2.0 mL) were placed in a water bath (Circulator Bath, FALC, Italy) at 85 °C for 35 min for gel formation. After heating, the samples were immediately placed on ice during 15 min and then, kept at room temperature (20 ± 1 °C) for 1 h. The tubes were then centrifuged at 740 g for 10 min at 20 °C (Centurion Scientific Ltd, Scansci, Portugal). The WHC was given by Eq. (6):

$$WHC (\%) = \frac{WAC}{WBC} \times 100 \quad (6)$$

where WAC is the weight of gel after centrifugation and WBC is the weight of gel before centrifugation.

2.8. Sensory analyses

Sensory analysis was carried out for egg tarts prepared with non-treated and treated LWE. The ingredients (LWE and salt) were mixed with a hand mixer for 30 s, and then pored into 8-mold silicone pie trays (each one with 7 cm in diameter at the top and 5 cm in diameter at the bottom). The samples were baked at 190 °C for 12 min in a pre-heated oven (Flama, Cesar, Portugal). The egg tarts were stored at ambient temperature for cooling before the sensory evaluation. A panel of




Fig. 1. Log₁₀ cycles reductions of *Salmonella Senftenberg* 775/W (ATCC 43845) in liquid whole egg treated by pressure treatments at 50 – 250 MPa during 5 min. N denotes the microbial load measured for each treatment and N₀ the initial microbial load (without any treatment). Different letters indicate significant differences ($p < 0.05$) between treatments.

13 untrained volunteers from the Chemistry Department at the University of Aveiro, Aveiro, Portugal (5 females and 8 males aged between 24 and 54 years) evaluated the sensory characteristics of egg tart. Each egg tart sample was randomly presented to the panellists in plastic dishes labeled with randomized three-digit codes. The panellists were served with water and crackers and instructed to cleanse their palate between tastings. Panellists were asked to evaluate each sample regarding acceptability based on 8 criteria (visual appearance, surface color, interior color, aroma, texture when cutting, texture when cracking, overall sensation when chewing, flavour) using a 9-point hedonic scale (1 = 'dislike extremely', 9 = 'like extremely'). In addition, panellists were asked to order the samples according to their preference, from the one they liked least to the one they liked the most.

2.9. Statistical analyses

The data were tested at a 0.05 probability level ($p < 0.05$) and the effect of each condition was tested with a one-way analysis of variance (ANOVA), followed by a multiple comparisons test (Tukey's Honestly Significant Difference, HSD) to identify statistically significant differences between treatments (SPSS Statistics v.28).

3. Results and discussion

3.1. Microbial analyses: pressure treatments

Fig. 1 presents the evaluation of the inactivation of *S. Senftenberg* 775/W using moderate pressure (MP, 50 – 250 MPa/5 min) and high pressure (225 – 250 MPa/5 min) treatments. In general, as expected, an improvement in *S. Senftenberg* 775/W inactivation was obtained (the initial load was $6.94 \pm 0.08 \log_{10}$ CFU/mL) with increasing pressure, consistent with what other authors have reported (Bari et al., 2008; Monfort et al., 2012). MP treatments in the range of 50 – 90 MPa induced a decrease from 0.17 ± 0.11 to $0.39 \pm 0.16 \log_{10}$ cycles ($p \geq 0.05$), but at pressure ≥ 125 MPa a significantly higher inactivation was achieved ($0.53 \pm 0.09 - 2.46 \pm 0.18 \log_{10}$ cycles, $p < 0.05$). Nevertheless, other studies, despite applying similar or stronger conditions (200 – 250 MPa/5 – 30 min), obtained a lower inactivation (about 1 log₁₀ cycles) for *E. coli* K12 (*S. Enteritidis* surrogate in liquid egg) and *S. Enteritidis* PT4 E10 compared to our study, suggesting a greater resistance to pressure of these two microorganisms (İşiker et al., 2003; Monfort et al., 2012). Therefore, the pressure conditions tested were not sufficient to ensure the LWE safety, although, higher pressures were not considered, since treatments over 300 MPa may have a drastic impact on the physical properties of LWE (Monfort et al., 2012).

3.2. Microbial analyses: pressure treatment followed by thermal pasteurization

In this study, the use of pressure pre-treatments (50 – 250 MPa/5 min, at room temperature), aiming to cause sublethal damages in microorganisms and reduce their resistance to heat, before a shorter TP (60 °C/1.75 min) was evaluated to determine the lethal effect on *S. Senftenberg* 775/W, and compared to commercial TP (60 °C/3.5 min) (Fig. 2). When the MP pre-treatments were conducted at 50 and 70 MPa, about 3.4 log₁₀ cycles of the *S. Senftenberg* 775/W population were inactivated ($p \geq 0.05$), whereas at 90 MPa lethality improved slightly ($3.8 \pm 0.1 \log_{10}$ cycles, $p < 0.05$), and a similar lethal effect ($p \geq 0.05$) was obtained with increasing pressure up to 125 MPa. Moreover, the treatments carried out at pressures ≥ 200 MPa caused even greater inactivation ($p < 0.05$), resulting in reductions of at least 4.95 log₁₀ cycles, with counts reaching below the quantification limit ($\leq 2.00 \log$ CFU/mL) for 200 MPa and below the detection limit ($\leq 1.00 \log$ CFU/mL) for 225 and 250 MPa. These findings are more encouraging than those of previously published works, which reported reductions of 1.5 to 3.1 log₁₀ cycles for *E. coli* K12 DH 5 α (*S. Enteritidis* surrogate in liquid egg) in LWE treated at 200 – 300 MPa/3 – 30 min followed by TP at 52 – 55 °C/2 – 3.5 min (Monfort et al., 2012).

Otherwise, the shorter TP alone reduced *S. Senftenberg* 775/W counts by $3.01 \pm 0.12 \log_{10}$ cycles, but the commercial TP achieved a greater decrease of at least 4.95 log₁₀ cycles (counts below the quantification limit, $\leq 2.00 \log$ CFU/mL). Compared to the latter, the combined treatments induced a comparable (at 200 MPa, $p \geq 0.05$) or greater (at pressures ≥ 225 MPa, $p < 0.05$) inactivation. The synergistic effect observed at pressures ≥ 200 MPa suggest that *S. Senftenberg* 775/W was sublethally injured by pressure pre-treatments, possibly reducing their thermal resistance, and leading to the subsequent TP being more effective than when applied alone. In addition, the effects induced by pressure pre-treatments may be due to damage of the cell membrane and its structure, changes in mechanisms of nutrient absorption and release of metabolites; changes in proteins structure (denaturation, aggregation or gel formation) (Thakur & Nelson, 1998; Yaldagard et al., 2008), disruption of non-covalent bonds (mainly affecting tertiary and quaternary structures) (Knorr, 1999); and multimeric disintegration ribosomes (Niven et al., 1999). Thus, when this technology is applied at sublethal intensities, the damage caused will likely be to the same structures but to a lesser extent.

According to the results discussed previously, two combined treatments were chosen for further investigation: 200 MPa/5 min – 60 °C/1.75 min, which corresponded to a processing condition that accomplished a similar inactivation level to commercial TP, and did not cause a large increase in viscosity (as confirmed by a simple test of flow time




Fig. 2. Log₁₀ cycles reductions of *Salmonella* Senftenberg 775/W (ATCC 43845) in liquid whole egg treated by pressure pre-treatments (50 – 250 MPa/5 min) before a shorter thermal pasteurization (TP, at 60 °C/1.75 min) or TP only (60 °C/1.75 min and 60 °C/3.5 min). N denotes the microbial load measured for each treatment and N₀ the initial microbial load (without any treatment). The symbols # and * means that microbial counts were below the quantification limit (≤ 2.00 log CFU/mL) or detection limit (≤ 1.00 log CFU/mL), respectively. Different letters indicate significant differences ($p < 0.05$) between treatments.

along a glass surface at a 45° angle), and 50 MPa/5 min – 60 °C/1.75 min. Due to the intention of the authors to conduct additional studies involving the sequential combination of two nonthermal pre-treatments (pressure and ultrasound), in order to enhance microbial inactivation, the latter case was evaluate to determine if it would result in lower quality changes, despite having caused a lower microbial reduction. In addition, a post-TP refrigeration storage evolution of *S. Senftenberg* 775/W and *L. innocua* was examined, although only for the 200 MPa /5 min – 60 °C/1.75 min treatment, which achieved microbial inactivation comparable to that of commercial TP.

3.3. Microbial analyses: assessment of post thermal pasteurization refrigeration storage

The impact of MP-TP (200 MPa/5 min followed by 60 °C/1.75 min), shorter TP and commercial TP on *S. Senftenberg* 775/W and *L. innocua* (*L. monocytogenes* surrogate) counts during storage at 4 °C, was evaluated (Fig. 3). LWE was inoculated with an initial load of 6.62 ± 0.11 log CFU/mL of *S. Senftenberg* 775/W (Fig. 3a), and immediately after treatments (day 0), a reduction of 4.15 log₁₀ cycles was observed for shorter TP, and at least 4.62 log₁₀ cycles were obtained for MP-TP and commercial TP ($p < 0.05$). During storage, *S. Senftenberg* 775/W counts decreased in all samples, with a faster reduction in commercial TP- followed by MP-TP- and shorter TP-treated samples, with counts reaching below the detection limit (≤ 1.00 log CFU/mL) after 3, 8 and 42 days of storage, respectively. No further variation until the end of storage was observed.

Concerning *L. innocua* (Fig. 3b), the initial load was 6.99 ± 0.03 log CFU/mL, and the three treatments (day 0) reduced the counts by at least 4.99 log₁₀ cycles (below the quantification limit, ≤ 2.00 log CFU/mL). Regarding to commercial TP- and MP-TP-treated LWE, the loads did not change during 23 and 14 days (≤ 2.00 log CFU/mL), respectively, but then increased until day 42, with the increase being slightly more pronounced in MP-TP-treated samples (up to 4.92 ± 0.26 log CFU/mL) and remaining similar until the end of the experiment. A greater and faster *L. innocua* growth was observed in shorter TP-treated LWE, with levels of 5.5 ± 0.21 log CFU/mL after 30 days of storage and no significant variations until day 61. In addition, the prevalence and growth of *L. innocua*, during storage, suggests that, despite this microorganism counts being below the quantification limit after treatments, the remaining cells were able to multiply under refrigerated conditions, although to a less extent and later for commercial TP, followed by MP-TP and shorter TP.

3.4. pH and color

The pH values measured range from 7.72 ± 0.01 (raw LWE) to 7.68 ± 0.01 for treated samples (Table 1). Regarding LWE color, in general, treated samples showed similar lightness (L*), redness (a*) and yellowness (b*) ($p \geq 0.05$) to non-treated sample. However, the MP-TP (200 MPa)-treated LWE exhibited a decrease in yellow color (lower b* value, $p < 0.05$), resulting in a significantly higher ΔE^* , which might be indicate a lower carotenoids content (Monfort et al., 2012). Nonetheless, the color variation produced by the four treatments would be not detected by the naked eye ($\Delta E^* < 3$) (De Souza & Fernández, 2011). Similar results were reported by Monfort et al. (2012) for LWE treated with a PEF pre-treatment (25 kV/cm, 75 kJ/kg) followed by a TP (60 °C/1 min) in the presence of TC.

3.5. Soluble protein content

The soluble protein content represents the most practical criteria for predicting the overall functionality of LWE, and a reduction often reflects a decrease in egg proteins functionality (Herald & Smith, 1989; Monfort et al., 2012). As can be seen in Table 1, all treatments reduced significantly the soluble protein content of non-treated LWE (3.16 ± 0.11 to 2.78 ± 0.10 g/100 g, $p < 0.05$), showing a decrement of about 7% for MP-TP (50 and 200 MPa)- and shorter TP-treated LWE, while a slightly higher loss (12%, $p < 0.05$) was caused by commercial TP. Heat may trigger the unfolding of egg proteins by exposing the hydrophobic groups buried inside, hence promoting aggregation by hydrophobic interactions, and reducing soluble protein content (Chang et al., 2022). Moreover, the negative impact on MP-TP-treated samples might be attributed to the thermal effects, that resulted in a less pronounced unfolding compared to commercial TP. Our results are supported by the findings of De Souza and Fernández (2013), who observed that a mild LWE pasteurization (60 °C/3.5 min) can cause proteins unfolding and aggregation, possibly affecting ovomucoid, livetins, and some LDL apoproteins. Likewise with MP-TP-treated LWE, previous works indicated a soluble protein decreased (about 12%) after TP (52 °C/3.5 min) preceded by a pressure treatment (300 MPa/5 min) in presence of TC (Monfort et al., 2012).

3.6. Apparent viscosity

In accordance with previous authors (Monfort et al., 2012), the apparent viscosity of raw LWE was 13.0 ± 1.0 mPa.s (Table 1), but all treated LWE samples showed higher apparent viscosity (20.53 ± 1.41




Fig. 3. Development of a) *Salmonella Senftenberg 775/W* (ATCC 43845) and b) *Listeria innocua* (ATCC 33090) up to 61 days of storage after moderate pressure followed by a shorter thermal pasteurization (TP) (200 MPa/5 min – 60 °C/1.75 min) or TP only (60 °C/1.75 min and 60 °C/3.5 min). Unfilled symbols on graphics and # or * mean that microbial counts were below the quantification limit ($\leq 2.00 \text{ log CFU/mL}$) or detection limit ($\leq 1.00 \text{ log CFU/mL}$), respectively, and N_0 corresponds to the initial load before each treatment. Different letters denote significant differences ($p < 0.05$) between each processing condition and storage days.

Table 1

Physicochemical properties, lipid oxidation and total carotenoids of raw and treated liquid whole egg (mean \pm standard deviation). Different letters along each row denote significant differences ($p < 0.05$) between processing conditions.

Properties	Raw	Thermal Pasteurization		Combined Treatments	
		60 °C/3.5 min	60 °C/1.75 min	50 MPa/5 min – 60 °C/1.75 min	200 MPa/5 min – 60 °C/1.75 min
pH	7.72 \pm 0.01 ^a	7.69 \pm 0.01 ^{bc}	7.70 \pm 0.01 ^b	7.68 \pm 0.01 ^c	7.69 \pm 0.01 ^c
L*	47.16 \pm 0.16 ^{ab}	47.24 \pm 0.07 ^{ab}	46.89 \pm 0.67 ^{bc}	46.65 \pm 0.72 ^c	47.35 \pm 0.13 ^a
a*	7.37 \pm 0.12 ^{ab}	7.40 \pm 0.06 ^a	7.13 \pm 0.39 ^b	7.19 \pm 0.45 ^{ab}	7.41 \pm 0.09 ^a
b*	11.30 \pm 0.16 ^a	10.96 \pm 0.22 ^{ab}	10.57 \pm 0.73 ^b	10.55 \pm 0.74 ^b	10.62 \pm 0.16 ^b
ΔE^*	-	0.46 \pm 0.16 ^b	0.38 \pm 0.09 ^b	0.40 \pm 0.11 ^b	0.64 \pm 0.04 ^a
Soluble Protein (g/100 g whole egg)	3.16 \pm 0.11 ^a	2.78 \pm 0.09 ^c	2.93 \pm 0.10 ^b	2.94 \pm 0.17 ^b	2.95 \pm 0.19 ^b
Viscosity (mPa.s, at shear rate of 50 s ⁻¹)	12.98 \pm 1.03 ^c	20.62 \pm 1.33 ^b	20.53 \pm 1.41 ^b	28.64 \pm 2.26 ^a	30.64 \pm 4.17 ^a
Lipid Oxidation (TBARS, $\mu\text{g}/100 \text{ g}$ whole egg)	13.29 \pm 0.56 ^a	13.11 \pm 0.51 ^a	12.92 \pm 0.58 ^a	13.15 \pm 0.73 ^a	13.36 \pm 0.57 ^a
Total Carotenoids (mg/100 g whole egg)	0.92 \pm 0.02 ^a	0.90 \pm 0.03 ^a	0.94 \pm 0.01 ^a	0.93 \pm 0.02 ^a	0.81 \pm 0.04 ^b

– 30.64 ± 4.17 mPa.s, $p < 0.05$). The increment found for TP-treated LWE (about 59%) is in line with the soluble protein decrease (Table 1), probably due to protein unfolding and consequent aggregation of unfolded proteins (Jaekel & Ternes, 2009). MP-TP (50 and 200 MPa)-treated LWE showed an increase up to 49% and 136% higher ($p < 0.05$) than TP- and non-treated LWE, respectively. However, despite the soluble protein data did not show a significant difference between shorter TP- and MP-TP-treated samples, the higher impact caused by MP-TP on viscosity, may indicate more marked protein aggregation under these conditions. Consequently, the results suggest that, in addition to thermal effects, the MP pre-treatment might also have an impact on viscosity *per se*. In fact, according to previous studies, increasing pressure and treatment time resulted in a LWE viscosity increment, possibly due to a greater protein-protein interaction under pressure (Anson & Mirsky, 1931; Monfort et al., 2012).

3.7. Secondary lipid oxidation

Secondary lipid oxidation (TBARS, expressed as MDA content) was assessed (Table 1). The MDA content of raw LWE was 13.29 ± 0.56 $\mu\text{g}/100$ g, being comparable to values reported by other authors (Hur et al., 2003). No significant differences were noticed among raw and treated LWE, with values ranging from 12.92 ± 0.58 to 13.36 ± 0.57 μg MDA/100 g ($p \geq 0.05$). These findings are consistent with what was reported in a previous study (De Souza & Fernández, 2011), with no significant effect on TBARS values being observed for pasteurized LWE (60 °C/3.5 min).

3.8. Total carotenoids content

The non-treated LWE presented a total carotenoids content of 0.92 ± 0.02 mg/100 g (Table 1), which is consistent with values reported by Fredriksson et al. (2006). As shown in Table 1, for all treatments, no significant differences ($p \geq 0.05$) were observed compared to raw LWE, with exception of MP-TP (200 MPa) (a reduction of about 12%). The lack of effect noted for the TP is corroborated by published data when egg yolk (EY) was heated at 61.5 °C for 3.5 min (Wenzel et al., 2010). On the other hand, the significant loss observed for MP-TP-treated LWE at the highest pressure (~12%), may be attributed to degradation reactions that occur during processing (Cano et al., 2019), suggesting that the decrease was likely a result of the pressure pre-treatment. Indeed, published studies reported that MP treatment (200 MPa/6 min) reduced the total carotenoid content of persimmon fruit (Cano et al., 2019). Furthermore, the carotenoids reduction resulted in a less yellow color and a greater ΔE compared to the other treated LWE, however, these differences are not considered noticeable by the consumer as $\Delta E^* < 3$ (De Souza & Fernández, 2011).

3.9. Foaming properties

The foaming properties of LWE is attributed to the egg white (EW) proteins, while EY components are considered as an inhibitor due to the competition by yolk compounds (proteins and lipids) with EW proteins in the air bubble interfaces (Li et al., 2019). Table 2 summarizes the influence of the studied treatments on foaming properties of LWE. All treatments improved the FC of non-treated LWE (47 – 67%, $p < 0.05$), but had no impact on FS. Additionally, the treated LWE samples exhibited comparable foaming properties to each other ($p \geq 0.05$), and so, suggesting that the impact on MP-TP-treated LWE was related with the thermal effects. The improvement of FC could probably be associated with the conditions used for foam production. As reported above, the treatments led to the formation of aggregates by hydrophobic interactions, which are quite weak and can probably be dissociated during the homogenization used for foaming (9500 rpm/2.5 min), resulting in the formation of smaller aggregates. Consequently, these aggregates can

likely move faster to the foam interface, than their corresponding native proteins, thus improving FC (Guilmineau & Kulozik, 2006; Liang & Kristinnsson, 2007). Similar findings were described by Van der Plancken et al. (2007), when heated EW solutions (pH 7.6) at 60 °C. Actually, other works also reported no significant changes on FC of EW treated at 200 MPa (Yang et al., 2009) or EW solutions (pH 7.6) pressure-treated (0.1 – 700 MPa) at 60 °C (Van der Plancken et al., 2007).

3.10. Emulsifying properties

Egg emulsifying properties are mainly associated to yolk compounds, however the whole egg is also a good emulsifier (Hatta et al., 1997). As indicated in Table 2, the parameters used to describe emulsifying properties of LWE were EAI and ESI. For all treatments, in general, it was observed a significant reduction ($p < 0.05$) of EAI (11 – 38%) and ESI (22 – 53%) compared to non-treated LWE, however, a less pronounced effect was found for MP-TP-treated samples. This decrease might be attributed to the proteins unfolding, which promotes aggregation by hydrophobic interactions, and consequently, these aggregated proteins could not adsorb at the interface (Chang et al., 2022; Le Denmat et al., 1999). In opposition to what was suggested for foaming properties, the homogenization conditions used for emulsions production were less intense (9500 rpm/1.5 min) and were likely insufficient to allow the dissociation of aggregates formed with the treatments. Also, when less aggregates adsorb at the interface, the stability of the formed emulsions reduce, since this property is related to the interfacial area that can be coated by proteins (Pearce & Kinsella, 1978).

Further, LWE pasteurized at commercial conditions presented an ESI 1.6-fold lower than shorter TP-treated LWE ($p < 0.05$), showing that the pasteurization time had a significant effect on this parameter. On the contrary, Lechevalier et al. (2017) highlighted that pasteurization at 60 °C (maximum) optimized the emulsifying properties of LWE, and only at 66 °C the pasteurization time has a significant effect. Regarding MP-TP-treated samples, similar or higher emulsifying properties than shorter-TP-treated LWE was detected, consistently with Chen et al. (2019) work, who noticed an increase in EAI and ESI when using pressures up to 200 MPa, possibly indicating that in our work the thermal effects counteract the positive effect of MP pre-treatments. Additionally, enhanced emulsifying properties (1.3 – 1.7-fold higher) was found for MP-TP (200 MPa)-treated LWE compared to commercial TP LWE.

3.11. Texture and water holding capacity of LWE gels

The texture properties (maximum force, area under the curve, and slope at 10% of height) of thermal-induced LWE gels were evaluated by uniaxial compression tests. The typical force-time curves were characterized by a rise in force until the end of the test, and the obtained results are shown in Table 2.

The gels formed from raw LWE presented a significantly lower maximum force ($p < 0.05$) than those obtained from commercial TP- and MP-TP (200 MPa)-treated LWE (25 – 43% higher), and an increment of about 20% was found with the treatment time increase (1.75 to 3.5 min). As presented in Table 2, the area under the curve calculated for gels produced from raw and treated LWE showed non-significant differences among samples ($p \geq 0.05$), with the exception of gels produced with MP-TP (200 MPa)-TP-treated LWE, which presented a higher value ($p < 0.05$). Moreover, these last gels also demonstrated a significant increment ($p < 0.05$) in slope value relative to the other gels, although no significant changes ($p \geq 0.05$) were found between TP treated samples and raw LWE. Furthermore, the higher hardness and rigidity (15 – 43%) of gels from MP-TP (200 MPa)-treated LWE as compared to the gels treated by only temperature, suggest that the pre-treatment at the highest pressure was responsible for the observed increase. For MP-TP (50 MPa)-treated gels, although there were no significant differences compared to shorter TP and non-treated gels, an increasing trend was

Table 2

Functional properties of raw and treated liquid whole egg (mean \pm standard deviation). Different letters along each row denote significant differences ($p < 0.05$) between processing conditions.

Properties	Raw	Thermal Pasteurization		Combined Treatments	
		60 °C/3.5 min	60 °C/1.75 min	50 MPa/5 min – 60 °C/1.75 min	200 MPa/5 min – 60 °C/1.75 min
Foaming Capacity (%)	39.3 \pm 7.3 ^b	60.5 \pm 7.9 ^a	58.6 \pm 6.5 ^a	65.5 \pm 2.8 ^a	57.6 \pm 9.5 ^a
Foaming Stability (%)	82.9 \pm 8.1 ^a	70.3 \pm 6.2 ^a	72.1 \pm 5.7 ^a	73.7 \pm 10.6 ^a	72.0 \pm 9.6 ^a
Emulsifying Activity Index (m ² /g)	65.8 \pm 4.2 ^a	44.2 \pm 4.8 ^c	40.8 \pm 1.0 ^c	45.1 \pm 1.6 ^c	58.5 \pm 1.9 ^b
Emulsifying Stability Index (min)	1.3 \pm 0.3 ^b	0.6 \pm 0.0 ^d	1.0 \pm 0.2 ^c	1.5 \pm 0.2 ^a	1.0 \pm 0.0 ^{bc}
Maximum Force (N)	79.0 \pm 11.2 ^b	98.6 \pm 9.7 ^a	79.1 \pm 7.9 ^b	80.9 \pm 12.1 ^b	113.2 \pm 9.7 ^a
Area under the curve (N.sec)	639 \pm 31 ^b	689 \pm 51 ^b	631 \pm 32 ^b	688 \pm 51 ^b	800 \pm 37 ^a
Slope at 10% Height	0.62 \pm 0.01 ^{bc}	0.63 \pm 0.04 ^{bc}	0.61 \pm 0.01 ^c	0.66 \pm 0.02 ^b	0.70 \pm 0.03 ^a
Water Holding Capacity (%)	99.0 \pm 0.6 ^a	99.0 \pm 0.9 ^a	99.1 \pm 0.7 ^a	99.4 \pm 0.4 ^a	98.6 \pm 0.7 ^a




Fig. 4. Comparative diagram of sensory attributes of the egg tart prepared from raw and treated liquid whole egg (9-point hedonic scale: 1-disliked extremely, 9-liked extremely).

observed. Therefore, the findings indicate that gels produced from commercial TP more closely resembled gels produced from raw LWE than those produced from MP-TP (200 MPa)-treated LWE.

Concerning WHC, as can be seen in Table 2, non-significant differences ($p \geq 0.05$) were found among the WHC of gels produced from raw and treated LWE, as also observed by Monfort et al. (2012) and (2012) in LWE supplemented with 2% TC and treated by HP (300 MPa/5 min) or PEF (25 kV/cm) followed by TP (52 – 55 °C/2 - 3.5 min).

3.12. Sensory analysis

Sensory analysis of an egg-derived product (egg tart) prepared with non-treated and treated LWE (200 MPa/5 min followed by 60 °C/1.75 min, 60 °C/3.5 min) was assessed, and the results are shown in Fig. 4. On a 9-point hedonic scale, the range of attributes were from 5.2 to 7.3, which is equivalent to “neither like nor dislike slightly” and “like moderately”, respectively. The sensory characteristics scores (interior color, texture when cracking, overall feeling when chewing, flavour and global acceptability) were highest for non-treated samples, while the scores of visual appearance, surface color, aroma and texture when cutting were higher for the LWE thermally treated. Beside, the egg tarts prepared with MP-TP-treated LWE received the lowest scores for all sensory characteristics. For this sample, the lower score for texture when cracking is con-

sistent with the highest texture properties observed for thermal-induced gels. Yet, there were no significant differences ($p \geq 0.05$) between the three egg tarts. Egg tarts obtained a similar global acceptability by the panellists, although they were ordered by preference as follows: egg tart produced from MP-TP-treated LWE, raw LWE and commercial TP-treated LWE (from the one panellist liked least to the one they liked the most).

4. Conclusions

The application of pressure treatments alone (50 – 250 MPa/5 min) was ineffective in controlling *S. Senftenberg* 775/W, but when followed by a shorter TP, inactivation was significantly increased (at least 6.09 log₁₀ cycles), obtaining similar or even higher lethality than commercial TP, using a pressure pre-treatment ≥ 200 MPa. On the other hand, all the treatments caused protein unfolding and aggregation, resulting in a drop in soluble protein content and an increase in viscosity. Consequently, these changes in treated LWE resulted in enhanced FC (47 – 88%), higher texture properties (13 – 43%) and reduced emulsifying properties (18 – 53%) in comparison to non-treated LWE. Compared to commercial TP, the MP-TP treatment presented higher viscosity, emulsifying and gelling properties, equivalent foaming properties, and a reduced total carotenoids content. Additionally, egg tarts produced with

non-treated and treated LWE had a similar global acceptability. Hence, the findings hint the possibility to use sequentially combined treatments to enhance inactivation of *S. Senftenberg* 775/W, using a shorter TP and maintaining/improving some properties of pasteurized LWE.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

CRedit authorship contribution statement

Ana C. Ribeiro: Conceptualization, Investigation, Methodology, Formal analysis, Visualization, Writing – original draft. **Francisco J. Barba:** Writing – review & editing. **Xavier Barber:** Writing – review & editing. **J.A. Lopes da Silva:** Conceptualization, Methodology, Writing – review & editing, Supervision. **Jorge A. Saraiva:** Conceptualization, Methodology, Writing – review & editing, Supervision.

Data availability

Data will be made available on request.

Acknowledgments

This work received financial support from PT national funds FCT/MCTES, Fundação para a Ciência e Tecnologia and Ministério da Ciência, Tecnologia e Ensino Superior) through the financial support for the Associate Laboratory LAQV-REQUIMTE (UIDB/50006/2020 and UIDP/50006/2020) and Associate Laboratory LAQV-REQUIMTE (Laboratory of Bromatology and Hydrology, University of Porto) (Project AgriFood XXI (NORTE-01-0145-FEDER-000041)). Ana C. Ribeiro thanks FCT/MCTES (Fundação para a Ciência e Tecnologia and Ministério da Ciência, Tecnologia e Ensino Superior) and ESF (European Social Fund) through NORTE 2020 (Programa Operacional Região Norte) for her PhD grants ref. SFRH/BD/129540/2017 and COVID/BD/151876/2021.

References

- Anson, M. L., & Mirsky, A. E. (1931). The reversibility of protein coagulation. *Journal of Physical Chemistry*, 35(1), 185–193. [10.1021/j150319a015](https://doi.org/10.1021/j150319a015).
- Bari, M. L., Ukuku, D. O., Mori, M., Kawamoto, S., & Yamamoto, K. (2008). Effect of hydrostatic pressure pulsing on the inactivation of *Salmonella* Enteritidis in liquid whole egg. *Foodborne Pathogens and Disease*, 5(2), 175–182. [10.1089/fpd.2007.0055](https://doi.org/10.1089/fpd.2007.0055).
- Cano, M. P., Gómez-Maqueo, A., Fernández-López, R., Welti-Chanes, J., & García-Cayuela, T. (2019). Impact of high hydrostatic pressure and thermal treatment on the stability and bioaccessibility of carotenoid and carotenoid esters in astringent persimmon (*Diospyros kaki* Thunb. var. Rojo Brillante). *Food Research International*, 123, 538–549. [10.1016/j.foodres.2019.05.017](https://doi.org/10.1016/j.foodres.2019.05.017).
- Chang, K., Jiang, W., & Liu, J. (2022). Effect of subcritical water treatment on the structure and foaming properties of egg white protein. *Food Hydrocolloids*, 124(Part A), Article 107241. [10.1016/j.foodhyd.2021.107241](https://doi.org/10.1016/j.foodhyd.2021.107241).
- Chen, S., Wang, X., Xu, Y., Zhang, X., Wang, X., & Jiang, L. (2019). Effect of high pressure treatment on interfacial properties, structure and oxidative stability of soy protein isolate-stabilized emulsions. *Journal of Oleo Science*, 68(5), 409–418. [10.5650/jos.ess18228](https://doi.org/10.5650/jos.ess18228).
- De Souza, P. M., & Fernández, A. (2011). Effects of UV-C on physicochemical quality attributes and *Salmonella* Enteritidis inactivation in liquid egg products. *Food Control*, 22(8), 1385–1392. [10.1016/j.foodcont.2011.02.017](https://doi.org/10.1016/j.foodcont.2011.02.017).
- De Souza, P. M., & Fernández, A. (2013). Rheological properties and protein quality of UV-C processed liquid egg products. *Food Hydrocolloids*, 31(1), 127–134. [10.1016/j.foodhyd.2012.05.013](https://doi.org/10.1016/j.foodhyd.2012.05.013).
- De Souza, P. M., Müller, A., Beniaich, A., Mayer-Miebach, E., Oehlke, K., Stahl, M., Greiner, R., & Fernández, A. (2015). Functional properties and nutritional composition of liquid egg products treated in a coiled tube UV-C reactor. *Innovative Food Science and Emerging Technologies*, 32, 156–164. [10.1016/j.ifset.2015.09.004](https://doi.org/10.1016/j.ifset.2015.09.004).
- European Commission, (EC). (2007). Commission regulation (EC) No 1441/2007 of 5 December 2007 amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs. *Official Journal of the European Union*, L 322, 12–29.
- (2021). European food safety authority, (EFSA), & European centre for disease prevention and control, (ECDC). The European Union one health 2020 zoonoses report. *EFSA Journal*, 19(12), 1–324. [10.2903/j.efsa.2021.6971](https://doi.org/10.2903/j.efsa.2021.6971).
- Food and Agriculture Organization, (FAO). (2015). *Egg facts*. <http://www.fao.org/resources/infographics/infographics-details/en/c/284410/>.
- Food and Drug Administration, (FDA). (2002). Pasteurization requirements. *Code of Federal Regulations*, 9(590), 570.
- Franco, D., Rois, D., Arias, A., Justo, J. R., Marti-Quijal, F. J., Khubber, S., Barba, F. J., López-Pedrouso, M., & Lorenzo, J. M. (2020). Effect of breed and diet type on the freshness and quality of the eggs: A comparison between MOS (Indigenous Galician breed) and Isa brown hens. *Foods*, 9(3), 342–353. [10.3390/foods9030342](https://doi.org/10.3390/foods9030342).
- Fredriksson, S., Elwinger, K., & Pickova, J. (2006). Fatty acid and carotenoid composition of egg yolk as an effect of microalgae addition to feed formula for laying hens. *Food Chemistry*, 99(3), 530–537. [10.1016/j.foodchem.2005.08.018](https://doi.org/10.1016/j.foodchem.2005.08.018).
- Guilmineau, F., & Kulozik, U. (2006). Impact of a thermal treatment on the emulsifying properties of egg yolk. Part 2: Effect of the environmental conditions. *Food Hydrocolloids*, 20(8), 1114–1123. [10.1016/j.foodhyd.2005.12.006](https://doi.org/10.1016/j.foodhyd.2005.12.006).
- Hatta, H., Hagi, T., & Hirano, K. (1997). Chemical and physicochemical properties of hen eggs and their application in foods. In *Hen Eggs: Their Basic and Applied Science* (pp. 117–133). CRC Press. 1st ed..
- Herald, T. J., & Smith, D. M. (1989). Functional properties and composition of liquid whole egg proteins as influenced by pasteurization and frozen storage. *Poultry Science*, 68(11), 1461–1469. [10.3382/ps.0681461](https://doi.org/10.3382/ps.0681461).
- Hur, S. J., Kang, G. H., Jeong, J. Y., Yang, H. S., Ha, Y. L., Park, G. B., & Joo, S. T. (2003). Effect of dietary conjugated linoleic acid on lipid characteristics of egg yolk. *Asian-Australasian Journal of Animal Sciences*, 16(8), 1165–1170. [10.5713/ajas.2003.1165](https://doi.org/10.5713/ajas.2003.1165).
- Işiker, G., Gurakan, G. C., & Bayindirli, A. (2003). Combined effect of high hydrostatic pressure treatment and hydrogen peroxide on *Salmonella* Enteritidis in liquid whole egg. *European Food Research and Technology*, 217(3), 244–248. [10.1007/s00217-003-0759-3](https://doi.org/10.1007/s00217-003-0759-3).
- Jaekel, T., & Ternes, W. (2009). Changes in rheological behaviour and functional properties of hen's egg yolk induced by processing and fermentation with phospholipases. *International Journal of Food Science and Technology*, 44(3), 567–573. [10.1111/j.1365-2621.2008.01847.x](https://doi.org/10.1111/j.1365-2621.2008.01847.x).
- Jin, T., Zhang, H., Hermawan, N., & Dantzer, W. (2009). Effects of pH and temperature on inactivation of *Salmonella* Typhimurium DT104 in liquid whole egg by pulsed electric fields. *International Journal of Food Science & Technology*, 44(2), 367–372. [10.1111/j.1365-2621.2008.01759.x](https://doi.org/10.1111/j.1365-2621.2008.01759.x).
- Knorr, D. (1999). Novel approaches in food-processing technology: New technologies for preserving foods and modifying function. *Current Opinion in Biotechnology*, 10(5), 485–491. [10.1016/S0958-1669\(99\)00015-4](https://doi.org/10.1016/S0958-1669(99)00015-4).
- Koç, M., Koç, B., Susyal, G., Yilmazer, M. S., Ertekin, F. K., & Bağdatlıoğlu, N. (2011). Functional and physicochemical properties of whole egg powder: Effect of spray drying conditions. *Journal of Food Science and Technology*, 48(2), 141–149. [10.1007/s13197-010-0159-1](https://doi.org/10.1007/s13197-010-0159-1).
- Kruger, N. J. (2002). Quantification of proteins - The Bradford method for protein quantitation. In *The Protein Protocols Handbook* (pp. 15–21). Humana Press. 2nd ed.
- Le Denmat, M., Anton, M., & Gandemer, G. (1999). Protein denaturation and emulsifying properties of plasma and granules of egg yolk as related to heat treatment. *Journal of Food Science*, 64(2), 194–197. [10.1111/j.1365-2621.1999.tb15863.x](https://doi.org/10.1111/j.1365-2621.1999.tb15863.x).
- Lechevalier, V., Guérin-Dubiard, C., Anton, M., Beaumal, V., David Briand, E., Gillard, A., Le Gouar, Y., Musikaphun, N., Tanguy, G., Pasco, M., Dupont, D., & Nau, F. (2017). Pasteurisation of liquid whole egg: Optimal heat treatments in relation to its functional, nutritional and allergenic properties. *Journal of Food Engineering*, 195, 137–149. [10.1016/j.jfoodeng.2016.10.007](https://doi.org/10.1016/j.jfoodeng.2016.10.007).
- Li, X., Li, J., Chang, C., Wang, C., Zhang, M., Su, Y., & Yang, Y. (2019). Foaming characterization of fresh egg white proteins as a function of different proportions of egg yolk fractions. *Food Hydrocolloids*, 90, 118–125. [10.1016/j.foodhyd.2018.12.014](https://doi.org/10.1016/j.foodhyd.2018.12.014).
- Liang, Y., & Kristinsson, H. G. (2007). Structural and foaming properties of egg albumen subjected to different pH-treatments in the presence of calcium ions. *Food Research International*, 40(6), 668–678. [10.1016/j.foodres.2006.11.015](https://doi.org/10.1016/j.foodres.2006.11.015).
- Marco-Molés, R., Rojas-Graü, M. A., Hernando, I., Pérez-Munuera, I., Soliva-Fortuny, R., & Martín-Belloso, O. (2011). Physical and structural changes in liquid whole egg treated with high-intensity pulsed electric fields. *Journal of Food Science*, 76(2), 257–264. [10.1111/j.1750-3841.2010.02016.x](https://doi.org/10.1111/j.1750-3841.2010.02016.x).
- Monfort, S., Gayán, E., Condón, S., Raso, J., & Álvarez, I. (2011). Design of a combined process for the inactivation of *Salmonella* Enteritidis in liquid whole egg at 55 °C. *International Journal of Food Microbiology*, 145(2–3), 476–482. [10.1016/j.ijfoodmicro.2011.01.027](https://doi.org/10.1016/j.ijfoodmicro.2011.01.027).
- Monfort, S., Mañas, P., Condón, S., Raso, J., & Álvarez, I. (2012). Physicochemical and functional properties of liquid whole egg treated by the application of pulsed electric fields followed by heat in the presence of triethyl citrate. *Food Research International*, 48(2), 484–490. [10.1016/j.foodres.2012.04.015](https://doi.org/10.1016/j.foodres.2012.04.015).
- Monfort, S., Ramos, S., Meneses, N., Knorr, D., Raso, J., & Álvarez, I. (2012). Design and evaluation of a high hydrostatic pressure combined process for pasteurization of liquid whole egg. *Innovative Food Science and Emerging Technologies*, 14, 1–10. [10.1016/j.ifset.2012.01.004](https://doi.org/10.1016/j.ifset.2012.01.004).
- Nagata, M., & Yamashita, I. (1992). Simple method for simultaneous determination of chlorophyll and carotenoids in tomato fruit. *Journal of the Japanese Society for Food Science and Technology*, 39, 925–928.
- Niven, G. W., Miles, C. A., & Mackey, B. M. (1999). The effects of hydrostatic pressure on ribosome conformation in *Escherichia coli*: An *in vivo* study using differential scanning calorimetry. *Microbiology*, 145(2), 419–425. [10.1099/13500872-145-2-419](https://doi.org/10.1099/13500872-145-2-419).
- Pearce, K. N., & Kinsella, J. E. (1978). Emulsifying properties of proteins: Evaluation of a turbidimetric technique. *Journal of Agricultural and Food Chemistry*, 26(3), 716–723.
- Rahman, M. S (2015). Hurdle technology in food preservation. In *Minimally processed foods: technologies for safety, quality, and convenience* (pp. 17–33). Springer International Publishing. 1st ed.. [10.1007/978-3-319-10677-9](https://doi.org/10.1007/978-3-319-10677-9).
- Sheng, L., Wang, Y., Chen, J., Zou, J., Wang, Q., & Ma, M. (2018). Influence of high-intensity ultrasound on foaming and structural properties of egg white. *Food Research International*, 108, 604–610. [10.1016/j.foodres.2018.04.007](https://doi.org/10.1016/j.foodres.2018.04.007).

- Thakur, B. R., & Nelson, P. E. (1998). High-pressure processing and preservation of food. *Food Reviews International*, 14(4), 427–447. [10.1080/87559129809541171](https://doi.org/10.1080/87559129809541171).
- Van der Plancken, I., Van Loey, A., & Hendrickx, M. E. (2007). Foaming properties of egg white proteins affected by heat or high pressure treatment. *Journal of Food Engineering*, 78(4), 1410–1426. [10.1016/j.jfoodeng.2006.01.013](https://doi.org/10.1016/j.jfoodeng.2006.01.013).
- Vyncke, W. (1970). Direct determination of the thiobarbituric acid value in trichloroacetic acid extracts of fish as a measure of oxidative rancidity. *Fette, Seifen, Anstrichmittel*, 72(12), 1084–1087. [10.1002/lipi.19700721218](https://doi.org/10.1002/lipi.19700721218).
- Wenzel, M., Seuss-Baum, I., & Schlich, E. (2010). Influence of pasteurization, spray- and freeze-drying and storage on the carotenoid content in egg yolk. *Journal of Agricultural and Food Chemistry*, 58(3), 1726–1731. [10.1021/jf903488b](https://doi.org/10.1021/jf903488b).
- Yaldagard, M., Mortazavi, S. A., & Tabatabaie, F. (2008). The principles of ultra high pressure technology and its application in food processing/preservation: A review of microbiological and quality aspects. *African Journal of Biotechnology*, 7(16), 2739–2767. [10.5897/AJB07.923](https://doi.org/10.5897/AJB07.923).
- Yang, R. X., Li, W. Z., Zhu, C. Q., & Zhang, Q. (2009). Effects of ultra-high hydrostatic pressure on foaming and physical-chemistry properties of egg white. *Journal of Biomedical Science and Engineering*, 2(8), 617–620. [10.4236/jbise.2009.28089](https://doi.org/10.4236/jbise.2009.28089).
- Zhang, Q., Yang, L., Hu, S., Liu, X., & Duan, X. (2019). Consequences of ball-milling treatment on the physicochemical, rheological and emulsifying properties of egg phosvitin. *Food Hydrocolloids*, 95, 418–425. [10.1016/j.foodhyd.2019.04.064](https://doi.org/10.1016/j.foodhyd.2019.04.064).